

DIRECTIONS

WEEDS IN NON-CROP AREAS

	AMOUNTS TO USE	WHEN TO SPRAY
Lawns, Parks, Golf Courses, Cemeteries and Similar Turf Areas	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.
Fencerows, Road sides, Ditch banks and Vacant Lots	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.
Woody Plants* (vacant lots, fence rows, similar areas)	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.

WEEDS IN CROPPED AREAS

	AMOUNTS TO USE	WHEN TO SPRAY
Wheat* Barley and Rye	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.
Gram Sorghum	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.
Field Corn—post-emergence	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.
Hayfields, Pastures and Rangelands	1.0-4.0 qt 2.0-4.0 qt	Apply when weeds are in the early stages of growth. Spray after mowing, but before the next mowing. Do not mow for 24 hours after spraying. Do not apply to newly seeded lawns. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc. Do not apply to lawns containing desirable plants such as St. Augustine grass, Bahia grass, etc.

MIXING SPRAYS

For information on mixing sprays, see the label on the product container.

WATER HYACINTH CONTROL IN STILL WATER LAKES, PONDS, AND MARSHES

For information on water hyacinth control, see the label on the product container.

AERIAL APPLICATION

For information on aerial application, see the label on the product container.

BOAT APPLICATION

For information on boat application, see the label on the product container.

APPLICATION WITH LO DRIFT™ SPRAY ADDITIVE

For information on application with Lo Drift Spray Additive, see the label on the product container.

EMULSAMI E-3 BROADLEAF HERBICIDE

The oil soluble amine
form of 2,4-D

CAUTION: Keep out of reach of children.
See other cautions on right side panel.

ACTIVE INGREDIENTS:

Dodecylamine salt of 2,4-dichlorophenoxyacetic acid
Tetradecylamine salt of 2,4-dichlorophenoxyacetic acid

INERT INGREDIENTS:

*Isomer specific by AOAC method No. 6 D01-5
*Total 2,4-dichlorophenoxyacetic acid equivalent
or 27 pounds per gallon

Made in U.S.A.

UNION CARBIDE AGRICULTURAL PRODUCTS CO.

AMBLER, PA. 19002

Clinton, IA ■ St. Joseph, MO ■ Fremont, CA

EPA Reg No 264-179AA

EPA E

LSAMINE[®]

ROADLEAF HERBICIDE

soluble amine
D

Keep out of reach of children.
See instructions on right side panel.

2,4-dichlorophenoxyacetic acid* 47.3
% of 2,4-dichlorophenoxyacetic acid* 11.9
41.7

OAC method No. 6 D01 5
2,4-dichlorophenoxyacetic acid equivalent 31.2% by weight
on.

Made in U.S.A.

AGRICULTURAL PRODUCTS COMPANY, INC.
AMBLER, PA. 19002
■ St. Joseph, MO. ■ Fremont, CA

9AA

EPA Est. 264-PA 1

GENERAL INFORMATION

EMULSAMINE E 3 broadleaf herbicide is a concentrated formulation of 2,4-D as the new oil soluble amine.

EMULSAMINE E 3 broadleaf herbicide combines the effectiveness of low volatile ester forms of 2,4-D with maximum safety from volatility damage. Especially useful in areas of high summer temperatures.

CONTROLS THESE WEEDS

Annual and Biennial Weeds

bull thistle	mustard (wild)
burdock	peppergrass
cocklebur	pigweed
coffeeweed	lettuce (wild)
French weed	radish (wild)
galinsoga	ragweed (common)
jimsonweed	Russian thistle
kochia	sowthistle
lambsquarters	(common)
morningglory	velvetleaf

Perennial Weeds

alkali mallow	nettles
aster	orange hawkweed
bindweed	plantains
blue lettuce	povertyweed
blueweed	ragweed
bur ragweed	sowthistle
Canada thistle	stinging nettles
chicory	toadflax
dandelion	wild-garlic
docks	wild onion
goldenrod	

CAUTION

Harmful if swallowed or absorbed by skin. Do not allow contact with skin or clothing. If contacted, wash exposed skin with plenty of soap and water.

Causes irritation of eyes and skin. Do not get in eyes. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention.

This product is toxic to fish. Keep out of lakes, streams or ponds except for water hyacinth control. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate any body of water by washing of spray equipment or disposal of waste. Do not graze dairy animals on treated areas within 7 days after treatment.

Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, and ornamentals as this product may cause injury. Coarse sprays are less likely to drift. Do not use the same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water used for irrigation or domestic purposes. Do not use in or near a greenhouse.

Do not reuse container. Destroy when empty.

Do not use this product for purposes other than those recommended on this label.

LIMITED WARRANTY AND DISCLAIMER

