

WEEDS SUPER D WEEDONE CONTROLS

buckhorn plantain
 bull thistle
 burdock
 butter cup
 butter print
 Canada thistle
 chickweeds
 chicory
 clevers
 cinquefoil
 cocklebur
 dandelion
 decks
 frenchweed
 galinsoga
 ground ivy
 hawkweed
 heal-all
 henbit
 jewel weed
 jimson weed
 knotweed
 lambquarter
 mallow
 morning glory
 mustard
 oxalis
 pennywort
 pepperglass
 pigweed
 plantain
 poverty weed
 purslane
 ragweed
 sheep sorrel
 smartweed
 sowthistle
 speedwell
 vervains
 wild aster
 wild carrot
 wild garlic
 wild lettuce
 wild onion
 wild radish
 yarrow
 and many other
 broadleaf weeds

CONTENTS ONE QUART

SUPER D WEEDONE

RECEIVED
 DEC 31 1977
 U.S. FEDERAL INSECTICIDE
 AND FUNGICIDE ACT
 ECONOMIC POISON REGISTRY
 NUMBER NO. 264-189-AA

**CAUTION: Keep out of reach of Children.
 See other cautions on back panel.**

LAWN WEED CONTROL

FOR SOUTHERN GRASSES

(Bent, St. Augustine, Centipede)

Use one-half the recommended rate.

Complete directions on back panel.

ACTIVE INGREDIENTS:
 Diethanol amine salt of 2,4-dichlorophenoxyacetic acid* 20.3%
 Diethanol amine salt of dicamba** 1.9%
INERT INGREDIENTS 77.8%
 *2,4-Dichlorophenoxyacetic acid equivalent 13.8% by weight or 1 1/4 pounds per gallon.
 **3,6-Dichloro-o-anisic acid equivalent 1.3% by weight or 1/8 pound per gallon



AMCHEM PRODUCTS, INC.

First Name in Herbicide Research
 AMBLER, PA • Clinton, Iowa • St. Joseph, Mo • Fremont, Calif
 SUBSIDIARY OF UNION CARBIDE CORP.
 EPA Reg. No. 264-189-AA EPA Est. No. 264-PA-1

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem, therefore, makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damage arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 4793 10-77 TC 86

DIRECTIONS FOR USE ON LAWNS

Apply Super D WEEDONE any time weeds are growing the air is calm, to avoid spray drift that might injure desirable plants.
 Newly seeded lawns may be treated after they have been established. To re-seed small areas in treated lawns, wait 4 weeks before spraying. A good rain or deep watering should occur during the treatment.
 A few weeds, such as henbit that grow during the winter and garlic and onion that emerge early in the year, should be controlled in the spring.

AMOUNTS TO USE IN AMCHEM METER-MISER

SUPER-D WEEDONE	ADD WATER TO MAKE
12 oz.	5 gallons
6 oz.	2 1/2 gallons
2.5 oz.	1 gallon

For difficult-to-control weeds such as sheep sorrel, chickweed, clover, oxalis or yarrow, repeat application in the spring at the rate as above.

HOSE ATTACHED SPRAYERS—LAWN MODELS

Use 3 tablespoons of Super D WEEDONE diluted in 1 gallon of water per gallon mark on the sprayer jar for each 500 square feet of lawn to be sprayed. Adjust water pressure so that spray stream is fine misting. Take usual precautions to prevent spray drift.

AMOUNTS TO USE IN KNAPSACK OR PRESSURE SPRAYERS

AREA SPRAYED	SUPER-D WEEDONE
200 sq. ft. (10' x 20')	1 tablespoon
3,200 sq. ft. (40' x 80')	1 cup (8 fl. oz.)
6,500 sq. ft. (65' x 100')	1 pint
13,000 sq. ft. (130' x 100')	1 quart

Where difficult-to-control weeds such as sheep sorrel, chickweed or branched knotweed are a problem, the application rate recommended above.

IMPORTANT

Do not use Super-D WEEDONE for controlling weeds in flower beds, shrubs or ornamental plantings, Lippia and other plants. This treatment may injure bentgrass, St. Augustine and newly seeded lawns. To control the weeds in these areas, use the rate recommended above.

No weed spraying should be attempted during periods of drought. Lawn is watered thoroughly several days before treatment.

A separate sprayer should be kept for weed control. Related chemicals are difficult to clean from equipment. Do not use this product within the drip line of desirable plants. Injury to these plants may result.

WEEDS IN FENCEROWS & SIMILAR AREAS

To control large areas of woody brush and vines use Super D WEEDONE BK-32.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, nose, mouth and clothing. Avoid spray drift to susceptible plants as this product is also a herbicide. (Coarse sprays are less effective than fine sprays.) Do not store in same spray equipment for other purposes. Do not store in insecticides or fungicides. Under very high temperatures this product may injure susceptible plants in the immediate vicinity. Contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy when empty. Do not use this product for purposes other than those indicated on the label.

CONTENTS ONE QUART

SUPER D WEEDONE

RECEIVED
Dec 21 1977
FEDERAL INSECTICIDE
ACT
ELECTRONIC PERSON REGISTRY
UNDER NO. 264-187

CAUTION: Keep out of reach of Children.
See other cautions on back panel.

LAWN WEED CONTROL

FOR SOUTHERN GRASSES

(Bent, St. Augustine, Centipede)

Use one-half the recommended rate.

Complete directions on back panel.

ACTIVE INGREDIENTS:
Triethylamine salt of 2,4-dichlorophenoxyacetic acid* 20.3%
Triethylamine salt of dicamba** 1.9%
INERT INGREDIENTS 77.8%
*chlorophenoxyacetic acid equivalent 13.8% by weight or 1 1/4 pounds per gallon.
**chloro-o-anisic acid equivalent 1.3% by weight or 1/8 pound per gallon.

AMCHEM PRODUCTS, INC.
First Name in Herbicide Research
AMBLER, PA ■ Clinton, Iowa ■ St. Joseph, Mo ■ Fremont, Calif
SUBSIDIARY OF UNION CARBIDE CORP.
EPA Reg. No. 264-169-AA EPA Est. No. 264-PA-1

WARRANTY
Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem, therefore, makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damage arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 4793 10-77 TC 86

DIRECTIONS FOR USE ON LAWNS
Apply Super D WEEDONE any time weeds are growing actively. Spray when the air is calm, to avoid spray drift that might injure desirable plants.
Newly seeded lawns may be treated after they have been mowed at least twice. To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period.
A few weeds, such as henbit that grow during the winter, as well as wild garlic and onion that emerge early in the year, should be sprayed in early spring.

AMOUNTS TO USE IN AMCHEM METER-MISER®

SUPER-D WEEDONE	ADD WATER TO MAKE	SQ. FEET COVERED
12 oz.	5 gallons	5,000
6 oz.	2 1/2 gallons	2,500
2.5 oz.	1 gallon	1,000

For difficult-to-control weeds such as sheep sorrel, branched knotweed, chickweed clover, oxalis or yarrow, repeat application in two weeks using same rate as above.

HOSE ATTACHED SPRAYERS—LAWN MODELS
Use 3 tablespoons of Super D WEEDONE diluted with water to the 3 to 5 gallon mark on the sprayer jar for each 500 square feet (25' x 20') of lawn area to be sprayed. Adjust water pressure so that spray streams about 10-15 feet with no misting. Take usual precautions to prevent spray drift.

AMOUNTS TO USE IN KNAPSACK OR PRESSURE SPRAYERS

AREA SPRAYED	SUPER-D WEEDONE	WATER
200 sq. ft. (10' x 20')	1 tablespoon	1 gallon
3,200 sq. ft. (40' x 80')	1 cup (8 fl. oz.)	10 gallons
6,500 sq. ft. (65' x 100')	1 pint	20 gallons
13,000 sq. ft. (130' x 100')	1 quart	30 gallons

Where difficult-to-control weeds such as sheep sorrel, oxalis, yarrow, clover, chickweed or branched knotweed are a problem, use double the concentration recommended above.

IMPORTANT
Do not use Super-D WEEDONE for controlling weeds in flower or vegetable beds, shrubs or ornamental plantings, Lippia and Dichondra lawns.
This treatment may injure bentgrass, St. Augustine grass, centipede grass, and newly seeded lawns. To control the weeds in these types of grasses use half the rate recommended above.
No weed spraying should be attempted during periods of drought, unless the lawn is watered thoroughly several days before treatment.
A separate sprayer should be kept for weed control because 2,4-D and related chemicals are difficult to clean from equipment.
Do not use this product within the drip line of desirable trees and shrubs as injury to these plants may result.

WEEDS IN FENCE ROWS & SIMILAR AREAS
To control large areas of woody brush and vines, ask your dealer for WEEDONE BK-32.

CAUTION
Harmful if swallowed. Avoid contact with skin, eyes, or clothing.
Avoid spray drift to susceptible plants as this product may injure tomatoes, beans, peas and ornamentals. (Coarse sprays are less likely to drift). Do not use same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Under very high temperature of use, vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation or domestic purposes. Do not use in a greenhouse.
Do not reuse container. Destroy when empty.
Do not use this product for purposes other than those recommended on this label.



87