
DIRECTIONS

WEEDONE GARDEN WEEDER kills weed seedlings as they sprout. It will maintain weed control for 6-8 weeks or more, depending on weather conditions.

Apply to bare soil immediately after weeding. It will not kill growing weeds. Do not disturb treated soil.

Apply directly from this shaker canister. The drawing shows the approximate number of particles desired per square inch. Contents will treat 250 square feet.



Established Plants

Perennial flowers Shrubs and trees

Spread on the soil immediately after weeding and cleaning up each bed in the spring. If no rain falls within 3 to 4 days after application, water the area lightly to activate the chemical.

Transplanted Plants

Annual flowers Vegetables—tomato, pepper Perennial flowers and shrubs

Prepare the bed and set out the transplants as usual. Allow about two weeks to overcome the "shock" of transplanting and start growing. Pull or hoe out any weeds that started to grow during this period. Spread evenly, keeping as much off the foliage as possible. Do not use on pansies.

Treat vegetables only once during the season. If necessary, flowers and shrubs may receive a second application later in the summer.

Seeded Vegetables

Lima Beans Pumpkin, squash Seedling asparagus

Prepare a good seed bed as usual, then plant the seeds. Spread the same day as planting or within a few days after planting, before weeds emerge. If no rain falls within 3 to 4 days after application, water the area lightly to activate the chemical and to insure adequate moisture for plant growth. Apply once during the season.

Do not use on seeded flowers prior to emergence, as many will be injured.

Annual Flowers, Perennial Flowers, Shrubs & Trees

arborvitae ash aster azalea celosia chrysanthemum	dahlia daylilly dogwood holly iris ivy gladioli	juniper marigold peony petunia pine privet	rhododendron rose snapdragon spiraea spruce yew zinnia
--	---	---	--

Weeds Controlled

foxtails	carpet weed	lambsquarter
crabgrass	seedling Johnsongrass	fall panic grass
pigweed	ragweed	purslane
chickweed	'smartweed	velvet leaf

Sprouts coming from roots of established perennial weeds such as horsenettle, hedge bindweed, nutgrass, quackgrass, and Johnsongrass are not controlled.

CAUTION

Avoid contact with skin, eyes, or clothing.

Avoid unintentional mixing with fertilizer, seeds, insecticides, fungicides, or other herbicides.

Do not reuse container. Destroy when empty.

Do not use this product for purposes other than those recommended on the label.

U. S. Patent No. 3,014,063

Form No. 2151 B -- 2-69PL

Printed in U.S.A.

GARDEN WEEDER

Prevents Weeds in Flower Gardens, Vegetables, Shrubs



DIRECTIONS

fact and explanate following

Fransplanted Plants

Agentina de Companya de Co Companya de Co

and the second of the second o

Service Vicability

The property of the property o

A compared to the second secon

- : **!!!**

, ·

WARRANTY


Amchem was and the proconditions very label directives is authorized from a bread

WEEDINE

CAUTION KEEP OUT OF REACH OF CHILDREN SEE OTHER CAUTIONS ELSEWHERE ON LABEL

MET WEIGHT

11/2 POUNDS





ACTIVE INGREDIENTS:

