

INSTRUCTIONS

Apply before weeds are 3 to 4 inches tall. For best results apply and remove. Treat one or two weeks later when reapplying to keep ground clear of broadleaf weeds and

Mix 9 gallons of Liquid AMIZINE in a minimum of 50 gallons of water. Use in nozzles as well as those in the suction hose of a sprayer. Use 8002 or larger fan type nozzle or its equivalent. For 1 gallon of AMIZINE per 1,000 sq. ft. in sufficient water for good

Controlled weeds start to turn white or brown within a week. All weeds growing actively are killed more rapidly than

Apply into the ground if no rain falls within a week after application. Does not stain metal, stone or concrete.

METER-MISER

Apply the spray mixture to the correct area. The spray pattern covers the entire width between the wheels. To keep wheel tracks about 3 inches.

Close valve before stopping sprayer.

WATER TO USE

Water Make	Area Covered Square Feet
gal	5000
gal	4000
gal	3000
gal	2000
gal	1000

No. 264 196

Printed in U.S.A.

ACCEPTED
MAY 7, 1971
AMCHEM



LIQUID AMIZINE
A General Weed Killer

For use on substations, storage yards, driveways, parking areas, baseball diamonds, similar areas.

CAUTION: Keep out of reach of children. See other cautions on right side of label.

ACTIVE INGREDIENTS:

Amitrole (3-amino-1,2,4-triazole)*	5.1%
Simazine (2-chloro-4,6-bis-(ethylamino)-s-triazine)**	10.2%
INERT INGREDIENTS:	84.7%

*0.44 pounds per gallon **0.88 pounds per gallon

44

44A

AMCHEM PRODUCTS, Inc.

AMBLER, PA • Clinton, Iowa • St. Joseph, Mo • Fremont, Calif

M...
SHAKE CON...
A. Where hose available. Pour contents into MISER tank. Use the 5 gallon nozzle provided with sprayer. Rock METER...
B. Where hose available, or tankful. Pour contents into 5 gallon bucket. Stir, then pour into sprayer. For preparation of mix chemical container according to instructions. Pour into M...
C. Double Spray In areas where desired, or in skipped stream spray at half-rate first from end...
IMPORTANT: Apply after mixing. Clean sprayer several times with water. Before using for other purposes, clean tank and wheels thoroughly. Some of the weeds killed are: bluegrass, Canada thistle, chrysanthemum, weed, crabgrass, dandelion, foxtail, kochia, lambsquarters, nightshade.



LIQUID TRIAZINE Weed Killer

lawns, storage yards, driveways,
baseball diamonds, similar areas.

Keep out of reach of children.
Directions on right side of label.

Triazine* 5.1 %
(ethylamino)-s-triazine)** 10.2 %
..... 84.7 %

*0.88 pounds per gallon

44A

AMCHEM PRODUCTS, Inc.

St. Joseph, Mo. • Fremont, Calif.

MIXING SPRAYS SHAKE CONTAINER BEFORE USING

A. Where hose water under pressure is available.

Pour contents of this container into METER-MISER tank. Add water under pressure to the 5 gallon mark (bottom of funnel provided with sprayer).

Rock METER-MISER back-and-forth several times, then spray.

B. Where hose water under pressure is not available, or for mixing less than a tankful.

Pour contents of this container into a 5-gallon bucket. Add water to make 5 gallons. Stir, then pour into METER-MISER tank.

For preparing less than 5 gallons of spray, mix chemical and water in a separate container according to table in directions panel. Pour into METER-MISER tank.

C. Double Spray Volume.

In areas where higher volume sprays are desired, or to minimize the possibility of skipped streaks, use double coverage. Mix spray at half-strength. Cover the area twice, first from end to end, then from side to side.

IMPORTANT: Apply spray as soon as possible after mixing.

Clean sprayer after use each day by flushing several times with clean water.

Before using for weed control in turf, spray out a tankful of detergent solution to thoroughly clean tank and sprayer head. Also, clean wheels thoroughly.

Some of the weeds controlled are:

bluegrass	pigweed
Canada thistle	quackgrass
chrysanthemum	ragweed
weed	smartweed
crabgrass	saw thistle
dandelion	plantain
foxtail	purslane
kochia	velvet grass
lambsquarters	wild oats
nightshade	witchgrass

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing.

Avoid spray drift to susceptible plants such as cotton, beans, tomatoes and ornamentals. (Coarse sprays are less likely to drift.) Do not use same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water used for irrigation or domestic purposes. Do not use in a greenhouse.

Do not reuse container. Rinse empty container thoroughly with water and discard or bury in a safe place. Do not burn.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

45

DIRECTIONS

Apply Liquid AMIZINE in spring or early summer, before weeds are 3 to 4 inches tall. For best results on larger weeds, cut or mow them to ground line and remove. Treat one or two weeks later when re-growth appears. Use Liquid AMIZINE every spring to keep ground clear of broadleaf weeds and grasses.

For conventional high volume ground sprayers, mix 9 gallons of Liquid AMIZINE in a minimum of 50 gallons of water for each acre to be sprayed. Screens in nozzles as well as those in the suction hose should be no finer than 50 mesh. Use a Tee-Jet 8002 or larger fan type nozzle or its equivalent. For knapsack sprayers, apply $\frac{1}{2}$ gallon of Liquid AMIZINE per 1,000 sq. ft. in sufficient water for good coverage; generally 2 to 3 gallons.

Liquid AMIZINE acts slowly but thoroughly. Treated weeds start to turn white or brown within a week and are usually dead in about three weeks. Small weeds growing actively are killed more rapidly than weeds in a dormant or dried condition.

In arid areas, Liquid AMIZINE should be watered into the ground if no rain falls within a week after treatment. Liquid AMIZINE is non-flammable and does not stain metal, stone or concrete.

USING THE AMCHEM METER-MISER*

The AMCHEM METER-MISER is calibrated to accurately apply the spray mixture to the correct area, when pushed at such a speed that the spray pattern covers the entire width between the wheels. To avoid skips, walk in straight lines and over-lap wheel tracks about 3 inches.

Open valve after sprayer is in motion. Close valve before stopping sprayer.

AMOUNTS TO USE

Liquid AMIZINE	Add Water To Make	Area Covered Square Feet
1 gal	5 gal	5000
$\frac{4}{5}$ gal	4 gal	4000
$\frac{3}{5}$ gal	3 gal	3000
$\frac{2}{5}$ gal	2 gal	2000
$\frac{1}{5}$ gal	1 gal	1000

Form No. 2899A 1-71 TC

USDA Reg. No. 264 196

Printed in U.S.A.

AC
MAY

A
A G

For use
parking

ACTIVE ING
Amitrole
Simazine
INERT ING

*0.44 pou

AM
AMBLER