

AMITROL-T.

CAUTION: Keep out of reach of children

See other cautions on left panel.

CONTROLS

**poison ivy
poison oak
weed grasses
in lawns**

This product affects all turf grasses.
Treat only grasses you
want to control.

CONTENTS 32 FL. OZ. (1 QT.)

| | |
|------------------------------------|-------|
| ACTIVE INGREDIENT | |
| Amitrole (3-amino 1,2,4 triazole)* | 21.6% |
| INERT INGREDIENTS | 78.4% |

*Two pounds of amitrole per gallon



A MCHM COMPANY



AMCHEM PRODUCTS, INC.

First Name in Herbicide Research
AMBLER, PA • Clinton, IA • St. Joseph, MO • Fremont, CA

EPA Reg No 264-226 AA

EPA Est. 264-PA-1

AMITROL-T contains amitrole, one of the most effective chemicals in a knife for controlling poison oak, poison ivy, Canada thistle, milkweed and certain other weeds and grasses. The product is non-volatile, is easy to use from spray equipment and does not sterilize soil when used as directed. This product effects all turf grasses. Treat only the areas you wish to control. If exposed to temperatures of 100° or below, shake container before using.

DIRECTIONS FOR USE

1. Fence rows, vacant lots, roadsides

Use a knapsack or similar pressure sprayer producing medium size droplets. Watering cans or fine sprayers are not recommended. Avoid drift by spraying when the air is quiet. Wet weed to apply thoroughly. A gallon of spray solution is usually enough to cover about 100 square feet. 25 x 10 inches infested with weeds.

POISON IVY, POISON OAK

Treat from the time leaves are fully sized in the spring until they begin to fall in the fall. Mix 1/4 fluid ounce (2 tablespoons) of this product in 1 1/2 gallons of water. Wet a patch of ivy or poison oak leaves and stems to a height of 3 to 6 feet. Poison ivy or poison oak growing under or on large shade trees can be treated if the spray is kept off the tree leaves. Never burn dead poison ivy or poison oak.

QUACKGRASS, TALL FESCUE, BERMUDAGRASS, OTHER WEED GRASSES, ALSO CANADA THISTLE, MILKWEED, HORSETAIL RUSH

Mix 1/4 cup (2 fluid ounces) of this product in 1 gallon of water. Spray quackgrass or other weedy grasses when top growth is 6 to 8 inches long.

When leaves become whitish, turn under, covering the grass completely with soil.

Spray Bermudagrass in spring when shoots are young and growing vigorously. Re-spray any green regrowth that appears.

Spray thistles between the bud and bloom stages. Do not mow treated plants. Spray milkweed in early summer during the pre-bloom stage. Spray horsetail when growing vigorously. If regrowth appears, spray again in August or the following spring.

2. Late summer to early fall LAWN RENOVATION of areas infested with quackgrass, tall fescue, muhlenbergia, orchardgrass or Bermudagrass

Soil must be moist and plants actively growing. Apply 1 pint of this product in 10 to 12 gallons of water per 2000 square feet (40' x 50'). If no rain falls within 5 to 7 days after application, water heavily. Ten to fourteen days after application, dig, rotovate or plow treated area. (Raking, verticutting or aerifying will not be satisfactory.) Prepare good seedbed and plant grass seed.

Avoid walking from treated to non-treated lawns if some of the chemical gets on your shoes. It may discolor desirable grasses.

In areas where late fall seeding is practical or a heavy stand of bermudagrass exists, a second application should be made for maximum control. Apply treatment to regrowth after it is 2-3" tall, rotovate again in 10 days and seed. Maintain good soil moisture between treatment and rotovation. Note: Rainfall or heavy watering within 12 hours after treatment will reduce weed control.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

Avoid spraying of desirable plants (shrubs, flowers, vegetables, turfgrasses) as injury may result. Spray drift may cause whitened new growth, but unless plants have been sprayed heavily, most recover.

After spraying, rinse equipment thoroughly with water, pumping it through the entire spray system. Do not store near fertilizers, seeds, insecticides or use in a greenhouse. Keep livestock off treated areas.

Do not reuse empty container. Dispose in a safe place. Do not burn.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 2585-E (7-71)

U.S. Patent No. 2,892,000

Cameraman's Note
Poor Copy

ACCEPTED
UNDER
FUNGICIDE
FOR ECONOMIC
CONTROL UNDER NO. 1
TO ATTACHED

264-226