


Proposed EMULSAVERT D Label

USDA Reg. No. 264-X

December 12, 1967



Contents

Gallons

Proposed EMULSAVERT D Label

USDA Reg. No. 264-X

December 12, 1967

Page 2

A M C H E M

EMULSAVERT D

An "Invert Emulsion" formulation of 2,4-D

Controls many annual, biennial and perennial broadleaf weeds on roadsides, fence rows and ditchbanks.

CAUTION: Keep out of reach of children. See other cautions on the right panel.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxy acetic acid*	9.1 %
N,N-dimethylethylamine salt of 2,4-dichlorophenoxy acetic acid*	34.6 %

INERT INGREDIENTS

56.3 %

*Total 2,4-dichlorophenoxy acetic acid equivalent 24.0 % by weight or 2 pounds per gallon

AMCHEM PRODUCTS, INC.

Ambler, Penna.

EMULSAVERT D makes a thicker, more viscous spray mixture than standard water, oil-water or oil mixtures. EMULSAVERT D is as effective as low volatile 2,4-D formulations without the problem of volatility at high temperatures.

DIRECTIONS

Add the required amount of EMULSAVERT D to the oil in the spray tank. Then add water while agitating the mixture vigorously until it becomes thick.

Always add water to the oil-concentrate mixture, never the reverse.

To Control:

Amounts per acre:

	EMULSAVERT D	Oil	Water	Total Volume/Acre
Annual and biennial broadleaf weeds	2 qts	4½ gal.	15 gal.	20 gal.
Perennial broadleaf weeds	4 qts	9 gal.	30 gal.	40 gal.

Treat annual and biennials such as ragweed, lambsquarter, dandelion and kochia when weeds are young and growing vigorously.

Treat perennials such as Canada Thistle, bindweed, stinging nettle and leafy spurge when weeds are actively growing and near the bud stage but before flowering. Most perennial weeds will need a repeat application. Treat when regrowth is 1 to 2 feet tall.

Do not apply to newly seeded turf areas until the seedlings are well established and have been mowed at least three times.

Use TOC 40, 60 or 80 nozzles (off center nozzles). Adjust spray swath, truck speed and spray pressure to provide the required volume of spray per acre. (Note: Nozzle capacity charts show nozzle delivery with water and will probably be different for invert emulsions). Since thickened sprays offer more resistance to flow than normal solutions, be sure sprayer hoses have sufficient capacity to deliver EMULSAVERT D to the nozzle without a great increase in sprayer pressure.

December 12, 1967

Page 3

CAUTION

Harmful if swallowed.

Avoid contact with skin, eyes or clothing.

Avoid spray drift to susceptible plants as this product may injure cotton, tobacco, beans, peas, ornamentals, etc. Course sprays are less likely to drift. Do not use the same spray equipment for other purposes.

Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water used for irrigation or domestic purposes. Do not use in a greenhouse.

Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands located away from water supplies. Do not reuse containers for other purposes. Destroy when empty.

Keep out of any body of water. Fish may be killed.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed.

Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.