


Proposed WEEDONE MCPA ESTER Label

Contents
Gallons

USDA Reg. No. 264-X

December 23, 1968



Anchem

WEEDONE MCPA

ESTER

Kills Broadleaf Weeds in Small Grains

CAUTION: Keep out of reach of children.

See other cautions on right panel.

ACTIVE INGREDIENTS:

Eutoxyethanol ester of 2-methyl-4-chlorophenoxyacetic acid* 66.7%

INERT INGREDIENTS: 33.3%

*2-methyl-4-chlorophenoxyacetic acid equivalent 44.5% by weight or 4.0 pounds per gallon.

Anchem Products, Inc.

Aabler, Pa. 19002

Proposed WEEDONE MCPA ESTER Label

USDA Reg. No. 264-X

December 23, 1968

Page 2

CAUTION

Harmful if swallowed. Avoid contact with skin eyes or clothing. In case of contact, flush eyes with plenty of water. Wash exposed skin with plenty of soap and water.

Avoid spray drift to susceptible plants as this product may injury cotton, beans, peas, grapes, ornamentals. Coarse sprays are less likely to drift. Do not use same spray equipment for other purposes unless thoroughly cleaned.

Do not reuse empty containers. Destroy containers by burying in a safe place. Do not burn.

Do not store near fertilizers, seeds, insecticides or fungicides. Under very high temperature, vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation or domestic purposes. Do not use in a green house.

Do not forage or graze treated grain fields within two weeks after treatment with this product. Do not feed treated straw to livestock.

Avoid contamination of lakes, streams or ponds as this product may kill fish.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Anchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Anchem therefore makes no other express or implied warranty, and no agent of Anchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Anchem's liability.

DIRECTIONS

Use the lower rate where susceptible weeds such as labasquarters, mustards, ragweed, stinkweed, wild radish, yellow rocket, purslane, peppergrass, shepherdspurse, henbane, and annual sow thistle are a problem. Use the higher rate where less susceptible weeds such as dandelion, plantain, pigweed, pennycress, kochia and perennial sow thistle are a problem.

| CROP | Amount per Acre | When to Spray |
|--|-----------------|--|
| SMALL GRAINS | | |
| Spring planted (not seeded to legumes) | ½ to 1½ pint | Wheat, <u>oats</u> , barley, rye Treat in spring when grain is fully tillered and 8 to 10 inches tall. Do not spray grain in the boot to dough stage. Use at least 8 to 10 gallons of water per acre. |
| Fall planted | ½ to 1½ pint | Apply in early spring when annual broadleaf weeds are small and grain is fully tillered, but before grain is in the jointed stage. Use at least 8 to 10 gallons of water per acre. |
| FLAX | 1/4 to 3/8 pint | Where susceptible weeds such as mustard are the main problem. |
| | ½ pint | Where weeds such as pigweed, labasquarters, stinkweed, ragweed, cocklebur are the problem. Apply when flax is 2 to 3 inches tall but before bud stage. Do not spray after flax is in the bud stage as severe damage may result. |
| OATS | ½ to 1 pint | Spray after oats are fully tillered (4 to 6 inches tall) but before they are in the boot to dough stage. Apply the lower rate where annual weeds are the major problem, the higher rate for perennial weeds. Use at least 8 to 10 gallons of water per acre. |
| RICE | 1½ to 2½ pints | Treat 7 to 10 weeks after planting when rice is fully tillered and 6 to 8 inches above water. Do not spray after rice is in the boot stage. Use the higher rate where curly indigo, sedge or bulrush are the major problems. Use lower rate for arrow head, water plantain and redstem. Do not spray when temperatures are over 90° F. |

Where MCPA resistant weeds are present in wheat and barley, the combination of WEEDONE MCPA ESTER and BROMINAL may be more effective. See special instructions for mixing with MCPA on the BROMINAL label.

The above recommendations are general. Because of variations in soil, moisture conditions, weed population, follow the recommendations of your extension weed control specialist.

To convert local recommendation into terms of WEEDONE MCPA ESTER, use the following table:

| MCPA acid | 1 1/4 lb. (20 oz.) | 1 lb. (16 oz.) | 3/4 lb. (12 oz.) | 1/2 lb. (8 oz.) | 3/8 lb. (6 oz.) | 1/4 lb. (4 oz.) |
|--------------------|-----------------------|-------------------|---------------------|--------------------|--------------------|--------------------|
| WEEDONE MCPA ESTER | 2 1/2 pt. | 2 pt. | 1 1/2 pt. | 1 pt. | 3/4 pt. | 1/2 pt. |

Important: Clovers and other legumes may be injured or killed by these rates. Do not use on susceptible turfgrasses such as St. Augustine, centepede, carpetgrass, bentgrass or ground covers such as dichondra.