DIRECTIONS

ORNAMENTAL PLANTINGS

ORNAMENTAL WEEDER controls germinating annual broadleaf weeds and grasses in ornamental plantings, ground covers and conifer seed beds. Preemergence application of ORNAMENTAL WEEDER prevents germination of these weeds for 6 to 8 weeks after treatment.

Apply ORNAMENTAL WEEDER granules to weed free ornamental plantings at the rate of 100 lbs. per acre. ORNAMENTAL WEEDER must be applied in the spring before weeds germinate, or to freshly cultivated soil. It will not control weeds which have already germinated or emerged. It will not control perennial weeds.

Do not incorporate ORNAMENTAL WEEDER into the soil.

If no rain falls within one week of application, apply ½ inch of overhead irrigation. This will wash the chemical into the soil where it kills germinating weeds.

ORNAMENTAL WEEDER may be used on the following:

	TIME FOR APPLICATION		
	Immediately after Transplanting	One month after transplanting	Established stock (6 wks or more)
arborvitae			X
azalea			X
[illegible]			X
chrysanthemum			X
[illegible] aster	X	X	X
[illegible]			X
[illegible]			X
[illegible]			X
[illegible]			X
English ivy	X	X	X
hemlock			X
holly	X	X	X
[illegible]			X
[illegible]			X
juniper	X	X	X
[illegible]			X
marigold			X
multiflora rose	X	X	X
[illegible]			X
[illegible]			X
[illegible]			X
red maple	X	X	X
[illegible] oak	X	X	X
[illegible]			X
[illegible]	X	X	X
[illegible]			X
[illegible]			X
[illegible] (Zinnia)			X
[illegible]			X
[illegible]	X	X	X
[illegible]	X	X	X
[illegible]			X

[illegible] ORNAMENTAL WEEDER on [illegible] eggplant [illegible] [illegible] the [illegible] soils.

CONTENTS 25 POUNDS

ORNAMENTAL WEEDER

A GRANULAR HERBICIDE for preemergence weed control in ornamental planting, ground covers and gladiolus

CAUTION: Keep out of reach of children.
See other cautions on right panel.

ACTIVE INGREDIENTS:

Ammonium salt of chloramben (3-amino-2,5-dichlorobenzoic acid)*	3.9%
Ammonium salts of related aminodichlorobenzoic acids*	0.4%
INERT INGREDIENTS:	95.7%

*Total aminodichlorobenzoic acid equivalent	4.0% by weight
Chloramben acid equivalent	3.6%
Related aminodichlorobenzoic acid equivalents	0.4%

A [illegible] AMERICAN CHEMICAL COMPANY

AMCHEM PRODUCTS, INC.
First Name in Herbicide Research
AMBLER, PA ■ Clinton, Iowa ■ St. Joseph, Mo. ■ Fremont, Calif.

EPA Reg. No. 264-243-7A EPA Est. 264-PA-1

DIRECTIONS

WEED CONTRO[illegible] CORMS AND C[illegible]

ORNAMENTAL [illegible] plied to gladiolu[illegible] cormel producti[illegible] the growth cycle.

FOR FLOWER [illegible]

two treatments

1. Apply 100 [illegible] WEEDER p[illegible] immediately [illegible] with bulbs [illegible]
2. At lay by, [illegible] (50-75 days [illegible] the same r[illegible] WEEDER a[illegible] application.

FOR FLOWER [illegible]

one treatment

Apply 150 [illegible] WEEDER per [illegible] either at planti[illegible]

FOR CORMEL F[illegible]

Cormels must [illegible] inches deep. F[illegible] weed free soil [illegible] lay by. Use 10[illegible] WEEDER per [illegible] incorporate.

Weeds Controlle[illegible] WEEDER Granul[illegible]

chickweed
coffeeweed *(Sesba[illegible]*
crabgrass
fall panicgrass
foxtails
kochia

DIRECTIONS

ORNAMENTAL PLANTINGS

ORNAMENTAL WEEDER controls germinating annual broadleaf weeds and grasses in ornamental plantings, ground covers and conifer seed beds. Preemergence application of ORNAMENTAL WEEDER prevents germination of these weeds for 6 to 8 weeks after treatment.

Apply ORNAMENTAL WEEDER granules to weed free ornamental plantings at the rate of 100 lbs. per acre. ORNAMENTAL WEEDER must be applied in the spring before weeds germinate, or to freshly cultivated soil. It will not control weeds which have already germinated or emerged. It will not control perennial weeds.

Do not incorporate ORNAMENTAL WEEDER into the soil.

If no rain falls within one week of application, apply ½ inch of overhead irrigation. This will wash the chemical into the soil where it kills germinating weeds.

ORNAMENTAL WEEDER may be used on the following:

	TIME FOR APPLICATION		
	Immediately after Transplanting	One month after transplanting	Established stock (6 wks or more)
[illegible]			X
[illegible]			X
[illegible]			X
[illegible]			X
[illegible]	X	X	X
[illegible]			X
[illegible]			X
[illegible]	X	X	X
[illegible]	X	X	X
[illegible]			X
[illegible]	X	X	X
[illegible]			X
[illegible]	X	X	X
[illegible]			X
[illegible]	X	X	X
[illegible]			X
[illegible]	X	X	X
[illegible]	X	X	X
[illegible]	X	X	X
[illegible]			X
[illegible]			X
[illegible]	

ting annual broadleaf weeds and grasses in
nifer seed beds. Preemergence application
nination of these weeds for 6 to 8 weeks

s to weed free ornamental plantings at the
WEEDER must be applied in the spring
ated soil. It will not control weeds which
ot control perennial weeds.
DER into the soil
ation, apply ½ inch of overhead irrigation.
e it kills germinating weeds.

the following:

TIME FOR APPLICATION	
One month after transplanting	Established stock (6 wks or more)
	X
	X
	X
	X
X	X
	X
X	X
	X
X