

- Controls growing of perennial grasses and broadleaf weeds.
- Prevents sprouting of annual weeds and grasses.
- Do not apply or use where runoff from the treated area may contact roots or foliage of desirable trees, shrubs or plants. Do not use on lawns, walks or driveways.

DIRECTIONS

Apply FENAVAR GRANULAR before weed growth begins in the spring. In areas of low or seasonal rainfall FENAVAR GRANULAR may be applied in the fall or winter while weeds are dormant. Rainfall is necessary to wash the chemical into the soil. There it contacts the roots of perennial weeds and kills annual weeds and grasses as they germinate. Do not apply when ground is frozen. If FENAVAR GRANULAR is applied when weeds are growing, the action is slow and dead weeds may remain standing in the treated area during the growing season.

AMOUNTS TO USE

Apply 75 pounds of FENAVAR GRANULAR per acre to control annual weeds and grasses such as foxtail, ryegrass, crabgrass, cheat, lambsquarters, pigweed and ragweed.

Apply 150 pounds of FENAVAR GRANULAR per acre to control biennial and perennial weeds and grasses such as smooth brome, bluegrass, bahiagrass, quackgrass, broomsedge, horsenettle, dandelion and wild carrot.

Repeat applications of the 150 pound per acre rate may be necessary to control johnsongrass and bermudagrass.

TO APPLY

Use a spreader that gives even distribution of granules. Calibrate spreaders accurately and re-check calibration periodically. To calibrate PCB, Amoco and Seymour Spreaders, use the following procedure:

1. Set spreader to treat a 10 foot swath. (Slide should be wide open)
2. Calibrate yourself to walk 3 mph. (Measure 100 feet and walk it in 23 seconds for 3 mph.) Check your speed regularly.
3. Set opening regulator at desired setting.

Rate of Granule	Spreader Setting 10 foot swath — 3 mph
40 lb/A	4
75 lb/A	6
150 lb/A	8

4. For more uniform coverage, apply half the rate and treat area twice. (The left edge of the spreader swath is straighter than the right edge, so a two-way treatment will give more uniform edges.)

For small areas, use the following amounts:

Square Feet	Area to Treat (Examples)	Amount FENAVAR GRANULAR	
		@ 75 lb/A Rate	@ 150 lb/A Rate
5,000	(50' x 100')	9 lb.	17 lb.
10,000	(100' x 100')	17 lb.	35 lb.
20,000	(100' x 200')	35 lb.	70 lb.
30,000	(100' x 300')	50 lb.	100 lb.

FENAVAR[®]

GRANULAR

GENERAL WEED KILLER

for industrial areas, pole yards, outdoor storage areas, parking lots and utility substations.

CAUTION: Keep out of reach of children. See other cautions on right side panel.

ACTIVE INGREDIENTS:

Sodium salt of 2,3,6-trichlorophenylacetic acid* 3.63
 Sodium salt of bromacil (5-bromo-3-sec-butyl 6-methyluracil)** 2.9

INERT INGREDIENTS: 93.47

*fenac acid equivalent 3.33% by weight
 **bromacil equivalent 2.67% by weight



AMCHEM PRODUCTS, INC

First Name in Herbicide Research
 AMBLER, PA ■ Clinton, Iowa ■ St. Louis, Mo ■ Fort Worth, Texas

EPA Reg. No. 244-245

25 POUNDS

FENAVAR GRANULAR

GENERAL WEED KILLER

Industrial areas, pole yards,
dr storage areas, parking lots
utility substations.

Caution: Keep out of reach of children.
See other cautions on right side panel.

INGREDIENTS:

of 2,3,6-trichlorophenylacetic acid* 3.63%
of bromacil (5-bromo-3-sec-butyl-6-methyluracil)** 2.9%
INGREDIENTS: 93.47%

acid equivalent 3.33% by weight
cil equivalent 2.67% by weight

AMCHEM PRODUCTS, INC.

First Name in Herbicide Research
AMBLER, PA ■ Clinton, Iowa ■ St. Louis, Mo ■ Ft. Worth, Tex

EPA Reg. No. 264-245

Some of the weeds controlled by recommended rates of FENAVAR GRANULAR:

- alfalfa
- bermudagrass
- bluegrass
- chrysanthemum weed
- crabgrass
- dandelion
- downy brome grass
(cheat grass)
- filaree
- foxtail
- johnsongrass
- kochia
- lambsquarters
- mustard
- nightshade
- pigweed
- plantain
- puncture vine
- purslane
- quackgrass
- ragweed
- Russian thistle
- ryegrass
- smartweed
- sowthistle
- tarweed
- velvet grass
- wild oats
- witchgrass

CAUTION

HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. IN CASE OF CONTACT, WASH THOROUGHLY WITH SOAP AND WATER.

Most ornamental shrubs are susceptible to this mixture and will be injured if contacted by granules or runoff from treated areas. Do not use this product near ornamentals or other valuable plants with roots extending into treated areas. For controlling weeds near ornamental shrubs, AMIZINE is suggested.

Do not use this material to control weeds in flower or vegetable beds, field crops or in turf.

Do not use the same spreader for other purposes. Do not contaminate or store near feed, foodstuffs, fertilizers, seeds, insecticides, fungicides, or use in a greenhouse.

Do not reuse containers. Destroy when empty. Do not contaminate water used for irrigation, domestic or fish purposes.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 1-7508 (10-74) Printed in U.S.A.

ACCEPTED WITH COMMENTS