

WEEDONE PREEMERGENCE CRABGRASS CONTROL

# WEEDONE PREEMERGENCE CRABGRASS CONTROL

NET CONTENTS

8.3 POUNDS

## CRABGRASS CONTROL



ACCEPTED
JUL 18 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 264-249

Controls crabgrass and other annual weeds  
in home lawn, golf course and athletic field turf.



TREATS 2,500 SQUARE FEET

CAUTION: Keep out of reach of children. See other cautions on back panel.

ACTIVE INGREDIENTS

Dimethyltetrachloroterephthalate\*\*

INERT INGREDIENTS

\*Reg. TM of Diamond Shamrock Corp.

\*\*Use as a preemergence weed killer patented under U.S. Pat. No. 3,421,111



AMCHEM PRODUCTS, INC.

AMCHEM PRODUCTS, INC. • 1000 N. 10TH ST. • PITTSBURGH, PA. 15222

EPA Reg No 264-249AA EPA Est 264 PA 1

### WEEDONE PREEMERGENCE CRABGRASS CONTROL

1000 N. 10TH ST. PITTSBURGH, PA. 15222

WEEDONE PREEMERGENCE CRABGRASS CONTROL

# WEEDONE PREEMERGENCE CRABGRASS CONTROL

See  
Plan  
←

## DIRECTIONS

### USE IN HOME LAWN, GOLF COURSE AND ATHLETIC FIELD TURF

#### When to Apply:

**WEEDONE PREEMERGENCE CRABGRASS CONTROL** Granular kills crabgrass and other annual weeds as they germinate. Apply in early spring, before these weeds sprout. This is usually about the time Forsythia blossoms start to fall. Spread on a calm day to avoid drift.

Before spreading, rake the area to remove sticks, leaves or other material.

Among the common lawn weeds controlled are:

common chickweed  
tall panicle  
green foxtail  
hairy (large) crabgrass  
lambquarters

purple  
smooth (small) crabgrass  
witchgrass  
yellow foxtail  
goosegrass

#### How to Apply:


Apply contents of this bag (5.2 pounds) evenly to 2,500 square feet (example 50 ft x 50 ft., or 25 ft x 100 ft.) Use a lawn spreader in good condition. Do not try to spread over more than 2,500 sq. ft. as this would give poor results.

To prevent skips or misses, cover the area two times, spreading half of the bag contents from end to end, then the remaining contents from side to side.

Be sure to shut off spreader when stopping to turn around.

Overlap wheel tracks by 2 to 3 inches to avoid missed or overdone strips. If grass is wet, check hopper opening occasionally to be sure material is flowing freely.

Do not rake the lawn or disturb soil surface after application. Water in lightly if no rain falls within a week after application.



#### Suggested Spreader Settings:

Spreader	Width	Setting
CENTRAL	20"	7 1/2
HERITAGE HOUSE	20"	6
LAWN BEAUTY	24"	8 1/2
SCOTT'S	24"	6 1/2
SEARS	20"	7 1/2

#### Re-Seeding:

Areas seeded in the fall may be treated with **WEEDONE PREEMERGENCE CRABGRASS CONTROL** the following spring. However, we do not recommend using this product on young spring seeded turf.

To seed into areas treated with this product, wait at least 60 days. If seeding must be done sooner, increase the amount of seed by 50 per cent to assure a satisfactory stand.

### USE IN NURSERY STOCK OR ORNAMENTAL PLANTINGS

Apply uniformly at the rate of 5.2 pounds of **WEEDONE PREEMERGENCE CRABGRASS CONTROL GRANULAR** per 2,500 to 3,500 square feet of planted beds. Spread prior to grass seed germination in the spring, or to freshly weeded beds in summer or fall. Weeds up of time of treatment are not controlled. Light watering or moderate rainfall following treatment will assure best results.

On newly transplanted beds, delay application for two or three weeks after transplanting, or until roots are established.

Among the many tolerant ornamentals are:

agrostis  
stachys  
arbor vitae  
cush  
cotton  
cotton

quercus  
br  
forythia  
laugave  
gardenia  
cotton

poonia  
pine  
paper  
pines  
redwood

### How to Apply


Apply contents of this bag (2.5 pounds) evenly to 2,000 square feet (example: 50 ft. x 50 ft., or 20 ft. x 100 ft.). Use a lawn spreader in good condition. Do not try to spread over more than 2,000 sq. ft., as this would give poor results.

To prevent clumps or misses, cover the area two times, spreading half of the bag contents from end to end, then the remaining contents from side to side.

Be sure to shut off spreader when stopping to turn around.

Overlap wheel tracks by 2 to 3 inches to avoid missed or overdone strips. If grass is wet, check hopper opening occasionally to be sure material is flowing freely.

Do not rake the lawn or disturb soil surface after application. Water in lightly if no rain falls within a week after application.



### Suggested Spreader Settings:

Spreader	Width	Setting
CENTRAL	20"	7 1/2
HERITAGE HOUSE	20"	8
LAWN BEAUTY	24"	8 1/2
SCOTT'S	24"	8 1/2
SCARF	20"	7 1/2

### No-Seeding:

Areas seeded in the fall may be treated with WEEDONE PREMIXED CRABGRASS CONTROL the following spring. However, we do not recommend using this product on young spring-seeded turf.

To seed into areas treated with this product, wait at least 60 days. If seeding must be done sooner, increase the amount of seed by 50 per cent to ensure satisfactory stand.

### USE IN NURSERY STOCK OR ORNAMENTAL PLANTINGS

Apply uniformly at the rate of 0.5 pounds of WEEDONE PREMIXED CRABGRASS CONTROL GRASS-LAK per 2,000 to 2,500 square feet of planted beds. Spread prior to grass weed germination in the spring, or to freshly weeded beds in summer or fall. Winds up at time of treatment are not controlled. Light watering or moderate rainfall following treatment will assure best results.

On newly transplanted beds, apply application for two or three weeks after transplanting, or until roots are established.

Among the many tolerant ornamentals are:

agave  
 anemone  
 arbor vitae  
 ash  
 aster  
 azalea  
 barberry  
 birch  
 bleeding heart  
 Boston ivy  
 boxwood  
 camellia  
 candy tuft  
 chrysanthemum  
 columbine  
 clematis  
 carnation  
 coral bells  
 cosmos  
 columbine  
 crab apple  
 dahlia  
 delphinium  
 dogwood  
 elm  
 English ivy

geranium  
 geranium  
 gladiolus  
 heather  
 holly  
 hollyhock  
 hydrangea  
 iris  
 Japanese andromeda  
 juniper  
 lilac  
 lily  
 locust  
 lupine  
 magnolia  
 maple  
 marigold  
 mock orange  
 morning glory  
 mountain laurel  
 nasturtium  
 oak  
 pachysandra

pearl  
 philodendron  
 poplar  
 privet  
 rhododendron  
 rose  
 Russian olive  
 salvia  
 snapdragon  
 spirea  
 spruce  
 strawflower  
 sunflower  
 sweet pea  
 sweetshrub  
 tree peony  
 tulip tree  
 variegated privet  
 verbena  
 viburnum  
 violet  
 walnut  
 weigela  
 willow  
 yew  
 zinnia

DO NOT use on the following plants:

bugle weed  
 button pink  
 carnation

gum  
 germander  
 mesembryanthemum  
 pansy

phlox  
 sweet william  
 tolanthera

## CAUTION

Avoid contact with eyes. Wash after using.

Do not use on dandelion lawns or on newly seeded lawns. Some thinning of fescue grass has been observed.

Do not store near seeds, fertilizers, and other pesticides. Do not contaminate food or feedstuffs.

Do not reuse bag. Discard in a safe place. Do not burn.

Do not use this product for purposes other than those described on this label.

## WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore, makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 2009E-6-75

Printed in U.S.A.



**AMCHEM PRODUCTS, INC.**

First Name in Herbicide Research

AMBLER PA ■ Clinton IA ■ St. Joseph MO ■ Fremont CA

# Code No. 112265