

thoroughly mix in the required amount of PAY-OFF according to dilution rates listed under *Amount to Apply*. Tank pressure should not exceed 20 psi so as to develop a *coarse spray*. The nozzle should be held so as to direct the spray downward into the top of the plant. Apply enough spray solution per plant so that liquid flows freely all the way down the stalk, so that the sucker buds from the top of the plant to the base are covered.

1. *Wind*. Do not spray when wind is blowing hard enough to turn over top leaves so that the underside is exposed. Leaf injury and poor coverage of sucker buds can result.
2. *Rain*. Do not spray just before or during rain or when plants are wet or covered with heavy dew.
3. *Drought*. Scald and similar injury can occur if sprays are applied to wilted leaves.
4. *Mixtures*. Compatibility with other pesticides is unknown. Do not mix or apply simultaneously with other pesticides.
5. *Drift*. Avoid drift to other crops when spraying.

Lot No.

Made and Printed in U.S.A.

**ACCEPTED**  
 264-250  
 10-13-76

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 264-250 SUBJECT  
 TO ATTACHED COMMENTS.

# NU-LAWN WEEDER

**CAUTION:** Keep out of the reach of children.  
 See other cautions on back panel.

CONTENTS

1 PINT



**in seedling turf**  
 Established Turf in California Only



**AMCHEM PRODUCTS, INC.**  
 First Name in Herbicide Research  
 AMBLER PA ■ Clinton Iowa ■ St Joseph Mo ■ Fremont Calif

EPA Reg. No. 264-250AA

EPA Est. 264-PA

**ACTIVE INGREDIENT:**  
 Bromoxynil (3,5-dibromo-4-hydroxybenzotrile)  
 ester of octanoic acid.....8.4%

**INERT INGREDIENTS:**  
 .....91.6%  
 \*bromoxynil equivalent 5.8% by weight or 0.5 pounds per gallon.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Avoid breathing spray mist. Keep all animals off treated areas for at least one week after treatment or until after first rainfall or irrigation.

**STATEMENT OF PRACTICAL TREATMENT**  
 If contact with eyes, flush with plenty of water and get medical attention. If contact with skin, wash with plenty of soap and water.

**ENVIRONMENTAL HAZARDS**  
 Keep out of lakes, streams, or ponds as this product is toxic to fish and wildlife. Do not contaminate water used for irrigation or domestic purposes. Clean sprayers thoroughly before and after using NU-LAWN WEEDER.

**DIRECTIONS FOR USE**  
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
**Storage:** Do not store near food stuffs, fertilizers, seeds, insecticides or fungicides.  
**Disposal:** Do not re-use empty containers. Discard in a safe place.

**GENERAL INFORMATION**  
 See Attached Pamphlet for Detailed Instructions  
 NU-LAWN WEEDER effectively controls seedling broadleaf weeds in newly planted home lawns, golf course, ornamental, or athletic field turf. NU-LAWN can be applied to grasses at a younger stage than is recommended with standard lawn weed killers containing 2,4-D

other phenoxy chemicals. Competition from broadleaf weeds is removed earlier, allowing seedling grasses to grow and become established faster. NU-LAWN WEEDER can also be used on established turf in California to control or suppress seedling broadleaf weeds.

NU-LAWN weeder will control or suppress the following seedling broadleaf weeds:

mustards	soliva sp.
penny cress	Bur clover
smartweed	ragweed
henbit	shepherdspurse
knawel	wild radish
lambsquarters	yellow rocket
pepperweed	spiny fruited crowfoot
pigweed	black medic

This treatment will not control established perennial broadleaf weeds such as Canada thistle, bindweed, or horsetail rush.

**WARRANTY**  
 Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of the product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Form No. 2874-B 8-76

Screened in U.S.A.



# NU-LAWN WEEDER

weed  
control

in seedling turf  
Established Turf in California Only

264-250



AMCHEM PRODUCTS, INC.

First Name in Herbicide Research  
AMBLER PA • Clinton Iowa • St Joseph Mo • Fremont Calif

### DIRECTIONS FOR USE

#### NEW LAWNS

Apply one-half pint (8 fluid ounces) of NU-LAWN WEEDER to 2,500 square feet (example: 25 ft. by 100 ft. or 50 ft. by 50 ft.). For best results, spray after grasses emerge and before broadleaf weeds are past the 3 to 4 leaf stage. Weeds that form rosettes should be sprayed when these rosettes are less than 1 1/2 inches across. Applications to larger weeds will be less effective.

Do not use on cool season grasses when temperatures are above 80°F.

One pint NU-LAWN WEEDER treats 5,000 square feet of grasses such as Merion, Park, Delta or common Kentucky bluegrasses, Pennlawn, Chewings, Illahee or Alta fescues, Highland, Seaside or Astoria bentgrass, perennial ryegrass.

Mix NU-LAWN WEEDER in enough water for uniform coverage of all weeds. For power sprayer equipped with booms, use at least 3 gallons of water per 2,500 square feet, or 50-100 gallons per acre.

When using the Amchem METER-MISER<sup>®</sup>, mix 8 ounces of NU-LAWN WEEDER with 5 gallons of water (one tankful). This quantity of spray mixture will cover 5,000 square feet of area. When the tank is empty repeat the procedure and apply the same amount of spray mixture to the same area, walking at right angles to the direction of the first application.

When using Amchem's WEEDONE<sup>®</sup> LAWN WEED SPRAYER, place one-half pint (8 fluid ounces) of NU-LAWN WEEDER in the sprayer jar and fill the remaining space with water. Spray this mix over 2,500 square feet of lawn, first from end to end and then from side to side.

### DIRECTIONS FOR USE

#### ESTABLISHED LAWNS IN CALIFORNIA

For seedling weeds such as soliva sp., mustard, lambs-quarter, pigweed in established lawns (2-3 months after seed germination) apply one-half pint NU-LAWN WEEDER to 2,500 sq. ft. Two quarts will treat 10,000 sq. ft. If weeds are at the 3-4 leaf stage, a repeat treatment will be needed in 10-14 days for control or suppression. To control or suppress hard to kill weeds such as Black Medic, Bur Clover, and Spiny Fruited Crowfoot, apply 1 pint NU-LAWN WEEDER to 2,500 sq. ft.

Do not apply more than 1 pint of NU-LAWN WEEDER per 2,500 sq. ft. to the same area within one year.

Mix NU-LAWN WEEDER in enough water for uniform coverage of all weeds. For power sprayers equipped with booms, use at least 3 gallons of water per 2,500 square feet, or 50 gallons per acre.

When using the Amchem METER-MISER<sup>®</sup>, mix one-half to one pint (8-16 fluid ounces) of NU-LAWN WEEDER with 5 gallons of water (one tankful). This quantity of spray mixture will cover 5,000 square feet of area. When the tank is empty repeat the procedure and apply the same amount of spray mixture to the same area, walking at right angles to the direction of the first application.

When using Amchem's WEEDONE<sup>®</sup> LAWN WEED SPRAYER, place one-half to one pint (8-16 fluid ounces) of NU-LAWN WEEDER in the sprayer jar and fill the remaining space with water. Spray this mix over 2,500 square feet of lawn, first from end to end and then from side to side.

Temporary turfgrass discoloration may occur particularly if excessive overlapping of spray solution occurs, or if NU-LAWN is applied at temperatures of 85°F or above. This will usually disappear following two mowings leaving no permanent damage.

Form No. 4054 (1-6-76)