

115 7 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR FEDERAL POISON REGISTER
 LISTED UNDER NO. 264 23 SUBJECT
 TO ATTACHED COMMENTS.

CAUTION

Harmful if swallowed.
 Avoid contact with skin, eyes or clothing; do not inhale dust.
 Do not store near fertilizers, seeds, insecticides, or fungicides, or allow contamination of food or feed.
 Clean sprayers thoroughly before using to apply CORIBEN-WP. To clean sprayer after using CORIBEN-WP, rinse all parts several times with water.
 Avoid spray drift to adjacent crops as injury may result. Coarse sprays are less likely to drift.
 Do not contaminate water used for irrigation or domestic purposes by cleaning equipment or disposal of excess spray.
 Do not reuse empty containers. Wash thoroughly with water and detergent. Dispose of rinsings and containers in a safe place not accessible to children, wildlife or livestock.
 Do not use this product for purposes other than those recommended on this label.

MIXING SPRAYS

Make certain spray equipment is clean (free of scale, rust, dirt, oil, and pesticide deposits). Screens should be 50-mesh or coarser. Booms, and hoses should be maximum of 1/2" diameter. MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION. Agitate by mechanical or hydraulic means. With hydraulic system, use a separate agitation line in addition to by-pass line; both lines should terminate at bottom of tank to minimize foaming. Spray booms must be shut off while starting, turning, slowing or stopping, or injury to crop may result.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the product as full discharge of Amchem's liability.

DIRECTIONS

WEED CONTROL IN FIELD CORN

BROADCAST APPLICATION:

Apply 4 pounds of CORIBEN-WP per acre. For best results plant and treat in the same operation.

Use on clay loam or clay soils in Missouri, Minnesota, Iowa, Ohio, Illinois, Kansas, Nebraska and Indiana.

Plant corn at least 2 to 2 1/2 inches deep. Do not use CORIBEN-WP on corn grown in light-textured sandy soils.

BAND APPLICATION:

Cost for chemicals are reduced by treating a band over the crop row. Reduce the amount of CORIBEN-WP applied per acre in proportion to the actual area treated. (Banded treatments reduce chemical residues in soil by applying less herbicide per acre.)

Mount the band sprayer on the planter so that treating and planting are carried out in the same operation. Maintaining proper nozzle height above the ground is very important when making band application. Make final adjustments while actually treating the field.

Pounds CORIBEN-WP Per Banded Acre		
Row Width	Band Width	Pounds/Acre
30 inch	12 inches	1.6
	14 inches	1.9
36 inch	12 inches	1.3
	14 inches	1.5
40 inch	12 inches	1.2
	14 inches	1.4

Weeds growing in the middle between chemically treated rows should be cultivated as soon as necessary. Do not cover treated bands with fresh soil.

If heavy rain and extended cold weather occur after planting, corn injury may result.

GENERAL

- CORIBEN-WP has
- Excellent crop tolerance
- No chemical accidents
- Available power
- Controls annual and perennial grasses.
- Shows cultivation and deep weed control
- Consistent weed control under weather conditions

HOW CORIBEN-WP

CORIBEN-WP is a pre-emergent herbicide. It kills germinating weeds, the seed while the seed is in the soil, while the seed is on the surface.

CORIBEN-WP controls broadleaves and grasses.

- giant foxtail
- bristly foxtail
- green foxtail
- yellow foxtail
- pigweeds
- smartweeds
- lamb's quarters
- crabgrass
- fall panicum
- ragweed
- velvetleaf
- barnyardgrass
- carpetweed
- purslane
- wild mustard

Velvetleaf, jimsonweed, and other weeds often escape the chemical control of this herbicide. Under certain conditions, weeds in the chemical zone are not controlled.

Some weeds from which weeds such as horseweed, milkweed, dogfennel, and other weeds are not controlled.

CORN

ults plant and treat

owa, Ohio, Illinois,

CORIBEN-WP on corn

the crop row. Re-
proportion to the
residues in soil by

g and planting are
bottle height above
ication. Make final

acre

Pounds/Acre

1.6
1.9
1.3
1.5
1.2
1.4

ted rows should be
ands with fresh soil.
planting, corn injury

GENERAL INFORMATION

CORIBEN-WP has many desirable features:

- Excellent crop tolerance.
- No chemical accumulation in soils.
- A wettable powder that mixes easily in water.
- Controls annual broadleaf weeds and annual grasses.
- Shallow cultivation after application does not destroy weed control.
- Consistent weed control with varied weather conditions.

HOW CORIBEN-WP WORKS

CORIBEN-WP is adsorbed through the roots of germinating weeds. Rainfall carries AMIBEN into the soil while the less soluble atrazine remains near the surface.

CORIBEN-WP controls the following Annual Broadleaves and Grasses:

giant foxtail	<i>Setaria faberii</i>
bristly foxtail	<i>Setaria glauca</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria lutescens</i>
pigweeds	<i>Amaranthus</i> spp.
smartweeds	<i>Polygonum</i> spp.
lambquarters	<i>Chenopodium album</i>
crabgrass	<i>Digitaria</i> spp.
fall panicum	<i>Panicum dichotomiflorum</i>
ragweed	<i>Ambrosia artemisiifolia</i>
velvetleaf	<i>Abutilon theophrasti</i>
barnyardgrass	<i>Echinochloa crusgalli</i>
carpetweed	<i>Mollugo verticillata</i>
purslane	<i>Portulaca oleracea</i>
wild mustard	<i>Brassica kaber</i>

Velvetleaf, Jimsonweed, buffalobur, wild oats and cocklebur often germinate in soil at depths below the chemical penetration. Even though populations of these weeds are reduced under normal rainfall conditions, weed seeds which germinate outside the chemical zone are not affected.

CONTENTS

8 POUNDS



CORIBEN-WPTM

PREEMERGENCE WEEDKILLER

EPA REG. No. 261-113

Controls Annual Broadleaf Weeds and Annual Grasses in Field Corn

CAUTION: Keep out of reach of children. See other cautions on right side panel

ACTIVE INGREDIENTS:

Sodium salt of amiben (3-amino-2,5-dichlorobenzoic acid)*	27.7%
Atrazine (2-chloro-4-(ethylamino)-6-isopropylamino-s-triazine)	50.0%
INERT INGREDIENTS:	22.3%

*3-amino-2,5-dichlorobenzoic acid equivalent 25.0% by weight

U.S. Pat. 3,014,063 and 2,819,855

ACCEPTED
264-253

5
2
4
2
5
3

CAUTION

- Harmful if swallowed.
- Avoid contact with skin, eyes or clothing, do not inhale dust.
- Do not store near fertilizers, seeds, insecticides, or fungicides, or allow contamination of food or feed.
- Clean sprayers thoroughly before using to apply CORIBEN-WP. To clean sprayer after using CORIBEN-WP, rinse all parts several times with water.
- Avoid spray drift to adjacent crops as injury may result. Coarse sprays are less likely to drift.
- Do not contaminate water used for irrigation or domestic purposes by cleaning equipment or disposal of excess spray.
- Do not reuse empty containers. Wash thoroughly with water and detergent. Dispose of rinsings and containers in a safe place not accessible to children, wildlife or livestock.
- Do not use this product for purposes other than those recommended on this label.

MIXING SPRAYS

Make certain spray equipment is clean (free of scale, rust, dirt, oil, and pesticide deposits). Screens should be 50-mesh or coarser. Booms, and hoses should be maximum of 1/2" diameter. MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION. Agitate by mechanical or hydraulic means. With hydraulic system, use a separate agitation line in addition to by-pass line; both lines should terminate at bottom of tank to minimize foaming. Spray booms must be shut off while starting, turning, slowing or stopping, or injury to crop may result.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the application of this product, it is impossible to guarantee results, although label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risk and in the event of crop injury, to accept the product as full discharge of Amchem's liability.

DIRECTIONS

WEED CONTROL IN FIELD CORN

BROADCAST APPLICATION:

Apply 4 pounds of CORIBEN-WP per acre. For best results plant and treat in the same operation.
Use on clay loam or clay soils in Missouri, Minnesota, Iowa, Ohio, Illinois, Kansas, Nebraska and Indiana.
Plant corn at least 2 to 2 1/2 inches deep. Do not use CORIBEN-WP on corn grown in light-textured sandy soils.

BAND APPLICATION:

Cost for chemicals are reduced by treating a band over the crop row. Reduce the amount of CORIBEN-WP applied per acre in proportion to the actual area treated. (Banded treatments reduce chemical residues in soil by applying less herbicide per acre.)
Mount the band sprayer on the planter so that treating and planting are carried out in the same operation. Maintaining proper nozzle height above the ground is very important when making band application. Make final adjustments while actually treating the field.

Pounds CORIBEN-WP Per Banded Acre		
Row Width	Band Width	Pounds/Acre
30 inch	12 inches	1.6
	14 inches	1.9
36 inch	12 inches	1.3
	14 inches	1.5
40 inch	12 inches	1.2
	14 inches	1.4

Weeds growing in the middle between chemically treated rows should be cultivated as soon as necessary. Do not cover treated bands with fresh soil.

If heavy rain and extended cold weather occur after planting, corn injury may result.

BEST DOCUMENT AVAILABLE

NS
FIELD CORN

For best results plant and treat
Minnesota, Iowa, Ohio, Illinois,

do not use CORIBEN-WP on corn

band over the crop row. Re-
1 per acre in proportion to the
duce chemical residues in soil by

that treating and planting are
ing proper nozzle height above
band application. Make final

Banded Acre	
	Pounds/Acre
	1.6
	1.9
	1.3
	1.5
	1.2
	1.4

nically treated rows should be
r treated bands with fresh soil.
our after planting, corn injury

GENERAL INFORMATION

CORIBEN-WP has many desirable features:

- Excellent crop tolerance.
- No chemical accumulation in soil.
- A wettable powder that mixes easily in water.
- Controls annual broadleaf weeds and annual grasses.
- Shallow cultivation after application does not destroy weed control.
- Consistent weed control with varied weather conditions.

HOW CORIBEN-WP WORKS

CORIBEN-WP is adsorbed through the roots of germinating weeds. Rainfall carries AMIBEN into the soil while the less soluble atrazine remains near the surface.

CORIBEN-WP controls the following Annual Broadleaves and Grasses:

- | | |
|-----------------|--------------------------------|
| giant foxtail | <i>Setaria faberii</i> |
| bristly foxtail | <i>Setaria glauca</i> |
| green foxtail | <i>Setaria viridis</i> |
| yellow foxtail | <i>Setaria lutescens</i> |
| pigweeds | <i>Amaranthus spp.</i> |
| smartweeds | <i>Polygonum spp.</i> |
| lambquarters | <i>Chenopodium album</i> |
| crabgrass | <i>Digitaria spp.</i> |
| fall panicum | <i>Panicum dichotomiflorum</i> |
| ragweed | <i>Ambrosia artemisiifolia</i> |
| velvetleaf | <i>Abutilon theophrasti</i> |
| barnyardgrass | <i>Echinochloa crusgalli</i> |
| carpetweed | <i>Mollugo verticillata</i> |
| purslane | <i>Portulaca oleracea</i> |
| wild mustard | <i>Brassica kaber</i> |

Velvetleaf, jimsonweed, buffalobur, wild oats and cocklebur often germinate in soil at depths below the chemical penetration. Even though populations of these weeds are reduced under normal rainfall conditions, weed seeds which germinate outside the chemical zone are not affected.

Sprouts coming from roots of established perennial weeds such as horseweed, hedge bindweed, grass, milkweed, doanone, quackgrass and johnson-grass are not controlled.

COR
PR
V

Controls
at

CAUTION
See (

ACTIVE INGREDIENTS:
Sodium salt of amiben
Atrazine (2-chloro-4-
INER INGREDIENTS: ..
*3-amino-2,5-dichloro



BEST DOCUMENT AVAILABLE

GENERAL INFORMATION

CORIBEN-WP has many desirable features:

- Excellent crop tolerance.
- No chemical accumulation in soils.
- A wettable powder that mixes easily in water.
- Controls annual broadleaf weeds and annual grasses.
- Shallow cultivation after application does not destroy weed control.
- Consistent weed control with varied weather conditions.

HOW CORIBEN-WP WORKS

CORIBEN-WP is adsorbed through the roots of germinating weeds. Rainfall carries AMIBEN into the soil while the less soluble atrazine remains near the surface.

CORIBEN-WP controls the following Annual Broadleaves and Grasses:

giant foxtail	<i>Setaria faberii</i>
bristly foxtail	<i>Setaria glauca</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria lutescens</i>
pigweeds	<i>Amaranthus</i> spp.
smartweeds	<i>Polygonum</i> spp.
lambsquarters	<i>Chenopodium album</i>
crabgrass	<i>Digitaria</i> spp.
fall panicum	<i>Panicum dichotomiflorum</i>
ragweed	<i>Ambrosia artemisiifolia</i>
velvetleaf	<i>Abutilon theophrasti</i>
barnyardgrass	<i>Echinochloa crusgalli</i>
carpetweed	<i>Mollugo verticillata</i>
purslane	<i>Portulaca oleracea</i>
wild mustard	<i>Brassica kaber</i>

Velvetleaf, jimsonweed, buffalobur, wild oats and cocklebur often germinate in soil at depths below the chemical penetration. Even though populations of these weeds are reduced under normal rainfall conditions, weed seeds which germinate outside the chemical zone are not affected.

Sprouts coming from roots of established perennial weeds such as horsetail, bedstraw, wild radish, quackgrass, and nutcracker, quackgrass, and nutcracker are not controlled.

CONTENTS

8 POUNDS



CORIBEN-WPTM PREEMERGENCE WEEDKILLER

EPA REG. No. 264-253

Controls Annual Broadleaf Weeds and Annual Grasses in Field Corn

CAUTION: Keep out of reach of children.
See other cautions on right side panel.

ACTIVE INGREDIENTS:

Sodium salt of amiben (3-amino-2,5-dichlorobenzoic acid)*	27.7
Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	50.0
INERT INGREDIENTS	22.3

*3-amino-2,5-dichlorobenzoic acid equivalent 25.0% by weight

U.S. Pat. 3,014,063 and 2,819,855

32

AMCHEM PRODUCTS, Inc.
Originator of 2,4-D-2,4,5-T and Amiblen Weed Killer

AMBLER, PA. • Clinton, Iowa • St. Joseph, Mo. • Fremont, Calif.