264-257



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

201

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Bayer_CropScience____ Laura Phelps 2 T.W. Alexander Drive RTP, NC 27709

MAY 1 1 2011

Subject: ETHREL brand Ethephon Plant Growth Regulator Registration No.: 264-257 Date Submitted: 3/14/11 Decision No.: 447178

> PREP brand Ethephon Plant Growth Regulator for Cotton Registration No.: 264-380 Date Submitted: 3/14/11 Decision No.: 447179

PREP brand Ethephon Plant Growth Regulator for Cotton and Tobacco Registration No.: 264-418 Date Submitted: 3/14/11 Decision No.: 447180

CERONE brand Ethephon Plant Growth Regulator Registration No.: 264-377 Date Submitted: 3/14/11 Decision No.: 447181

Dear Ms. Phelps:

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10.

The Agency acknowledges the removal of the patent number from the label.

The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have questions concerning this letter, please contact Dominic Schuler at 703-347-0260 or via email at schuler.dominic@epa.gov.

Sincerely Tony Kish

Product Manager 22 Fungicide Branch Registration Division (7504P)

	S&12152
Please read instructions on reverse before continued form. United States Environmental Protection Ag Washington, DC 20460	Jency Approved, <u>B No. 2070-0060</u> Registration Amendment × Other
Application for	Pesticide - Section I
1. Company/Product Number 264-257	2. EPA Product Manager 3. Proposed Classification Mr. Tony Kish Image: Classification
4. Company/Product (Name) Ethrei brand Ethephon Plant Growth Regulator	PM# Restrict
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience LP P. O. Box 12014; 2 T. W. Alexander Dr. Research Triangle Park, NC 27709	 6. Expedited Review. In accordance with FIFRA Section 3(c){3 (b){i}, my product is similar or identical in composition and labeling to: EPA Reg. No
	Product Name
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below. Explanation: Use additional page(s) if necessary. (For section I and Bayer CropScience is submitting a notification for Ethrel brand E label.	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below. Section II.)
Se 1. Material This Product Will Be Packaged In:	ction - III
Yes* Yes No If "Yes" No. per If "Yes"	er Soluble Packaging 2. Type of Container Yes Metal Plastic Glass Yes" No. per Kage wgt container Other (Specify)
be submitted 3. Location of Net Contents Information 4. Size(s) Retail Con	
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled	On Labeling accompanying product
	ction - IV
1. Contact Point IComplete items directly below for identification of inc	
Name	Affairs Crossielist
•	
Certification I certify that the statements I have made on this form and all atta I acknowledge that any knowingly false or misleading statement both under applicable law.	دورد د. دوردد 6. Date Application Accord Received
Certification I certify that the statements I have made on this form and all atta I acknowledge that any knowingly false or misleading statement both under applicable law. 2. Signature	دون دور دور دور دور الم الم الم الم الم الم الم الم الم الم

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

Bayer CropScience



March 10, 2011

Document Processing Desk (NOTIF) Registration Division (7505P) Office of Pesticide Programs U.S. Environmental Protection Agency One Potomac Yard (South Building) 2777 S. Crystal Drive Arlington, VA 22202

Mr. Tony Kish, RD Team 22

Subject: Ethrel brand Ethephon Plant Growth Regulator. No. 264-257; Removal of Patent Number

Dear Mr. Kish:

Bayer CropScience is submitting a notification for Ethrel brand Ethephon Plant Growth Regulator to remove the patent number from the label.

This notification is consistent with the provisions of PR Notice 98-10. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Section 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and I may be subjected to enforcement action and penalties under Section 12 and 14 of FIFRA".

Enclosed are an EPA Form 8570-1 and one copy of the revised Ethrel brand

Please contact me if you should have further questions.

Sincerely,

Bayer CropScience P. O. Box 12014 2. T. W. Alexander Drive Research Triangle Park, NC 27709

ειοι

Bayer CropScience

,

aura Phelps

Laura Phelps Regulatory Affairs, Herbicides & PGRs Bayer CropScience 2 T. W. Alexander Drive RTP, NC 27709 office: 919-549-2302 mobil:919-352-8075 email: laura.phelps@bayer.com



	0000 0000 0000 000 000 000 000 000 000
ΓίΓι ι ι ι ι ι	Ĺ
	ι ο ο ο ο ι ο ο ο ο ο ο ο ι
	ίζζε ί ι ι ίζι ί
	ι
	ί ζ ίιιίι ι
	ι ι ιιςιίι ι

ETHREL[®] brand Ethephon Plant Growth Regulator

For Pineapple and Sugarcane

FOR COMMERCIAL OR AGRICULTURAL USE ONLY. NOT FOR RESIDENTIAL USE

For Flower induction and Maturity Concentration in Pineapple

For Flower Prevention and Biomass Increase in Sugarcane (Hawaii Only)

ACTIVE-INGREDIENT:-

*This product contains 4 pounds ethephon per gallon.

EPA Reg. No. 264-257

EPA Est. No. 264-PA-01

KEEP OUT OF REACH OF CHILDREN PELIGRO DANGER

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID				
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 			
· · · · · · · · · · · · · · · · · · ·	Call a poison control center or doctor for treatment advice.			
IF SWALLOWED:	Immediately call a poison control center or doctor for treatment advice.			
	Do not induce vomiting unless told to do so by a poison control center or doctor.			
	Have person sip a glass of water if able to swallow.			
	Do not give anything by mouth to an unconscious person.			
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.			
	Rinse skin immediately with plenty of water for 15-20 minutes.			
	Call a poison control center or doctor for treatment advice.			
	For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.			
Have the p	roduct container or label with you when calling a poison control center or doctor or going for treatment.			

NOTE TO PHYSICIAN: Treat symptomatically. Consideration should be given to the possibility that overexposule to materials other than this product may have occurred. No specific antidote is available. Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

DANGER HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirts and long pants, chemical resistant gloves made of any waterproof material (such as Nitrile, Butyl, Neoprene and/or Barrier Laminate), shoes plus socks and protective eyewear.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as

1

NUTIFICATIONS

MAY 1 1 2011

specified in the WPS.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this products concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

SPRAY DRIFT

Avoid spray drift. Do not apply when weather conditions may cause drift. Do not allow this product to drift on to non-target areas. Drift may result in illegal residues or injury to adjacent crops and vegetation, in the form of leaf yellowing and defoliation. To avoid spray drift, DO NOT apply aerially when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet size will also reduce spray drift.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- 1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory below:

AERIAL DRIFT REDUCTION ADVISORY

[This section is advisory in nature and does not supersede the mandatory label requirements].

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater neight is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the

field, the applicator should compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

WIND

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide

regulations.

Read entire label before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is coveralls over long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof materials, such as nitrile, butyl, neoprene, or barrier laminate, chemical resistant footwear plus socks, and protective eyewear. For overhead exposure, chemical-resistant headgear is also required.

Notify workers of the application by warning them orally and posting warning signs at entrances to treated areas.

IMPORTANT

Do not use this product for purposes other than those listed on the label. Do not exceed the rate of ETHREL® prand Ethephon per acre per year recommended on this label.

Do not apply ETHREL® brand Ethephon through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide wastes are hazardous. Do not contaminate water, food, or feed by storage or disposal. Improper disposal of excess pesticide, spray-mixture, or-rinsate-is-a-violation-of-Federal-law.-If-these-wastes-cannot-be-disposed-of-by-use-according-to-label-instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by state and local authorities, by open burning. If burned, stay out of smoke.

GENERAL INFORMATION

When ETHREL® brand Ethephon Plant Growth Regulator for Pineapple and Sugarcane is applied to plants, the product decomposes within the plant tissue to release ethylene, a naturally occurring plant hormone.

PINEAPPLE FLOWER INDUCTION

Application of ETHREL® brand Ethephon will induce uniform flower initiation of pineapple plants.

PINEAPPLE MATURITY CONCENTRATION

Application of ETHREL® brand Ethephon will stimulate uniform shell color of pineapple fruit.

SUGARCANE FLOWER PREVENTION

A foliar spray of ETHREL® brand Ethephon will reduce or prevent flowering of sugarcane.

SUGARCANE BIOMASS INCREASE

ETHREL[®] brand Ethephon application to prevent flowering can result in increased biomass accumulation and increased recoverable sugars.

For recommendations on rates, spray volumes, and spray equipment under varying temperature and rainfall conditions consult your Extension specialist, Bayer CropScience Representative or farm agronomist for his experience with this product in your area.

SPRAY PREPARATION

Add 1/2 to 3/4 of the required amount of water to the spray tank. Start agitation. Add the required amount of ETHREL® brand Ethephon, and the remaining amount of water. Prepare only as much spray solution as can be used on the day of mixing. Use of previous day's spray mix may result in reduced activity.

Do not spill the concentrated product on spray equipment, or any airplane parts. ANY SPILLS SHOULD BE IMMEDIATELY RINSED WITH PLENTY OF WATER AS ETHREL[®] brand Ethephon IS CORROSIVE. Use of a nurse tank is highly recommended for avoiding possible spills of concentrated formulation on spray equipment.

EQUIPMENT CLEANING

Because of the acidic nature of this product, prolonged exposure to spray deposit will damage acrylic plastics, certain paints and metals.

Rinse thoroughly with detergent and water all exposed acrylic plastic-type materials (e.g. aircraft windshields), and painted surfaces within an hour after exposure to spray deposits.

At the end of each day, rinse thoroughly with detergent and water all metal parts of the aircraft and the associated spray equipment exposed to the spray deposits.

CROP AND USE	RATE PER ACRE	SPRAY VOLUME	APPLICATION TIMING	SPECIFIC COMMENTS	RESTRICTIONS
PINEAPPLE					
Flower Induction ETHREL [®] applied to pineapple plants will stimulate-uniform	1 to 2 quarts	100 to 500 gal/A	'Bearing age' about 12 months after planting. Generally about 6 months prior to desired harvest.	Proper rate will vary with local growing conditions, varieties, plantation management practices and time of year. Use the higher rate where earlier harvest is desired. Apply when pineapple foliage is	forage treated with ETHREL [®] Do not harvest
Maturity Concentration ETHREL [®] application will stimulate uniform shell color development	1/2 to 2 quarts	100 to 300 gal/A	When the first fruit begin to change color.	dry. Use a broadcast spray, thoroughly cover both foliage and fruit. Use the higher rate during periods of cool or cloudy weather when normal ripening has slowed.	
SUGARCANE					
Flower Prevention ETHREL [®] application will prevent or reduce flowering and pithy tissue formation.	1 pint	7 gal/A minimum	Just prior to flower initiation.	Apply by fixed wing or helicopter using equip- ment designed to give uniform coverage. Actual biomass increase	treated with ETHREL [®] sooner than 2 months after the last application. Do not graze sugarcane
Biomass Increase ETHREL [®] used to				will be affected by the time between treatment and harvest.	forage treated with ETHREL [®] .
prevent flowering can also result in increased biomass accumulation and recoverable sugar yield.				Consider anticipated re- coverable sugars as well as biomass to determine optimum harvest date.	For Use in Hawaii Only.

Ć

9/10

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 1 GALLON



Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

ETHREL® brand Ethephon Plant Growth Regulator for Pineapple and Sugarcane (MASTER) Approved 11/18/02, Notification 03/14/2011