

DIRECTIONS

FOLIAGE STEM METHOD

This is the standard method for high volume sprays along fence rows, highways and rights-of-way. Apply spray to both foliage and stems of all plants from the time foliage is fully developed to 11/1 and to plants dormant. **For effective control, all leaves, stems and suckers must be thoroughly wet to ground line.** Some regrowth may be expected in resistant species such as ash, maple, oak and poplar. Add 1 gallon WEEDONE* IBK WOODY PLANT HERBICIDE to 100 gallons of water. Mix thoroughly before spraying. Use 100 to 300 gallons of spray mixture per acre depending on density and height of brush.

BASAL BARK METHOD

Wet the base of all stems until the spray puddles out 5 to 6 inches and coats area to the root collar at ground line. This spray may be applied in any season. Apply WEEDONE* IBK WOODY PLANT HERBICIDE with the basal bark method or scattered brush cut as a second spray on species resistant to a first foliage spray. Mix 1 to 4 gallons of WEEDONE* IBK WOODY PLANT HERBICIDE in 100 gallons of oil. Apply with low volume knapsack sprayer or power equipment. Concentration will depend on species present, volume of spray used, and season applied.

MODIFIED BASAL METHOD

Drench the base of plants, then wet remaining stems and leaves to run-off spraying the lower 4/5 of the plants from bottom up. Treat when brush is in full foliage. This method may be used where susceptible species have been controlled by prior sprays and resistant species such as maple, oak, persimmon and ash remain. **Drenching the base of the plant and wetting all stems to run-off is essential for complete control.**

Early season spraying: Add 1 to 1 1/2 gallons of WEEDONE* IBK WOODY PLANT HERBICIDE to 10 gallons of diesel oil and mix thoroughly. Add this mixture to 89 gallons of water.

During dry weather or the latter part of spray season: Add 1 gallon of WEEDONE* IBK WOODY PLANT HERBICIDE to 15 gallons of diesel oil and mix thoroughly. Add this mixture to 83 gallons of water. Stir thoroughly before spraying, particularly if material is allowed to stand for more than one hour after mixing.

TO PREPARE A SPRAY: Add one half the required amount of oil or water to the spray tank, then add this product with agitation and finally the balance of the water or oil with continued agitation.

CAUTION: If this material is to be used in straight or mixture, do not let water get into it or the finished mixture.

NOTE: This material forms an emulsion in water, not a solution. This tends to separate on standing. Provide agitation to prevent such separation and ensure a uniform spray mixture.

RAILROAD ON-TRACK APPLICATION

Use 1 gallon of WEEDONE* IBK WOODY PLANT HERBICIDE and 1 quart of TRANSVERT* in each 100 gallons of water and apply 100 to 200 gallons per acre to control mixed brush growing along railroad rights-of-way and under railroad communication lines. Apply from spray train or Hi-Rail equipment only. Avoid contact with skin, eyes or clothing.

CUT SURFACE METHOD

Stump

This method is applicable at any time of year, but is most effective when applied as soon as possible after tree is cut. Spray the entire stump, particularly the exposed roots. **A thorough drenching is essential for complete control.**

Mix 1 to 4 gallons of WEEDONE* IBK WOODY PLANT HERBICIDE in 100 gallons of oil. Apply with low volume knapsack sprayer using solid stream nozzle of medium range.

Frit

Make a frit by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark but do not remove chips. This method is recommended for cut trees 5 to 6 inches in diameter and larger. Treat freshly cut frits at any time of the year.

Mix 1 to 4 gallons of WEEDONE* IBK WOODY PLANT HERBICIDE in 100 gallons of oil. Pour in as much of this mixture as the frit will hold without wasting it away.

WEEDONE* IBK WOODY PLANT HERBICIDE is a most effective formulation for controlling:

ALDER	ELM	RASPBERRY
ASH	GALLBERRY	SASSAFRAS
ASPEN	GROUND CHERRY	SPICEBUSH
BIRCH	HONEYSUCKLE	SUMAC
BLACKBERRY	HORSE CHESTNUT	SWEET GUM
BOX ELDER	KUDZU	SYCAMORE
BUCKBRUSH	LOCUST	VIRGINIA CREEPER
BUTTERNUT	OAK	WALNUT
CATALPA	OSAGE ORANGE	WILD CHERRY
CHESTNUT	PERSIMMON	WILD GRAPE
CHABAPPLE	PINE	WILD ROSE
DEWBERRY	POISON IVY	WILLOW
DOGWOOD	POISON OAK	
ELDERBERRY	POPLAR	

(and many other species)

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suitable for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks, and in the event of damage arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

© 1984 Amchem Products, Inc.

Printed in U.S.A.
1-800-368-7222

WEEDONE*

IBK WOODY PLANT HERBICIDE

Contains
of 2,4,5
woody
and also

CAUTION
on right side

ACTIVE INGREDIENTS
*2,4,5-Trichlorophenoxyacetic acid
**2,4-Dichlorophenoxyacetic acid

INERT INGREDIENTS
*2,4,5-Trichlorophenoxyacetic acid
or 2 pounds
**2,4-Dichlorophenoxyacetic acid
or 2 pounds



