

DIRECTIONS		
WEEDS IN CROPS	AMOUNT PER ACRE	DIRECTIONS
Spring Seeded Wheat, Barley, Rye	1/2 pint	For annual broadleaf weeds. Spray after grain is fully tillered (about 4 to 6 inches tall) but before it is in the boot stage.
	1 1/2 to 2 1/2 pints	For perennial broadleaf weeds. Spray after grain is fully tillered (about 4 to 6 inches tall) and when weeds are nearing the bud stage. Do not spray grain in the boot to dough stage.
Fall Seeded Wheat, Rye	1 1/2 to 2 1/2 pints	Use the lower rate where annual broadleaf weeds are the problem. Spray in the spring before grain is in the boot to dough stage. Do not forage or graze treated grain fields within two weeks after treatment with this product. Do not feed treated straw to livestock.
For Aerial Application	1/2 to 1 pint	For aerial application to small grains, pastures or rangeland, use the recommended amount of this product in 5 to 10 gallons of water per acre.


**WEEDS AND BRUSH IN FALLOW LAND, RANGELAND, PASTURES, ROADSIDES, FENCEROWS**

Annual Broadleaf Weeds	1 1/2 to 2 1/2 pints	Spray when weeds are young and growing vigorously. Controls: mustards, galinsoga, ragweed, lambsquarter, pigweed, sunflower, cocklebur.
Perennial Broadleaf Weeds	2 1/2 to 4 pints	Spray when weeds are actively growing and near the bud stage. Repeat applications may be needed for complete control. Controls: Canada thistle, bindweed, wild garlic, dandelion, plantain, chicory, dock.
Brush	2 1/2 to 4 pints per 100 gal water	Spray to thoroughly wet plants when they are in full leaf and growing actively. Where practical, cut tall woody plants and spray sucker growth when 2 to 4 feet tall. Retreatment may be needed for some species. Controls: willow, Virginia creeper, cherry rose, Japanese honeysuckle, wild grape.
Aerial Brush Spraying	1 1/2 to 2 1/2 pints in 1 gallon diesel oil 2 to 4 gallons water	Aerial spraying is a specialized job. Secure qualified technical guidance and employ a competent, reliable applicator. Become familiar with state laws governing the use of herbicides. Treat in spring when brush is fully leaved and growing actively. Controls: shinnery oak, sand sage, buckbrush, rabbitbrush, big sagebrush.

Do not graze treated areas by dairy animals within 7 days after treatment, or until chemical has disappeared from milk stage where grass seed production is desired. Do not graze susceptible grass pastures as injury may result.

**TO PREPARE A SPRAY:** Add one-half the required amount of product to one-half the water with continued agitation and finally the balance of water with continued agitation.

The above recommendations are general. Because of variable conditions, consult your local agricultural experiment station or extension service for specific recommendations.



# WEEDON

## LV6

EMULSIFIABLE BROADLEAF HERBICIDE

Weed and brush control in grains, rangeland, pasture roadsides and fencerows

**CAUTION:** Keep out of reach of children. See on right panel.

**ACTIVE INGREDIENT:**  
2,4-Dichlorophenoxyacetic acid, butoxyethyl ester  
**INERT INGREDIENTS:** .....  
\*Isomer specific by AOAC method No. 6.D01-5  
2,4-Dichlorophenoxyacetic acid equivalent 5.7 lb./gal.

MADE IN U.S.A.

**UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.**

AMBLER, PA 19002  
Clinton, IA ■ St. Joseph, MO. ■ Fremont, CA

70  
EPA Reg. No. 264-271AA EPA Est. 264-PA-1, 26  
Refer to the first letter of the lot number for packaging information.

# WEEDONE

## DEFOLIANT BROADLEAF HERBICIDE

and brush control in small  
fields, rangeland, pastures,  
road sides and fencerows

**WARNING:** Keep out of reach of children. See other cautions on label.

**INGREDIENT:**  
2,4-Dichlorophenoxyacetic acid, butoxyethyl ester\* ..... 83.5%  
**OTHER INGREDIENTS:** ..... 16.5%  
\*As determined by AOAC method No. 6.D01-5  
2,4-Dichlorophenoxyacetic acid equivalent 5.7 lb./gal. or 57.5% wt.

MADE IN U.S.A.

## AMMONIUM CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.

AMBLER, PA 19002

Clinton, IA. ■ St. Joseph, MO. ■ Fremont, CA

EPA Reg. No. 264-271AA EPA Est. 264-PA-1-264-MO-1-264-CA-1  
Refer to the first letter of the lot number for packaging location

## CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid inhalation of spray mist.

Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, tobacco and most ornamentals as injury may result. Coarse sprays are less likely to drift. Do not use same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides. Do not use in or near a greenhouse. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

Do not contaminate any body of water by washing of spray equipment or disposal of waste.

Do not contaminate water intended for irrigation or domestic purposes.

Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands away from water supplies and not accessible to livestock. Do not reuse empty containers. Destroy in a safe place. Do not burn.

**Do not use this product for purposes other than those recommended on the label.**

## WARRANTY

The manufacturer warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. The manufacturer, therefore, makes no other express or implied warranty, and no agent of the manufacturer is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damage arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of the manufacturer's liability.

Form No. 54191 (Rev. 4-8-81)

Printed in U.S.A.