

264-271 pm23 10614

WEEDONE® LV6  
EMULSIFIABLE BROADLEAF HERBICIDE

Weed and brush control in small grains, field corn, rangeland, pastures,  
roadsides and fencerows

ACTIVE INGREDIENT:

2,4-Dichlorophenoxyacetic acid, butoxyethyl ester\* . . . . . 83.5%


INERT INGREDIENTS: . . . . . 16.5%

\*Isomer specific by AOAC method No. 6.D01-5

2,4-Dichlorophenoxyacetic acid equivalent 5.7 lb./gal. or 5.5% wt.

E.P.A. Reg. No. 264-271AA

E.P.A. Est. No. 264-271



KEEP OUT OF REACH OF CHILDREN

CAUTION


Precautionary Statements: See Inside Booklet

IN CASE OF EMERGENCY TELEPHONE 1-800-822-4357  
(24 HOURS A DAY) IN U.S.A.  
FOR GENERAL PRODUCT INFORMATION CALL 1-800-334-9745

RHONE POULENC AG PRODUCTS COMPANY  
P. O. Box 12014, T.W. Alexander Drive  
Research Triangle Park, N.C. 27709

WEEDONE is a registered trademark of RHONE POULENC NEDERLAND B.V.

Made in U.S.A.



PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing vapors or spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady gentle stream of water for at least 15 minutes. GET MEDICAL ATTENTION, preferable an ophthalmologist.

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. GET MEDICAL ATTENTION.

IF ON SKIN: Immediately wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen, if necessary. GET MEDICAL ATTENTION.

NOTE TO PHYSICIAN

No specific antidote is available.

All treatments should be based on observed signs and symptoms of distress in the patient. Over-exposure to materials other than this product may have occurred.

This product contains petroleum distillates. If large amounts (greater than 1 ml/kg body weight) of the product have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, was stomach by instilling 30-50 g. of activated charcoal in 3-4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.


ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

USE PRECAUTIONS

Do not apply when weather conditions favor drift from treated areas.

Do not use the same spray equipment for other purposes unless thoroughly cleaned.



Do not use in or near a greenhouse.

Do not apply WEEDONE® LV6 through any type of irrigation system.

Do not contaminate water used for irrigation or domestic purposes.


Do not apply WEEDONE® LV6 Emulsifiable Broadleaf Herbicide directly to, or permit spray mist to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by WEEDONE® LV6 Emulsifiable Broadleaf Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion, consult with local weather services before making an application.

Use coarse sprays to minimize drift. Do not apply with hollow cone type insecticide or other nozzles that produce fine spray droplets.

Drift from aerial or ground application may be reduced by: (1) applying as near to the target as possible in order to obtain coverage; (2) by increasing the volume of spray mix per acre; (3) by decreasing the pounds of pressure at the nozzle tips; (4) by using nozzles which produce a coarse spray pattern; (5) by not applying when wind is blowing toward susceptible valuable plants.

At high air or ground surface temperatures, vapors from this product may injure susceptible plants.



**DIRECTIONS FOR USE**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

**GENERAL INFORMATION**

**WEEDONE LV6 Emulsifiable Broadleaf Herbicide** contains the butoxyethyl ester of 2,4-D, the original and one of the most volatile esters. In cropland, WEEDONE LV6 is more effective than amines for controlling hard-to-kill weeds such as thistles, smartweeds, wild garlic, curled dock, tansy ragwort, and wild onions.

**TO PREPARE A SPRAY:** Add one-half the required amount of water to the spray tank, then add the WEEDONE LV6 with agitation. Finally, the balance of the water with continued agitation. This material forms an emulsion in water--not a solution to separate on standing. Provide agitation to prevent such separation and insure uniform spray mixtures.

**COMPATIBILITY:** If WEEDONE LV6 is to be tank mixed with fertilizers or with other pesticides, compatibility should be tested before mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not mix for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read the label of each tank-mix product used for precautionary statements, directions for use, geographic and other restrictions.

**COMBINATION WITH LIQUID NITROGEN FERTILIZER:** Use 1/3-1/2 pint WEEDONE LV6 per acre for weeding and feeding corn, grass pastures as directed on this label. Use the fertilizer at rates recommended by supplier or Extension Service. Fill the spray tank about half full with the liquid fertilizer, then add the WEEDONE LV6 with vigorous agitation, filling the tank with fertilizer. Apply immediately and continue agitation in the spray tank during application. Spraying during very cold weather (near freezing) is not advisable. Do not allow mixture to stand overnight. Incompatibility has been encountered with some fertilizer brands or under some environmental conditions. If in doubt, test a small sample before application. NOTE: Fertilizers can increase foliage contact burn of herbicides. Reducing the rate and concentration will reduce the hazard of leaf burn.

Weeds controlled by properly applied treatments of WEEDONE LV6 include the following:

Annual and biennial weeds

annual yellow sweetclover  
\*beggarticks  
bull thistle  
coffeeweed  
common evening primrose  
\*common broomweed  
common burdock  
common cocklebur  
common lambsquarters  
croton (Texas or woolly)  
hairy galinsoga  
Jimsonweed

\*knotweed  
\*mallow (venice, dwarf, little)  
marshelder  
morning glory (common, ivy, woolly)  
mustards (except blue mustard)  
pepperweeds (except perennial)  
\*\*pigweeds (Amaranthus spp.)  
\*prickly lettuce  
ragweed (common, giant)  
rough fleabane  
\*Russian thistle

salsify (western or common)  
\*smartweeds (annual species)  
sowthistle (annual or spiny)  
sunflower  
velvetleaf  
\*vervains  
vetches  
\*wild carrot  
wild lettuce  
wild parsnips  
wild radish

Perennial weeds

\*alfalfa  
\*blindweed (hedge, field and European)  
blue lettuce  
\*Canada thistle  
catnip  
chicory  
dandelion  
\*docks  
\*dogbanes

\*goldenrod  
\*ground ivy  
\*hawkweed (orange)  
healall  
\*hoary cress  
Jerusalem-artichoke  
\*many-flowered aster  
\*nettles (including stinging)  
plantains

sowthistle (perennial)  
\*tansy ragwort  
\*vervains  
\*western ironweed  
\*wild garlic  
\*wild onion

\* These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for application.

\*\* Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply WEEDONE LV6 only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

RECOMMENDATIONS FOR APPLYING WEEDONE LV6 EMULSIFIABLE BROADLEAF HERBICIDE

WEEDS IN CROPS

AMOUNT OF WEEDONE LV6  
PER ACRE

DIRECTIONS

CEREAL GRAINS

Postemergence

WHEAT, BARLEY, RYE  
(not underseeded with legumes)  
Annual and biennial weeds

1/3 - 1 1/3 pints\*

Perennial broadleaf weeds

2/3 - 1 1/3 pints\*

Apply after grain is fully tillered (usually about 4-8" high) but not forming joints in the stem. Do not spray grain in boot to dough stage.

RECOMMENDATIONS FOR APPLYING WEEDONE LV6 EMULSIFIABLE BROADLEAF HERBICIDE

WEEDS IN CROPS

AMOUNT OF WEEDONE LV6 PER ACRE	DIRECTIONS
Wild onion or garlic	1-1 1/3 pints*
Emergency Weed Control in Wheat Perennial broadleaf weeds	2 pints*
Preharvest	2/3-1 1/3 pints*

Apply 1 pint when grain is fully tillered and wild garlic or onion plants are small. Apply 1 1/3 pints after harvest in the crop stubble. For control of new fall growth of wild onion or garlic refer to fallowland use directions.

Apply when weeds are approaching bud stage, but do not spray grain during the boot to dough stage. The 2 pints per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.

Apply when grains are in the hard dough stage to suppress large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to induce succulent weed growth.

\* Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage. Do not apply to grain in the seedling stage.

RESTRICTIONS AND LIMITATIONS FOR USE ON SMALL GRAINS:

- For aerial application on grain, apply WEEDONE LV6 in 2 or more gallons of water per acre. For ground application, use a minimum of 10 gallons of water per acre.
- Do not mix with oil for crop uses.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Do not feed treated straw to livestock if a preharvest treatment or emergency treatment as described above is applied.

Handwritten mark: a stylized signature or initials.

RECOMMENDATIONS FOR APPLYING WEEDONE LV6 EMULSIFIABLE BROADLEAF HERBICIDE

WEEDS IN CROPS

	AMOUNT OF WEEDONE LV6 PER ACRE	DIRECTIONS
<b>FIELD CORN</b>		
Preplant	2/3-1 1/3 pints	To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn. Apply 7 to 14 days before planting. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth. Use high rate for control of less susceptible weeds or cover crops such as alfalfa.
Preemergence	1 1/3-2 pints	To control broadleaf weeds and suppress annual grasses, apply 3 to 5 days after planting, but before corn emerges. Use high rate on soil high in organic matter. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth.
Postemergence		
Annual broadleaf weeds	1/3 pint	Avoid spraying just after corn leaves unfold, as injury may occur. Apply when weeds are small and corn is less than 8" tall. If corn more than 8" tall must be sprayed, use drop pipes.
Perennial broadleaf weeds	1/2 pint	Spray when weeds are in the bud to bloom stage. If corn is more than 8" tall, use drop pipes to keep spray off corn leaves. <u>Do not spray when corn is in tassel to dough stage.</u>  2,4-D may make corn brittle. Winds or cultivation may cause stalk breakage while brittle.
Preharvest	2/3-1 1/3 pints	After the hard dough (denting) stage, to suppress weeds that interfere with harvest such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, and velvetleaf, and to decrease production of weed seeds, spray with air or ground equipment. The high rate will be needed for tough weeds under stress.

RESTRICTIONS AND LIMITATIONS FOR USE ON FIELD CORN:

- Do not forage or feed corn fodder for 7 days following application.

0468G/PENDING

7/19/52

RECOMMENDATIONS FOR APPLYING WEEDONE LV6 EMULSIFIABLE BROADLEAF HERBICIDE

WEEDS IN CROPS

	AMOUNT OF WEEDONE LV6 PER ACRE	DIRECTIONS
GRAIN SORGHUM (milo)		
Annual broadleaf weeds	1/2 pint	Treat when sorghum is 6 to 15" tall (to top of canopy). If sorghum is taller than 8" to top of canopy, use drop nozzles to keep spray off crop foliage. <u>Do not treat during the boot, flowering, or early dough stages.</u>
Perennial broadleaf weeds	3/4 pint	

RESTRICTIONS AND LIMITATIONS FOR USE ON SORGHUM:

- Do not forage or feed fodder for 7 days following application.
- A minimum application volume of 5 gallons per acre by air or 10 gallons per acre by ground is recommended.

SUGARCANE		
Preemergence	1 1/3 pints	Apply before canes appear for control of emerged broadleaf weeds. Apply after cane emerges and through layby. DO NOT USE IN CALIFORNIA
Postemergence	1 1/3-2 2/3 pints	

WEEDS AND BRUSH IN PASTURES, RANGELAND, AND FALLOW LAND

	AMOUNT OF WEEDONE LV6 PER ACRE	DIRECTIONS
ESTABLISHED GRASS PASTURES AND RANGELAND		
Annual broadleaf weeds	1 1/3 pints	Apply when weeds are small and actively growing and prior to bud stage. Do not apply to newly seeded areas until grass is well established. Do not apply when grass is in the boot to milk stage if grass seed production is desired. Bittergrass and legumes may be injured by this treatment.
Perennial and biennial broadleaf weeds	2-2 2/3 pints	

2-2-14



ESTABLISHED GRASS PASTURES AND RANGELAND

Sand sage, sagebrush,  
shinnery oak, and other  
2,4-D susceptible woody  
species

2/3-2 2/3 quarts

Apply aerially to brush in 2 or more gallons of water per acre. One gallon fuel oil may be included in the mixture. Consult state or local brush control specialists for most effective rate, volume and timing of spray applications.

FALLOW LAND AND CROP STUBBLE

Annual broadleaf weeds

2/3-1 1/3 pints

Use the lower rate when weeds are small (2-3" tall) and growing actively. Use the higher rate on older and drought-stressed plants.

Biennial broadleaf weeds

2/3-1 2/3 pints

Spray while musk thistles or other biennial species are in the seedling to rosette stage, and before flower stalks are initiated. The lower rate can be used in spring during rosette stage. In fall or after flower stalks have developed, use the highest rate.

Perennial broadleaf weeds

1 1/3-4 pints

Spray weeds in bud to bloom stage, or in good vegetative growth. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead.

Wild garlic and onion in  
crop stubble

2 2/3-4 pints

Apply to new regrowth of wild onion or garlic which occurs in the fall following harvest of small grains, soybeans, corn, or grain sorghum.

RESTRICTIONS AND LIMITATIONS FOR USE ON PASTURES, RANGELAND, AND FALLOW LAND:

- For ground application, use a minimum of 10 gallons of water per acre. For aerial application, use a minimum of 2 gallons of water per acre.
- Do not graze animals on treated areas within 7 days after treatment.
- Do not plant treated fallow land until three months after treatment or until chemical has disappeared from soil.



41206

RECOMMENDATIONS FOR APPLYING WEEDONE LV6 EMULSIFIABLE BROADLEAF HERBICIDE

FENCEROWS, ROADSIDES, DRAINAGE DITCHBANKS, VACANT LOTS, AIRFIELDS, RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY, AND OTHER NON-CROP AREAS.

	AMOUNT OF WEEDONE LV6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1 1/3-2 2/3 pints	Apply when most weeds are still young and growing vigorously.
Perennial and biennial broadleaf weeds	2-4 pints	Apply when weeds are actively growing and near the bud stage, but before flowering. For best results on tansy ragwort and musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on thistle, nettle, and bindweed. Treat wild onion or garlic in early spring and in fall when they are young and growing actively.
Cattails and tules	1-1 2/3 gallons	Mix 1 1/3 quarts WEEDONE LV6 in 2 quarts kerosene or diesel oil then add this mixture to 100 gallons of water. Apply 300 to 500 gallons of spray per acre, depending on the stand. The addition of a wetting agent such as SURFEL Spray Adjuvant is suggested.
Woody plants	2/3-1 1/3 gallons	Apply in 30-100 gallons of spray per acre, preferably through low-volume equipment such as DIRECTA-SPRA, WOBBLER, MINI-WOBBLER, OR SPIROMETER. Treat when 2,4-D-susceptible species are in full leaf and growing actively. Repeated applications, oil or wetting agent, and/or high volume spraying (300 to 500 gallons of spray per acre) may be needed for acceptable control of some species or if brush growth is dense.

RESTRICTIONS AND LIMITATIONS FOR USE IN NON-CROP AREAS:

- For aerial application to solid stands of susceptible brush, use 1 1/3-2 2/3 quarts in 3-12 gallons volume per acre. Two to 4 quarts of fuel oil may be included in this mixture.
- Applications will be less effective when conditions such as deficient soil moisture reduce brush growth.

11901

Read and follow the label of each tank-mix product used for precautionary statements, directions for use, geographic and other restrictions.

WEEDONE LV6 and GARLON 4  
or GARLON 3A Tank Mixtures  
for non-crop areas.  
Broadleaf Weed Control

1 1/3 -  
2 2/3 pints  
WEEDONE LV6  
Plus  
2-6 pints  
GARLON-4  
or  
3-8 pints  
GARLON-3A

For wider spectrum control of broadleaf weeds and woody plants.

Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre.  
Apply when broadleaf weeds are actively growing.

Woody Plant Control  
Broadcast Foliar Spray

2/3 -  
1 1/3 gallons  
WEEDONE LV6  
Plus  
1 1/2 - 3 quarts  
GARLON-4  
or  
2-4 quarts  
GARLON-3A

Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when woody plants are actively growing.

Woody Plant Control  
High Volume Leaf-Stem  
Treatment with Ground  
Equipment


2/3 -  
5 1/3 quarts  
WEEDONE LV6  
Plus  
1 1/2 to  
12 pints GARLON-4  
or  
2 to 16 pints  
GARLON-3A

Mix 2/3 to 1 1/3 to 2 quarts WEEDONE LV6 plus 1 1/2 to 3 pints GARLON 4 or 2 to 4 pints GARLON 3A in enough water to make 100 gallons of spray. Apply at a volume of 100 to 400 gallons of total spray per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems and root collars of plants to be controlled

Woody Plant Control  
Aerial Application  
(Helicopter only)

2/3 -  
1 1/3 gallons  
WEEDONE LV6  
plus  
3-4 quarts  
GARLON-4  
or  
4-6 quarts  
GARLON-3A

Apply in a total spray volume of 10 to 30 gallons per acre using drift control equipment such as the MICROFOIL boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.



11/20/11

WEEDONE LV6 and BANVEL  
Herbicide tank mixtures for  
non-crop areas.

Annual Broadleaf Weeds

1 1/3 -  
2 2/3 pints  
WEEDONE LV6  
plus  
1/2 - 1 1/2  
pints BANVEL

For wider spectrum control of broadleaf weeds and woody plants.

Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth.

Perennial and biennial  
broadleaf weeds

2-4 pints  
WEEDONE LV6  
plus  
1/2-6 pints  
BANVEL

Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches Rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3 inch rosette stage.

Woody Plant Control  
Broadcast, high volume,  
stem foliage or aerial  
application

2/3 -  
1 1/3 gallons  
WEEDONE LV6  
plus  
2-8 quarts  
BANVEL

Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons per acre) or apply aerially in enough water to deliver total spray volume of 10 to 30 gallons per acre using drift control equipment such as the MICROFOIL Boom or an effective drift control agent such as LO-DRIFT spray additive. Use the higher rates and volumes when plants are dense or under drought conditions.

---

SMALL AREA APPLICATIONS FOR NON-CROP USE

---


For control of broadleaf weeds in small non-crop areas with hand held or back-pack sprayers mix 2 2/3 fluid ounces WEEDONE LV6 per gallon of water. Thoroughly wet all weed foliage. Maintain agitation of mixture to prevent separation.

---

WEEDONE, SURFEL, DIRECTA-SPRA, WOBBLER, MINI WOBBLER, SPIROMETER, and LO-DRIFT are registered trademarks of Rhone-Poulenc Ag Company.

GARLON is a trademark of the DOW Chemical Co.

BANVEL is a trademark of Velsicol Chemical Corporation



12814


Fungicides or other pesticides.

#### PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



14 of 14

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYERS'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

