


DIRECTIONS FOR USE



FOLIAGE SPRAY - Spray WEEDONE® Super BK 32 herbicide after brush foliage is fully developed in the spring until plants begin to go dormant in the fall. All leaves, stems and suckers should be wet thoroughly to ground line. Some regrowth may be expected on resistant species such as ash, blackberry, maple, oak and persimmon. Treat any regrowth the following season.

BASAL SPRAY - Wet the lower 18 inches of all stems until material runs down in all sides and collects around the root collar at ground line. This treatment can be applied in any season.

STUMP SPRAY - To prevent stumps from resprouting, spray the entire stump, particularly bark and exposed roots. Spray stumps any time during the year, but as soon as possible after trees are cut. A thorough drenching is essential for complete control.

CONCENTRATE STUMP TREATMENT - For small (up to 3-inch diameter) stems, cut them as close to the ground as possible and apply undiluted WEEDONE Super BK 32 herbicide directly from the can to the surface of the freshly cut stump.

FRILL TREATMENT - This method is recommended for trees 5 to 6 inches in diameter and larger. Make frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark but do not remove chips. Treat freshly cut frills at any time during the year with 1 pint of WEEDONE Super BK 32 per 2 gallons of water. Pour in as much of this mixture as the frill will hold without wasting chemical.

AMOUNT TO USE KNAPSACK SPRAYERS Mix thoroughly before using.

Method	WEEDONE Super BK 32	Diluent
Foliage spray	4 ounces (1/2 cup) 1/2 pint	2-1/2 gallons water 5 gallons water
Basal spray	1 pint	2-1/2 gallons diesel oil or kerosene
Stump spray	1 pint	2-1/2 gallons diesel oil or kerosene

WEEDONE Super BK 32 woody plant herbicide is effective for controlling:

alder	fir	sassafras
ash*	honeysuckle	spicebush
aspen	kudzuvine	spruce
birch	maples*	sumac
blackberry	oaks	sweetgum
black cherry	osage orange	sycamore
black locust	persimmon*	tulip poplar
boxelder	pine	virginia creeper
buckbrush	poison ivy	wild cherry
dewberry	poison oak	wild grape
dogwood	poplar	wild rose
elderberry	raspberry	willow
elms	salmonberry	(and many other similar species)

* controlled by stump or basal treatment only.

WEEDONE Super BK 32 woody plant herbicide is not recommended for weed control in crops.

When diluting, the material must be mixed thoroughly before using. If the material is allowed to stand for any length of time after dilution, it should be stirred again before using.

WARRANTY

The manufacturer warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. The manufacturer, therefore, makes no other express or implied warranty, and no agent of the manufacturer is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of the manufacturer's liability.

Form No. 4982 12-79

Printed in U.S.A.

BEST DOCUMENT AVAILABLE

NET CONTENTS

1 - GAL.

WEEDONE

SUPER BK 32

WOODY PLANT HERBICIDE

Controls unwanted trees, shrubs and vines in fencerows, roadsides, vacant lots and similar non-crop areas.

CAUTION: Keep out of reach of children.
See other cautions on side panel.

ACTIVE INGREDIENTS:

2 - (2,4 - Dichlorophenoxy) propionic acid butoxyethyl ester*16.4%
2,4 - Dichlorophenoxyacetic acid, butoxyethyl ester**16.7%

INERT INGREDIENTS:66.9%

*2, (2,4-Dichlorophenoxy) propionic acid equivalent 11.5% by weight or 0.95 pounds per gallon
**2,4 - Dichlorophenoxyacetic acid equivalent 11.5% by weight or 0.95 pounds per gallon

**isomer specific by A.O.A.C. method 6.DO1-5



AGRICULTURAL PRODUCTS COMPANY, INC.
AMBLER, PA. 19002

Clinton, IA St. Joseph, MO Fremont, CA

EPA Reg. No. 264-307

(A) (S) (N)

EPA Est. No. 264-PA-1, 264-MO-1, 264-CA-1.

Refer to the first letter of the first number for packaging location.

PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals

CAUTION

Harmful if swallowed or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of contact immediately flush skin or eyes with plenty of water. Get medical attention if irritation persists. Contains petroleum distillates; if swallowed, do not induce vomiting. Get immediate medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate any body of water by washing of spray equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, tobacco, and ornamentals as this product may cause injury. Coarse sprays are less likely to drift. Do not use same spray equipment for other purposes.

Under very high temperature of use, vapors from this product may injure susceptible plants in the immediate vicinity.

Do not contaminate water used for irrigation or domestic purposes. Do not use in or near a greenhouse.

Do not use this product for purposes other than those recommended on this label.

STORAGE AND DISPOSAL STATEMENTS

Do not reuse container. Destroy by burying in a safe place away from water supplies or wrap and discard in trash. Do not burn. Do not store near fertilizers, seeds, insecticides or fungicides.

BEST DOCUMENT AVAILABLE