


### RHÔNE-POULENC AG COMPANY

## 2,4-DP Registration Standard Labeling Requirements Certification of Compliance Statement

I, being an authorized representative of Rhône-Poulenc Ag Company understand the labeling requirements specified in the Registration Standard for 2,4-Dichlorophenoxypropionic Acid and its inorganic salts, amines and esters, as amended and included therein by reference. Further, I certify that all containers of WEEDONE® Super BK32 Woody Plant Herbicide, EPA Reg. No. 264-307, released for shipment after March 1, 1991, will bear revised labeling in accordance with those requirements.

Company Representative Signature: Karen S. Shearen

Date: No Vember 5, 1990



## WEEDONE SUPER BK-32

**Woody Plant Herbicide** 

Controls unwanted trees, shrubs, and vines in fencerows, roadsides, vacant lots and similar non-crop areas.

**ACTIVE INGREDIENTS** 

\*2-(2,4-Dichlorophenoxy)propionic acid equivalent 11.5% by weight or 0.95 pounds per gallon.

\*\*2,4-Dichlorophenoxyacetic acid equivalent 11.5% by weight or 0.95 pounds per gallon.

\*Isomer specific by AOAC Method 6.DO1-5

EPA Reg. No. 264-307

EPA Est. No.

# KEEP OUT OF REACH OF CHILDREN CAUTION

For <u>PRODUCT USE</u> information Call 1-800-334-9745
For <u>MEDICAL</u> and <u>TRANSPORTATION Emergencies</u> <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577

## STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses to drink. Do not induce vomiting. Get medical attention. IF ON SKIN: Wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

#### NOTE TO PHYSICIAN

No specific antidote is available. All treatments should be based upon observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains petroleum distillates. If large amounts (greater than 1 ml/kg body weight) of the product have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30-50 grams of activated charcoal in 3-4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.

## **PRECAUTIONARY STATEMENTS**

#### CAUTION

## HAZARD TO HUMANS AND DOMESTIC ANIMALS

'armful if swallowed, absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. When undling this product, wear chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label. Wash thoroughly with soap and water after handling.

## **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect fish, aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, tobacco, and ornamentals as this product may cause injury. Coarse sprays are less likely to drift. Under very high temperature of use, vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation or domestic purposes. Do not use in or near a greenhouse.

CLEANING OF EQUIPMENT: Do not use same spray equipment for other purposes unless thoroughly cleaned. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water on the ground; spray or drain over a large area away from wells and other water or drain over a large area.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

## STORAGE AND DISPOSAL

#### STORAGE

Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE AND CONTAINER DISPOSAL

Do not reuse container. Securely wrap original container (bottle, can, jar or plastic jug) in several layers of newspaper and discard in trash.

#### **GENERAL INFORMATION**

WEEDONE® SUPER BK-32 Woody Plant Herbicide is not recommended for weed control in crops.

When diluting, the material must be mixed thoroughly before using. If the material is allowed to stand for any length of time after dilution, it should be stirred again before using.

## **GENERAL WEED LIST**

WEEDONE SUPER BK-32 WOODY PLANT HERBICIDE IS EFFECTIVE FOR CONTROLLING:

dogwood alder ash\* elderberry **aspen** elms bear clover fir birch gooseberry blackberry honevsuckle black cherry interior live oak black locust kudzuvine black oak madrone oxelder manzanita Juckbrush. manies\* canyon live oak oaks ceanothus osage orange chamise persimmon\* chinquapin pine coffeeberry poison ivv currant poison oak dewberry poplar

rasoberry salmonberry sassafras serviceberry spicebush SDIUCE sumac sweetgum sycamore tanoak tulio poplar Virginia creeper wild cherry wild grape wild rose willow yerba santa

(And many other similar species)

#### FOLIAGE SPRAY

Spray WEEDONE® SUPER BK-32 Woody Plant Herbicide after brush foliage is fully developed in the spring until plants begin to go dormant in the fall. All leaves, stems and suckers should be wet thoroughly to ground line. Some regrowth may be expected on resistant species such as ash, blackberry, maple, oak and persimmon. Treat any regrowth the following season.

## **BASAL SPRAY**

Wet the lower 18 inches of all stems until material runs down in all sides and collects around the root collar at ground line. This treatment  $\gamma$  in be applied in any season.

#### STUMP SPRAY

To prevent stumps from resprouting, spray the entire stump, particularly bark and exposed roots. Spray stumps any time during the year, but as soon as possible after trees are cut. A thorough drenching is essential for complete control.

## CONCENTRATE STUMP TREATMENT

For small (up to 3 inch diameter) stems, cut them as close to the ground as possible and apply undiluted WEEDONE® SUPER BK-32 directly from the can to the surface of the freshly cut stump.

### FRILL TREATMENT

This method is recommended for trees 5 to 6 inches in diameter and larger. Make frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark but do not remove chips. Treat freshly cut frills at any time during the year with 1 pint of WEEDONE® SUPER BK-3.2 Herbicide per 2 gallons of water. Pour in as much of this mixture as the frill will hold without wasting chemical.

Controlled by stump or basal treatment only.

Knapsack Sprayers

METHOD	WEEDONE® SUPER BK-32	DILUENT	
Foliage spray	4 Ounces (1/2 cup)	2 1/2 Gals. Water	
	1/2 Pint	5 Gais. Water	
Basal spray	1 Pint	2 1/2 Gals. Diesel Oil or Kerosene	
Stump spray	1 Pint	2 1/2 Gals. Diesel Oil or Kerosene	

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LÆBEL

DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

WEEDONE® is a registered trademark of Rhône-Poulenc ©1990 Rhône-Poulenc Ag Company Made in U.S.A.

EPA Submission: 11/5/90