




RHÔNE-POULENC AG COMPANY

2,4-DP Registration Standard  
Labeling Requirements  
Certification of Compliance Statement

I, being an authorized representative of Rhône-Poulenc Ag Company understand the labeling requirements specified in the Registration Standard for 2,4-Dichlorophenoxypropionic Acid and its inorganic salts, amines and esters, as amended and included therein by reference. Further, I certify that all containers of WEEDONE® DP Herbicide, EPA Reg. No. 264-308, released for shipment after March 1, 1991, will bear revised labeling in accordance with those requirements.

Company Representative Signature: Karen S. Shearer

Date: November 5, 1990



274

# WEEDONE® DP Herbicide

## FOR TURF WEED CONTROL

### ACTIVE INGREDIENTS:

\*2,4-Dichlorophenoxyacetic acid, butoxyethyl ester.....16.7%  
 \*\*2-(2,4-Dichlorophenoxy)propionic acid, butoxyethyl ester.....16.4%

### INERT INGREDIENTS:

\* 2,4-Dichlorophenoxyacetic acid equivalent 11.5% by weight or 0.95% pounds per gallon  
 \*\*2-(2,4-Dichlorophenoxy)propionic acid equivalent 11.5% by weight or 0.95 pounds per gallon  
 \*Isomer specific AOAC Method #6 D01-5.....66.9%

EPA Reg. No. 264-308

EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN CAUTION

For PRODUCT USE Information Call 1-800-334-9745.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. Get medical attention.  
**On Skin:** Wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.  
**If in Eyes:** Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.  
**If Inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

#### NOTE TO PHYSICIAN

No specific antidote is available. All treatments should be based upon observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred. This product contains petroleum distillates. If large amounts (greater than 1 ml/kg body weight) of the product have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30-50 grams of activated charcoal in 3-4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.

### PRECAUTIONARY STATEMENTS

#### CAUTION

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. When handling this product, wear chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instruction on this label. Wash thoroughly with soap and water after handling.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect fish, aquatic invertebrates and non-target plants. Do not apply to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

**IMPORTANT:** Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, tobacco, and ornamentals as this product may cause injury. Coarse sprays are less likely to drift. Do not contaminate water used for irrigation or domestic purposes. Do not use in or near a greenhouse.

**CLEANING OF EQUIPMENT:** Do not use same spray equipment for other purposes unless thoroughly clean. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

### STORAGE & DISPOSAL

#### STORAGE

Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

#### PESTICIDE AND CONTAINER DISPOSAL

Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

## GENERAL CAUTIONS AND RESTRICTIONS

**IMPORTANT:** Do not use WEEDONE® DP Herbicide for controlling weeds in flower or vegetable beds, in shrub or ornamental plantings, or on lippia or dichondra lawns. No weed spraying should be attempted during drought periods unless the lawn is watered thoroughly several days before treatment. A separate sprayer should be kept for killing weeds because 2,4-D and related chemicals are difficult to clean from equipment. Spray when the air is clam to avoid spray drift that might injure nearby desirable plants. Do not use on bentgrass greens or tees.

## MIXING SPRAYS

Fill tank of sprayer halfway with water. Pour in required amount of WEEDONE® DP Herbicide. Mix well. Add remainder of water and mix again. If the spray mixture is allowed to stand several hours, mix again before spraying.

## GENERAL INFORMATION

### USE ON LAWNS, PARKS, GOLF COURSES, ATHLETIC FIELDS OR INDUSTRIAL TURF AREAS

Apply WEEDONE® DP Herbicide any time broadleaf weeds are growing actively. Dandelion, plantain, wood sorrel and clovers are best treated in the fall or in spring before flower heads develop.

Winter weeds such as chickweeds and henbit should be treated in early spring.

Summer weeds such as oxalis, knotweed and spotted spurge should be sprayed when they are small. In areas with extended growing seasons, such as California, treatment in both spring and fall may be needed to control more resistant species.

Lawns to be over-seeded in the fall should be treated at least 4 weeks before the planned seeding date. A good rain (1/4 inch or more) or deep watering should occur during this period. Fall seeded lawns may be treated the following spring.

Spring-seeded lawns may be treated after the grasses have sprouted and been cut at least twice, generally 6 to 10 weeks after seeding, depending on germination and growth rate.

For established lawns, fall treatment fits into a good turf management program. Proper fertilization and mowing should be combined with chemical weed control to thicken the turf after weeds have died and to discourage more weeds from invading.

## GENERAL WEED LIST

### COMMON LAWN WEEDS

black medic	dandelion	knotweed
buttercup	*English daisy	little starwort
chickweeds (common, mouseear)	ground ivy	oxalis (wood sorrel)
clovers	healall	plantains (narrow or buckhorn; broadleaf)
cudweed	henbit	soliva
		*spurge

\*Treat in spring and again in fall

### OTHER WEEDS FOUND IN LAWNS

bindweed	kochia	vervain
dock	lambquarters	vetch
evening primrose	mallow	viclet
false dandelion	pigweed	wild carrot
fleabane	poorjoe	wild geranium
Florida pusley	ragweed	wild lettuce

This product is not effective on perennial veronics or on weed grasses.

## AMOUNTS TO USE

### Pressure Sprayers

WEEDONE® DP	AREA COVERED	WATER
3 Fl. Ozs. (6 Tablespoons)	1,000 Sq. Ft.	1 Gallon
1 Pint	5,000 Sq. Ft.	10 Gallons
1 Quart	10,000 Sq. Ft.	20 Gallons
1 Gallon	40,000 Sq. Ft.	80 Gallons

## HOSE ATTACHED SPRAYERS

### Lawn Models

Use 3 tablespoons of WEEDONE® DP Herbicide diluted with water to the 3 to 5 gallon mark on the sprayer jar for each 500 square feet (25' x 20') of lawn area to be sprayed. Adjust water pressure so that spray streams about 10 to 15 feet with no misting.

This treatment may injure bentgrass, St. Augustinegrass, centipedegrass, carpetgrass and newly seeded lawns. If it is necessary to control weeds in these grasses, use one-half the rate recommended above and make a second half-rate treatment 2 to 3 weeks later.

494

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

### NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

WEEDONE® is a registered trademark of Rhône-Poulenc.

©1990 Rhône-Poulenc Ag Company  
Made in U.S.A.

EPA Submission: 11/5/90