. 4 = = >

Ms. Lizbeth Huckaba
Rhone-Poulenc Ag Company
2 T.W. Alexander Drive
P.O. Box 12014
Research Triangle Park, NC 27709

Dear Ms. Huckaba:

Subject: Label Revisions - Revised Confidential Statement of Formula Sevin Brand 10% Bait Carbaryl Insecticide EPA Registration No. 264-312
Your Application Dated January 6, 1989

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

- Submit/cite all data required for registration/reregistration
 of your product under FIFRA section 3(c)(5) when the Agency
 requires all registrants of similar products to submit such
 data.
- 2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. The revised Confidential Statement of Formula (CSF) is not acceptable as submitted. According to our calculations, the lower limit comes to 10.04%. Submit a revised CSF to reflect this change.
 - b. Change the preharvest interval for alfalfa to 7 days.
 - c. Change the site name "Ornamental Plantings" to "Ornamental Plants."

Submit a revised CSF and final printed labeling with the revisions mentioned above within 60 days from the date of this letter.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

51616:I:CR-44:Kumar:E-3:KENCO:05/12/89:06/23/89:CL:VO:JH:AS

		CONCUR	RENCES			
SYMBOL					3 44	••••••
SURNAME	 				V Similar	* *
DATE						
	 			1	AUTO-DEELCI	AL FILE COR

EPA Form 1320-1 (12-70)

If these conditions are not on of a with, the registration will be subject to cancellation in accordance with PIFRA section o(e). Your release for shipment of the ordinat bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label to enclosed for your records.

Slacerely yours,

Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Baclocure

SEVIN® brand 10% Bait Carbaryl Insecticide

EPA Reg. No. 264-312

EPA Est. No. 10352-/GA-01

Made in U.S.A.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Page 2 for Additional PRECAUTIONARY STATEMENTS.

IN CASE OF AN EMERGENCY TELEPHONE (24 HOURS A DAY) IN THE USA
1-800-3344900071EDwith COMMENTS
to EPA Letter Dates

NET CONTENTS:

Under the Federal Insecticida, Fungicide, and Rodenticide Act as amended, for the pesticida registered under EPA Reg. No.

264-312

Rhône-Poulenc Ag Company
P. O. Box 12014, 2 T. W. Alexander Drive
Research Triangle Park, NC 27709

SEVIN is a registered trademark of Rhône-Poulenc for carbaryl insecticides.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

MAY BE HARMFUL IF SWALLOWED. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

STATEMENT OF PRACTICAL TREATMENT GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 or 2 glasses of water and induce vomiting be touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes, if needed, to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number 1-800-334-7577.

ENVIPONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment of disposal of wastes.

DIRECTIONS FOR USE


IT IS A VIOLATION OR FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when carbaryl is used in accordance with label directions. It SEVIN carbaryl insecticide is used in accordance with label directions these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops for which carbaryl tolerances are not established.



Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 10-20 pounds per acre of SEVIN® brand 10% Batt Carbaryl Insecticide will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs.

		Days Between Final Application
Crop	Dosage/Acre	and Harvest
Cucumbers, squash, melons, peanuts	10 pounds	None
Alfalfa, Peas	15 pounds	Days
Cotton	15 pounds	7 Days
Beans, carrots, corn, eggplants, okra, peppers, potatoes, tomatoes, and sweet com	10-20 pounds	None
Asparagus, strawberries	10-20 pounds	1 Day
Broccoli, brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	10-20 pounds	3 Days
Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard and turnips (tops)	10-20 pounds	14 Days

SITE	INSECT	OUNCES OF SEVIN® brand 10%BAIT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
Poultry Houses (chickens, turkeys, ducks, geese, gamebirds, and pigeons)	Darkling beetle (Lesser mealworm or litter beetle)	24	Apply bait uniformly to floor or littenance using appropriate equipment such as chest-carried fertilizer/seed spreader. It is not necessary to remove pirds during application, but uniform coverage may be difficult to achieve with birds present. Do not contaminate feeding or watering units.
			Optimum treatment timings, are: 1. Ten to 14 days after the birds are placed in the house and 2, two weeks after timing "#1" above. One application at each of the two timings listed above should be made for heavy infestations. If only one application can be made, timing #1 should be followed

SITE	INSECT	OUNCES OF SEVIN® brand 10%BAIT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
Poultry Houses (Cont.) (chickens, turkeys, ducks, geese, gamebirds, and pigeons)			For best control of high darkling beetle population: In addition to a bait application, walls, cracks, posts, crevices, and insulation areas should be treated with a sprayable or a flowable formulation of SEVIN® brand carbaryl insecticide. Spray applications should I made immediately following removal of the flock from thouse or immediately prior introducing the flock into thouse. All label directions and restrictions for the liquing application of SEVIN® brand carbaryl insecticide formulation used should be carefully followed.

SITE	INSECT	OUNCES OF SEVIN® brand 10%BAIT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
Turfgrass	Armyworms Crickets Cutworms Darkling ground beetle Grasshoppers Mole crickets Sowbugs	4 to 8	Treat entire area to obtain uniform coverage, giving particular attention to building perimeters, edges of sidewalks, and along porch an patio foundations. For best results, mow lawn, remove clippings, and thoroughly water area to be treated prior to application.
Ornamental plantings	Arthyworms Crickets Cutworms Darkling ground beetle Grasshoppers Mole crickets Sowbugs	4 to 8	Apply uniformly to soil around plants, shrubs, and trees. For best results, thoroughly water area to be treated prior to application.

-1

STORAGE AND DISPOSAL

STORAGE

Store unused product in original container only, in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

PESTICIDE DISPOSAL

Do not contaminate food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If bag is, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING ROM THE USE OR HANDLING OF THIS PRODUCT.