

-3/4

P/M

MAR 29 1985

**BEST AVAILABLE COPY**

Mr. J. S. Lowell  
Union Carbide Agricultural Products Company Inc.  
P.O. Box 17014  
Research Triangle Park, NC 27709

Dear Mr. Lowell:

**Subject: General Revision of Label in Response to the  
Carbaryl Registration Standard  
Sevin 50-W Carbaryl Insecticide  
EPA Registration No. 254-314  
Your Application Dated November 13, 1984**

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Submit five copies of finished printed labeling incorporating the comments listed below before you release the product for shipment:

1. In your Environmental Hazards paragraph delete the statement "Do not apply when weather conditions favor drift from area treated."
2. In your Bee Caution paragraph delete "Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions."

Sincerely yours,

Jay S. Ellenberger  
Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

**Enclosure**

CONCURRENCES

SYMBOL ▶	JOB: 85064; RD-85/01; Ellenberger; 3/13/85; D; 1/3/20/85; Kendrick; 898-1270; DAA						
SUBMITTER ▶							
DATE ▶							

2 of 22 # 264-314 PM 12

**BEST AVAILABLE COPY**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

SEVIN® brand 50-W  
Carbaryl Insecticide

**MAR 29 1985**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
264-314

For Agricultural Or Commercial Use Only

	Percent by wt.
ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-methylcarbamate) . . . . .	50% by wt.
INERT INGREDIENTS: . . . . .	50% by wt.
E.P.A. Reg. No. 264-314	E.P.A. Est. No. 477-MD-01

KEEP OUT OF REACH OF CHILDREN

**PRECAUTION**

See Page 2 for additional PRECAUTIONARY STATEMENTS.

IN CASE OF EMERGENCY TELEPHONE COLLECT (24 HOURS A DAY) IN THE U.S.A.  
(304) 744-3487.  
FOR GENERAL INFORMATION CALL 800-334-9745.

NET CONTENTS: \_\_\_\_\_

Made in U.S.A.

©UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.  
P. O. Box 12014, T. W. Alexander Drive  
Research Triangle Park, NC 27709

SEVIN is a registered trademark of Union Carbide Corporation for carbaryl insecticides.

APC Form No. \_\_\_\_\_

3/22 # 214-314 P.M.

## PRECAUTIONARY STATEMENTS

### CAUTION

**BEST AVAILABLE COPY**

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**MAY BE HARMFUL IF SWALLOWED.** Avoid breathing of dust and spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

#### ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

#### STATEMENT OF PRACTICAL TREATMENT

##### GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately, and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

**IF SWALLOWED:** Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF INHALED:** Move from contaminated atmosphere and call a physician.

##### NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

1 of 2 #. 264-314 Ph 12  
**BEST AVAILABLE COPY**

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Union Carbide Agricultural Products Company, Inc.

Consultation on therapy can be obtained at all hours by calling the Union Carbide emergency phone number (304) 744-3487.

#### ENVIRONMENTAL HAZARDS

~~This product is extremely toxic to aquatic and estuarine invertebrates. It is also highly toxic to birds and mammals. Do not apply to forest canopy and~~  
Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes.

#### BEE CAUTION

~~This product is highly toxic to bees exposed to direct treatment or residues on blooming crops and weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treated area.~~ Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

#### GENERAL INFORMATION

SEVIN® brand 50-W Carbaryl Insecticide is a dry powder for dispersion in water and application as an insecticide spray. READ THIS LABEL BEFORE USE. STRICTLY OBSERVE LABEL DIRECTIONS AND CAUTIONS, AND APPLICABLE FEDERAL AND STATE REGULATIONS.

#### GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENTS, STATEMENT OF PRACTICAL TREATMENT and

NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: Appropriate signal word (CAUTION), area treated with SEVIN brand 50-W Carbaryl Insecticide, date of application, appropriate clothing, and reentry interval (i.e., until sprays have dried).

PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when carbaryl is used in accordance with label directions. Pre-harvest and grazing use limitations are listed at the end of the directions for each crop use. Where no pre-harvest or grazing time limitations are stated, application can be made to the day of harvest and crops can be grazed without time delay if directions for use on the crop are followed. If SEVIN brand 50-W Carbaryl Insecticide is used in accordance with label directions, forage and feed crops, including almond hulls, bean vines, carrot tops, citrus pulp, cowpea hay, peanut hay, pea vines, rice straw, soybean hay, and sugar beet tops, may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

~~Do not plant rotational food and feed and crops not listed on this or other carbaryl labels in carbaryl treated soil.~~

~~Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops for which carbaryl tolerances are not established.~~

PLANT RESPONSE PRECAUTIONS:

To avoid possible injury on tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.

Do not use on Boston Ivy, Virginia creeper and maidenhair fern as injury will result. During early season, it may also injure Virginia and sand pines.

Observe label instructions on apple thinning and on combinations with certain herbicides on rice and soybeans.

*deleted  
18 mo. crop  
rotation  
restriction*

NOT REPRODUCIBLE COPY

### SPRAY PREPARATION

TO ASSURE A UNIFORM SUSPENSION, AGITATE, STIR OR RECIRCULATE ALL "SEVIN®" BRAND 50-W CARBARYL INSECTICIDE SPRAY MIXTURES PRIOR TO AND DURING USE. Remove oil, rust, scale, pesticide residues and water. Fill spray or mix tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of SEVIN® brand 50-W Carbaryl Insecticide and then the remaining volume of water. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURE FOR PROLONGED PERIODS.

### COMPATIBILITY

SEVIN brand 50-W Carbaryl Insecticide has been effectively used without plant injury with most common insecticides, miticides, fungicides, nutrients, adjuvants, and with petroleum oil as used on citrus. If compatibility with another product and the resulting crop response is unknown, it should be tested on a small scale. Curdling, precipitation, greasing, layer formation or increased viscosity are symptoms of incompatibility. Incompatibility will reduce insect control and may cause mechanical difficulties or plant injury. Observe all cautions and limitations on labeling of all products used in mixtures. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD "SEVIN®" BRAND 50-W CARBARYL INSECTICIDE TO 1/2 THE DESIRED AMOUNT OF WATER, MIX THOROUGHLY, AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE. DO NOT APPLY TANK MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS, EXCESSIVE RESIDUES OR PLANT INJURY.

Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime-sulfur and casein-lime spreaders. Do not use with summer oils on deciduous fruits.

### APPLICATION

On all crops use sufficient gallonage to obtain thorough and uniform coverage. Calibrate spray equipment to deliver the required volume. Use 50 mesh slotted strainers in spray system and 25 mesh slotted strainers behind nozzles. Avoid applications just before rainfall as poor insect control may result. To clean spray system after use, drain and flush with water. Refer to the Storage and Disposal directions for proper disposal of rinse water.

### INSECT CONTROL

Apply when insects or their damage appear. To maintain control repeat at 7 to 14 day intervals or as necessary unless a shorter interval is specified below. Where a dosage range is indicated, use lower rate on young plants and early instars and higher rate on mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

SEVIN® brand 50-W Carbaryl Insecticide does not control spider mites. If spider mites are a problem, use a registered miticide.

Regional differences have been noted in the susceptibility of certain strains of fall armyworm to SEVIN® brand 50-W Carbaryl Insecticide. If local experience indicates inadequate control, use an alternative pesticide.

FORAGE, FIELD AND VEGETABLE CROP INSECT CONTROL

Recommended dosages refer to pounds of SEVIN® brand 50-W Carbaryl Insecticide per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Use sufficient spray gallonage to obtain full coverage. This will usually vary from 3 to 40 gallons per acre depending on equipment, degree of insect infestation, and stage of crop growth. Keep spray mixture well agitated. Apply when insects or their damage appear and repeat at 7 to 10 day intervals or as necessary unless shorter interval is specified below. See specific directions below for grasshopper control.

ALFALFA AND CLOVERS

Use 1 to 2 pounds for blister beetles and Mexican bean beetle.

Use 2 pounds for alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, three-cornered alfalfa hopper, thrips and velvetbean caterpillar.

Use 2 to 3 pounds for alfalfa weevil larvae, armyworm, cloverhead weevil, corn earworm, cutworms, Egyptian alfalfa weevil larvae, essex skipper, European alfalfa beetle, fall armyworm, lygus bugs, stink bugs, webworms, and yellowstriped armyworm. Use higher rate in areas east of Rocky Mountains. On dense growth, use 25 to 40 gallons per acre with ground equipment to ensure adequate coverage. For alfalfa weevil larvae, if pretreatment damage is extensive, cut alfalfa and treat the stubble.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.



ASPARAGUS

Use 2 to 4 pounds for asparagus beetle on seedlings or spears.

Use 4 to 8 pounds for asparagus beetle and Apache cicada on ferns or brush growth in the post-harvest period.

Do not apply more often than once every 3 days.

Do not apply within 1 day of harvest.

BEANS (including blackeyed peas, ~~cowpeas~~, crowder or southern peas, dry beans, green beans, lima beans, navy beans and snap beans)

Use 1 to 2 pounds for blister beetles and Mexican bean beetle.

Use 2 pounds for alfalfa caterpillar, bean leaf beetle, cucumber beetles, flea beetles, green cloverworm, Japanese beetle, leafhoppers, three-cornered alfalfa hopper, thrips, velvetbean caterpillar, and western bean cutworm.

Use 2 to 3 pounds for armyworm, corn earworm, cutworms, European corn borer, fall armyworm, stink bugs, tarnished plant bug, and webworms.

Use 4 pounds for cowpea curculio.

In California, use 4 pounds for corn earworm, limabean pod borer, lygus bugs, and stink bugs.

**Do not apply to cowpeas within 3 days of harvest.**

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI**  
Use 1 to 2 pounds for flea beetles and harlequin bug.

Use 2 to 4 pounds for armyworm, fall armyworm, imported cabbageworm and corn earworm.

Do not apply within 3 days of harvest.

**CHINESE CABBAGE, COLLARDS, HANOVER SALAD, HORSERADISH, KALE, MUSTARD GREENS, RADISHES, RUTABAGAS AND TURNIPS.**

Use 1 to 2 pounds for flea beetles, harlequin bug and leafhoppers.

Use 2 to 3 pounds in a 5 to 7 day schedule for aster leafhopper.

Use 2 to 4 pounds for armyworm, fall armyworm, imported cabbageworm, corn earworm, tarnished plant bug and stink bugs.

Do not apply within 3 days of harvest of horseradish, radishes, rutabagas and turnips (roots).

Do not apply within 14 days of harvest of Chinese cabbage, collards, hanover salad, kale, mustard greens and turnips (tops).

**CARROTS, CELERY, PARSNIPS AND PARSLEY (DO NOT USE ON CELERY IN CALIFORNIA)**

Use 1 to 2 pounds for flea beetles and leafhoppers.

Use 2 to 3 pounds in a 5 to 7 day schedule for aster leafhopper, lygus bugs, and spittlebugs.

Use 2 to 4 pounds for armyworm, fall armyworm, corn earworm, tarnished plant bug and stink bugs.

No time limitation on carrots.

Do not apply within 3 days of harvest of parsnips.

Do not apply within 14 days of harvest of parsley and celery.



**CORN (FIELD, SWEET AND POP)**

Use 2 to 4 pounds for armyworm, chinch bugs, corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles, Southwestern corn borer, and leafhoppers. For larvae in whorl and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 1 to 6 day intervals starting when first silks appear and continuing until silks begin to dry. For chinch bugs use high gallonage ground application directed at the base of plants. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. OBSERVE BEE CAUTION.

Use 4 pounds for Western bean cutworm. Treat when infestation averages 15% and at 90% to 100% tassel emergence. Treatment after 100% silk emergence will reduce effectiveness.

Use 4 to 13 pounds in at least 15 gallons of water per acre for cutworm control. Spray in 12-inch band using 3/8 pound (6 ounces) per 1000 linear feet of row. For broadcast ground application, use up to 13 pounds in at least 20 gallons of water per acre.

**COTTON**

Use 1 to 2 pounds for early season control of cotton fleahopper, cotton leafworm, flea beetles, striped blister beetle, and thrips.

Use 2 to 4 pounds for mid and late season control of boll weevil, bollworms, cotton leafperforator, fall armyworm, leafrollers, leafhoppers, tarnished plant bug, and yellowstriped armyworm (cotton cutworm). Treat on a 5 to 7 day schedule for as long as necessary. May be applied after bolls open.

Use 2 to 4 pounds for light to moderate populations of lygus bugs in Western irrigated cotton.

Use 3 to 5 pounds for pink bollworm.

Use 4 pounds for stink bugs and saltmarsh caterpillar.

Aphid populations will be suppressed by repeated applications of this insecticide.

**Do not use [REDACTED] within 7 days of harvest.**

**CUCUMBER, MELONS, PUMPKIN AND SQUASH**

Use 1 to 2 pounds for pickleworm and melonworm; 2 pounds for cucumber beetles, flea beetles, leafhoppers and squash bug. Avoid excessive applications.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

**DANDELION, ENDIVE (ESCAROLE), LETTUCE AND SALSIFY**

Use 1 to 2 pounds for flea beetles, harlequin bug and leafhoppers.

Use 2 to 3 pounds in a 5 to 7 day schedule for aster leafhopper, lygus bugs, and spittlebugs.

**BEST AVAILABLE COPY**

10 of 22 #264-314 PM12

Use 2 to 4 pounds for armyworm, fall armyworm, imported cabbageworm, corn earworm, tarnished plant bug and stink bugs. For optimum control of corn earworm after lettuce heads begin to form, use 4 pounds in a 5 to 7 day schedule.

To avoid possible injury on lettuce, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

Do not apply within 3 days of harvest of head lettuce and salsify (roots).

Do not apply within 14 days of harvest of dandelion, endive (escarole), leaf lettuce and salsify (tops).

#### FORAGE GRASSES AND PASTURE

Use 2 to 3 pounds for armyworm, blackgrass bugs, chinch bugs, essex skipper, fall armyworm, range caterpillar, range crane fly, striped grass looper, and thrips. For thrips control in grasses grown for seed, use high spray pressure to improve penetration into boot.

Use 3 to 4 pounds for white grubs (green June beetle).

#### GARDEN BEET, SPINACH AND SWISS CHARD

Use 1 to 2 pounds for the flea beetles, harlequin bug and leafhoppers.

Use 2 to 3 pounds in a 5 to 7 day schedule for aster leafhopper.

Use 2 to 4 pounds for armyworm, corn earworm, fall armyworm, tarnished plant bug and stink bugs.

Do not apply within 3 days of harvest of garden beets (roots).

Do not apply within 14 days of harvest of garden beets (tops), spinach and Swiss chard.

#### OKRA

Use 2 to 4 pounds for corn earworm and stink bugs. Repeat at 5 to 7 day intervals or as necessary.

#### PEANUTS

Use 1 to 2 pounds for blister beetles and Mexican bean beetle.

Use 2 pounds for alfalfa caterpillar, bean leaf beetle, cucumber beetles, green cloverworm, Japanese beetle, leafhoppers, red-necked peanutworm, three-cornered alfalfa hopper, thrips and velvetbean caterpillar.

Use 2 to 3 pounds for armyworm, fall armyworm, corn earworm, stink bugs and webworms.

Use 4 pounds for white fringed beetle adults and cutworms.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

11 of 22

#264-314

PM 12

**BEST AVAILABLE COPY**

**PEAS**

Use 2 pounds for leafhoppers and Colorado potato beetle.

Use 2 to 3 pounds for armyworm.

Use 3 pounds for alfalfa caterpillar, cutworms, pea leaf weevil, pea weevil, and yellow-striped armyworm.

Use 5 pounds for alfalfa looper in Washington State only. Repeat as necessary.

**Do not apply within 3 days of harvest.**

**POTATO, TOMATO, EGGPLANT AND PEPPER**

Use 1 to 2 pounds for Colorado potato beetle, flea beetles and leafhoppers.

Use 2 to 4 pounds for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug and stink bugs.

Use 4 pounds for cutworm control.

**RICE**

CAUTION: May kill shrimp and crabs. Do not use in areas where these are important resources. ~~Do not use in rice fields in which crayfish and/or catfish farming are included in the cultural practice.~~

In the Mississippi Delta and Texas use 2 to 3 pounds for armyworm, fall armyworm, chinch bugs, and stink bugs.

DO NOT APPLY PROPANIL HERBICIDES WITHIN 15 DAYS BEFORE OR AFTER APPLICATION OF THIS PRODUCT OR PLANT INJURY WILL RESULT.

In California use 4 pounds for armyworm, leafhoppers and tadpole shrimp. For optimum tadpole shrimp control apply to the water when the pest first appears.

Do not apply within 14 days of harvest.

**SORGHUMS (milo, grain sorghum, sweet sorghum, and hybrids)**

Use 2 to 4 pounds for armyworm, chinch bugs, corn earworm, fall armyworm, stink bugs, and webworms. For chinch bugs use high gallonage ground application directed to base of plants.

Use 3 pounds for sorghum midge and Southwestern corn borer. Treat for sorghum midge when 25 to 30 percent of heads have emerged from boot and are in bloom. Repeat 3 to 5 days later if adults are still active. A third application may be necessary.

For optimum insect control, direct spray into forming heads.

Do not apply within 21 days of harvest of grain. No time limitations on forage.

Use 4 pounds for cutworm control.

**BEST AVAILABLE COPY**

**SOYBEANS**

DO NOT APPLY A COMBINATION OF THIS PRODUCT AND 2,4-DB HERBICIDES TO SOYBEANS AS CROP INJURY MAY RESULT.

Use 1 pound for light to moderate populations or 2 pounds for heavy infestations of bean leaf beetle, cucumber beetle, green cloverworm, and velvetbean caterpillar.

Use 1 to 1 1/2 pounds for light to moderate populations or 2 to 3 pounds for heavy infestations of corn earworm.

Use 1 to 2 pounds for blister beetles, grape colaspis, and Mexican bean beetle.

Use 2 pounds for alfalfa caterpillar, Japanese beetle, leafhoppers, three cornered alfalfa hopper, and thrips.

Use 2 to 3 pounds for armyworm, cutworm, fall armyworms, stink bugs, and webworms.

Use 4 to 5 pounds for painted lady (thistle caterpillar), saltmarsh caterpillar, and yellow striped armyworm.

Use higher rates for heavy infestations.

**SUGAR BEETS**

Use 2 to 3 pounds for armyworms, beet leaf beetle, fall armyworm, flea beetles, leafhoppers and webworms.

Use 3 pounds for cutworms.

Do not apply within 14 days of harvest.

**SWEET POTATOES - (DO NOT USE IN CALIFORNIA.)**

Use 13 Tbsps. per gallon of water as a preplant dip for sweet potato weevil. Just prior to planting, dip sweet potato cuttings in a suspension of SEVIN® brand 50-W Carbaryl Insecticide.

Use 2 to 4 pounds for corn earworm, cucumber beetles, flea beetles, sweet potato hornworm, sweet potato weevil, and tortoise beetles.

**TOBACCO**

**IN PLANT BEDS**

For tobacco flea beetle, use 4 level tablespoonfuls per gallon or 2 pounds per 50 gallons, and apply 6 gallons per 100 square yards.

For green June beetle grubs, use 1 pound per 100 gallons of water when insects or their damage appear. Apply only to areas that larvae have uprooted by sprinkling as a drench with 50 to 100 gallons per 100 square yards. May be applied before or after seeding. Avoid excessive application as plant injury may result.

**BEST AVAILABLE COPY****IN FIELDS:**

For budworms, fall armyworm, flea beetles, Japanese beetle, June beetles, hornworms and suckfly, use 2 to 4 pounds in 50 to 100 gallons of water per acre. Full coverage of plants is essential. Use lower rate on young plants up to knee height, higher rate on mature plants. Begin treatments when worms are small. Avoid excessive application in bud of plant as injury may result.

Treated fields may be entered immediately after foliage has dried. If late applications are necessary, allow 3 days before pruning or cutting.

**GRASSHOPPERS**

Apply 1 to 3 pounds as often as necessary. TO CONTROL GRASSHOPPERS ON THE ABOVE FORAGE FIELD AND VEGETABLE CROPS, FOLLOW PREHARVEST LIMITATIONS GIVEN AFTER EACH CROP. A 1 to 2 pounds rate is suggested for nymphs on small plants or sparse vegetation in pastureland, rangeland, ditchbanks and borders. A 2 to 3 pounds dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage.

**TREE FRUIT AND NUT INSECT CONTROL**

Recommended dosages refer to pounds of **SEVIN® brand 50-W Carbaryl Insecticide** per 100 gallons of dilute spray in conventional hydraulic-type sprayers. Apply dilute sprays for full coverage. Spray gallonage will vary according to tree size, density, planting distance and stage of growth. With concentrate spray, where lower volumes of spray are applied per acre, the concentration of **SEVIN® brand 50-W Carbaryl Insecticide** should be increased in the spray mixture so that the amount of **SEVIN® brand 50-W Carbaryl Insecticide** applied per acre is equivalent to a full-coverage dilute spray. Keep spray mixtures well agitated. Additional instructions appear with the crop categories below.

**ALMOND**

Use 2 pounds for peach twig borer, San Jose scale and fruit tree leafroller. Apply in "popcorn" or petal fall stages and again when the May brood of the peach twig borer begins to hatch or thereafter as needed.

Use 2 pounds for navel orangeworm. Timely and mid season applications to correspond to moth flight peaks. Make late season application at initiation of or up to 10% hull split.

Do not apply more than 10 pounds per acre.

Do not apply within 28 days of harvest.

**APPLES AND PEARS**

Application of **SEVIN® brand 50-W Carbaryl Insecticide** within 30 days after full bloom may also provide apple thinning; to avoid this, delay use until at least 30 days after bloom. Foliar injury may result from combination with summer oils west of the Rocky Mountains, or when used before second cover on York and McIntosh apples east of the Rocky Mountains.

West of the Rocky Mountains: Use 1 1/2 to 2 pounds for apple sucker, apple aphid, woolly apple aphid, bagworms, California pearslug (pear sawfly), codling moth, apple rust mite, pearleaf blister mite, pear rust mite, eyespotted bud moth, green fruitworm, lesser appleworm, lygus bugs, orange tortrix, psylla, tentiform leafminers, lecanium scales, oystershell scale, and San Jose scale. For optimum scale control apply when crawlers are present. For psylla control apply when eggs hatch or when young nymphs are present. When heavy aphid infestations are present, repeated applications may be necessary for optimum control.

East of the Rocky Mountains: Use 1 pound for apple mealybug, apple aphid, codling moth and white apple leafhopper. When heavy aphid infestations are present, repeated applications may be necessary for optimum control.

Use 2 pounds for apple maggot, bagworms, eastern tent caterpillar, European apple sawfly, eyespotted bud moth, fruittree leafroller, green fruitworm, Japanese beetle, lesser appleworm, pear psylla, periodical cicada, plum curculio, redbanded leafroller, rosy apple aphid, woolly apple aphid, apple rust mite, pearleaf blister mite, pear rust mite, tarnished plant bug, tentiform leafminers, yellow headed fireworm, forbes scale, lecanium scales, oystershell scale, and San Jose scale. For optimum scale control, apply when crawlers are present. To control rosy apple aphid, apply before leaves are curled.

Do not apply within 1 day of harvest of apples and pears.

#### APPLE THINNING

Many factors influence the degree of apple thinning obtained with SEVIN brand 50-W Carbaryl Insecticide. These include tree age, variety, nutrition, previous crop, pruning, degree of set, bloom, weather and use of other chemical thinners. When any of these favor reduced fruit set, caution should be observed in applying SEVIN brand 50-W Carbaryl Insecticide so that excessive thinning and possible yield reduction will not occur. Recommended dosages refer to pounds of SEVIN brand 50-W Carbaryl Insecticide per 100 gallons dilute spray.

Consult with your County Extension Service or other experts for advice on the proper use of this product.

Concentration: Use 1/2 to 1 pound on easily thinned varieties. Apply for full coverage. On hard-to-thin varieties, use 1 to 2 pounds.

Timing: Apply in one spray timed between 10 and 25 days after full bloom.

Varietal Response: Easily-thinned varieties include Cortland, Grimes, Jonathan, McIntosh, Orleans, Rome Beauty, Puritan, Red Delicious, Winesap, and Yellow Newton. Difficult-to-thin varieties include Baldwin, Ben Davis, Dutchess, Early McIntosh, Golden Delicious, Lady Apple, Northern Spy, Rhode Island Greening, Steele Red, Turley, Wealthy, Yellow Transparent, and York Imperial.

CITRUS FRUITS (such as grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats and hybrids).

Use 2 pounds for avocado leafroller, California orangedog, citrus cutworm, citrus root weevil, fruittree leafroller, orange tortrix, Western tussock moth and West Indian sugarcane borer (adults).

Use 1 1/2 to 2 pounds for black scale, brown soft scale, California red scale, citricola scale, citrus scale, and yellow scale.

Apply lower dosage early in the season or against light infestation and the higher dosage late in the season or against heavy infestation and situations involving infested fruit. May be used with petroleum oil as used in common practice on citrus.

Do not apply more than 40 pounds of SEVIN® brand 50-W Carbaryl Insecticide per acre per application.

Do not apply less than 10 gallons of dilute spray mixture per mature tree.

Do not apply within 5 days of harvest of these crops.

CHESTNUTS - DO NOT USE IN CALIFORNIA.

Use 6 pounds for chestnut weevil.

Make 4 applications at weekly intervals beginning in late July for adult chestnut weevil control. Last application should be made prior to shuck split.

#### FILBERT

Use 2 pounds for filbert aphid, filbert leafroller and fibertworm. Apply when leafroller eggs are hatching. Repeat on first appearance of adult fibert moths and again 3 to 4 weeks later. With dilute sprays, 400 gallons per acre is suggested.

#### OLIVES

Use 1 1/2 to 2 pounds with 1 1/2 gallons of summer oil, in sprays for olive scale control. For optimum scale control, apply when crawlers are present. A maximum of 2 applications is permitted.

Do not apply more than 30 pounds of this product per acre per application.

#### PEACHES APRICOTS AND NECTARINES

Use 2 pounds for codling moth, cucumber beetles, European earwig, Japanese beetle, apple pandemis, June beetles, lesser peachtree borer, orange tortrix, oriental fruit moth, peach twig borer, periodical cicada, plum curculio, fruittree leafroller, redbanded leafroller, variegated leafroller, tarnished plant bug, tussock moths, lecanium scale, San Jose scale and olive scale. For optimum scale control, apply when crawlers are present. For lesser peachtree borer control, spray limbs and trunk thoroughly, weekly during moth flight.

Do not apply more than 12 pounds of this product per acre per application on apricots.

**BEST AVAILABLE COPY**

Do not apply within 1 day of harvest of peaches.

Do not apply within 3 days harvest of apricots and nectarines.

#### PEACHES

Use 2 to 5 pounds for black margined aphid, fall webworm, hickory shuckworm, lesser webworm, pecan leaf phylloxera, pecan nut casebearer, pecan spittlebug, pecan weevil, twig girdler, walnut caterpillar.

Do not apply more than 14 2/5 pounds of SEVING brand 50-W Carbaryl Insecticide per acre per application.

#### PLUMS, PRUNES, AND CHERRIES

Use 2 pounds for black cherry aphid, nearly plum aphid, cherry maggots, cherry fruitworm, eyespotted bud moth, fruitree leafroller, redbanded leafroller, Japanese beetle, lesser peachtree borer, peach twig borer, plum curculio, prune leafhopper, European earwig, green fruitworm, rose chafer, variegated leafroller, brown soft scale, Forbes scale, lecanium scales, oystershell scale, and San Jose scale. For optimum scale control, apply when crawlers are present. For lesser peachtree borer control spray limbs and trunk thoroughly, weekly during moth flight.

Use 1 1/2 pounds for eastern tent caterpillar, codling moth, orange tortrix, and tussock moths.

Do not apply more than 12 pounds of this product per acre per application.

Do not apply within 1 day of harvest of these crops.

#### PISTACHIOS

Use 1 to 4 pounds per 100 gallons of water for navel orangeworm control. Apply 150 to 300 gallons of mixed spray per acre for full coverage at onset of hull split. Do not use within 14 days of harvest.

#### WALNUT

Use 1 pound for calico scale, codling moth, European fruit lecanium, filbert-worm, fruitree leafroller, and frosted scale. Apply 1000 gallons of diluted spray per acre for mature trees. For codling moth, apply first spray when average cross-sectional diameters of developing nuts are 1/2 to 3/4 inch. Repeat during middle or late June as needed.

Use 4 pounds for European earwig. Spray tree trunks to point of run-off.

#### SMALL FRUIT INSECT CONTROL

Recommended dosage refers to pounds of SEVING brand 50-W Carbaryl Insecticide per acre. Keep spray mixtures well agitated. Apply when insects or the damage appear. Repeat at 7 to 10 day intervals or as necessary.

Typical spray gallonage per acre ranges from 100-300 gallons for dilute sprays, and 30-100 gallons for concentrate sprays. Do not exceed maximum label rate per acre per application.



**BEST AVAILABLE COPY**

**BLACKBERRIES, RASPBERRIES AND DEWBERRIES (including Boysenberries and Loganberries)**

Use 2 to 4 pounds for Japanese beetle, leafrollers, leafhoppers, European raspberry aphid, rose chafer, snowy tree cricket.

In California, use 4 pounds for omnivorous leafroller and raspberry sawfly.

Do not apply within 7 days of harvest of these crops.

**BLUEBERRIES**

Use 3 pounds for blueberry maggot, cherry and cranberry fruitworms, European fruit lecanium and Japanese beetle. Apply 3 weeks before harvest and repeat as necessary.

**CRANBERRIES**

Use 3 to 6 pounds for cutworms, cranberry fireworms, cranberry fruitworm, Japanese beetle, leafhoppers, and rose chafer. Apply in late bloom and as needed at 7 to 10 day intervals.

Do not apply within 1 day of harvest.

**CAUTION:** May kill shrimp and crabs. Do not use in areas where these are important resources.

**GRAPES**

Use 2 to 4 pounds for European fruit lecanium, grape leafroller, grape leafhopper, western grapeleaf skeletonizer, and Western yellowstriped armyworm.

Apply just before first brood leafroller larvae emerge from rolls and as needed for leafhoppers.

Use 4 pounds for grape berry moth, Japanese beetle, June beetles, orange tortrix, omnivorous leafroller, redbanded leafroller, cutworms, eight spotted forester, and saltmarsh caterpillar.

**STRAWBERRIES**

Use 2 to 4 pounds for flea beetles, meadow spittlebug, omnivorous leaf tier (strawberry fruitworm), strawberry leafroller and strawberry weevil. Carbaryl may injure Early Dawn and Sunrise varieties on Delmarva Peninsula.

Do not apply within 1 day of harvest.

## TREES AND ORNAMENTALS INSECT CONTROL

For dilute spray ground applications to trees (including shade trees, sugar maple, shelter belts, forests, plantations, parks, and recreational areas), ornamentals (including roses, flowers and other herbaceous plants), woody plants and shrubs, apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage. To prepare small volumes of spray, use 2 tablespoonsful of SEVIN® brand 50-W Carbaryl Insecticide per gallon of water where rates of 2 pounds are indicated.

Use 2 pounds for control of:

Ants	Fall armyworm	Nantucket pine	Saddled prominent
Apple aphid	Flea beetles	tip moth	Sawflies (exposed)
Armyworm	Fuller rose beetle	Oak leafminers	Scale insects
Azalea leafminer	Gall midges	Oak leaf	Sowbugs
Bagworms	Gall wasps	skeletonizer	Spiny elm
Birch leafminer	Green striped	Oakworm complex	caterpillar
Blister beetle	mapleworm	Oleander caterpillar	Springtails
Boxelder bug	Grasshoppers	Olive ash borer	Spruce budworm
Boxwood leafminer	Gypsy moth	Orange-striped	Spruce needle-
Brown tail moth	Hackberry nipple-	oakworm	miner
Cankerworms	gall maker	Orange tortrix	Subtropical pine
Catalpa sphinx	Holly bud moth	Periodical cicada	tip moth
Chiggers	Holly leafminer	Pine sawfly	Tent caterpillars
Cooley spruce	Jackpine budworm	Pine spittlebug	Thorn bug
gall aphid	Japanese beetle	Pitch pine tip moth	Thrips (exposed)
Cutworms	Jeffrey pine	Plant bugs	Ticks
Cypress tip moth	needleminer	Poinsettia	Walnut caterpillar
Douglas-fir	June beetles	hornworm	Webworms
tussock moth	Lace bugs	Psyllids	Western hemlock
Eastern spruce	Leafhoppers	Puss caterpillar	loopers
gall aphid	Leafrollers	Redhumped	Western spruce
Elm leaf aphid	Locust borer	oakworm	budworm
Elm leaf beetle	Maple leafcutter	Rose aphid	Willow leaf
Elm spanworm	Mealybugs	Rose chafer	beetles
Eriophyid mites	Mimosa webworm	Roseslug	Yellow poplar
European pine			weevil
shoot moth			

Apply dilute sprays to obtain thorough coverage of upper and lower leaf surfaces. To control scale insects, treat trunks, stems, and twigs in addition to plant foliage. For optimum worm control, treat when in early instars. Addition of a sticker may improve residual control.

Applications for control of Maple leafcutter on sugar maple should be made when larvae are in 2nd instar, after mining, and as cases are being formed.

Use 40 pounds for control of Ips engraver beetles, mountain pine beetle, roundheaded pine beetle, and western pine beetle. Effective as a preventive treatment only. Repeat annually as required to prevent beetle attacks. Apply 1 gallon of spray per 50 square feet of bark in late May to early July, or prior to beetle attack. Treat tree trunks from ground level up, until trunk diameter is less than 5 inches.

**BEST AVAILABLE COPY**

**LAWN AND RECREATIONAL AREA INSECT CONTROL**

Use 2 pounds of SEVIN® brand 50-W Carbaryl Insecticide in 150 to 200 gallons of water for each 5000 square feet (4 tablespoonfuls in 3 to 4 gallons of water for each 100 square feet) of established lawn area for the control of:

- |                   |                 |                   |                            |
|-------------------|-----------------|-------------------|----------------------------|
| armyworm          | cutworms        | fleas             | mosquitoes                 |
| ants              | earwigs         | grasshoppers      | sod webworms               |
| bluegrass billbug | essex skipper   | green June beetle | (lawn moths)               |
| centipedes        | European chafer | leafhoppers       | sowbugs                    |
| chiggers          | fall armyworm   | lucerne moth      | springtails                |
| chinch bugs       | fiery skipper   | millipedes        | ticks                      |
|                   |                 |                   | yellow-striped<br>armyworm |

Use garden hose sprayers or pressure type equipment.

Insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

Use 3 pounds of this product in 150 to 200 gallons of water per 5000 square feet for control of chinch bugs and white grubs.

**IMPORTED FIRE ANT CONTROL**

To prepare small volumes of spray, use 1/2 ounce of SEVIN® brand 50-W Carbaryl Insecticide per gallon water.

SITE	Pounds of SEVIN® BRAND 50-W	SPECIFIC INSTRUCTIONS
Lawns, cemeteries and Recreational Areas (including turf, golf courses, and parks) Pastures Rangeland Forested Lands and Wastelands	3	Apply a total of 2 gallons of the diluted solution over the surface of each mound or at least 1 quart per 6 inches of mound diameter using a bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). Do not disturb mounds prior to treatment. Pour solution from a height of about 3 ft. to give sufficient force to break mound apex and flow into ant tunnels. For best results apply in cool weather, 65-80°F, or in early morning or late evening hours. Repeat application if mound activity resumes after 10 days. Treat new mounds as they appear. Pressurized spray may disturb the ants and causes migration, reducing product effectiveness.

**BEST AVAILABLE COPY**

**ADULT MOSQUITOES**

On pasture, rangelands and non-agricultural lands such as yards, parks, recreation areas, logging camps, military posts and adjacent forested lands or wastelands, use 1/2 to 1 pound per acre in mist blowers, 2 pounds per acre in suitable low pressure ground equipment. Apply in sufficient spray gallonage to effectively cover areas to be treated. For optimum control, apply in early morning or evening when air is calm and adult mosquitoes are active.

Use 2 pounds per 100 gallons in hydraulic sprayers or 2 tablespoonfuls per gallon in hand sprayers for treating backyards or other limited areas. Thoroughly cover lower shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate. Repeat at 7 to 10 day intervals or as necessary. Do not use on Boston ivy, Virginia creeper and maidenhair fern as injury will result.

**OBSERVE BEE CAUTION.**

**CAUTION:** May kill shrimp and crabs. Do not use in areas where these are important resources.

Use 50 pounds of SEVIN® brand 50-W Carbaryl Insecticide in 100 gallons of water for residual control in subtropical regions. Apply 4 gallons of prepared spray per 2000 square feet of surface area. Repeat in 3 to 6 months or when necessary.

**PEST CONTROL IN AND AROUND BUILDINGS**

**LIMITED TO PEST CONTROL OPERATORS ONLY**

For use in and around buildings such as homes, apartments, warehouses, barns, and municipal and recreation areas.

Use 40 pounds per 100 gallons of water (40 tablespoons per gallon of water) for ants and cockroaches.

Use 20 pounds per 100 gallons of water (20 tablespoons per gallon of water) for brown dog tick, earwigs, fleas, millipeds, and silverfish.

Apply as a coarse wet spray or with a paint brush, and thoroughly treat cracks and crevices, base boards, corners, walls, door sills and window frames, quarters of household pets, ant trails, outside perimeter of dwellings and other areas where insects tend to congregate.

Do not use as a space spray. Repeat when necessary, but not more than twice per week.

~~Do not use in areas where food is exposed or where food is prepared or processed.~~ Do not use in serving areas while food is exposed.

Avoid application to surfaces where visible spray residues are objectionable.

**BEST AVAILABLE COPY**

## POULTRY INSECT CONTROL

### CHICKENS, DUCKS, GAME BIRDS, GEESE, PIGEONS AND TURKEYS

Direct Mist Spray on Birds: Control northern fowl mite, chicken mite, lice and fleas by:

1. Misting with Electric Fog Machine: Mix 5/8 pound (10 ounces) of SEVIN® brand 50-W Carbaryl Insecticide in 1 gallon of spray. Use 1 1/2 gallons per 1000 hens in cages, on litter or slatted floor. Repeat in 4 weeks if necessary.

or

2. Spraying with knapsack or cylinder type compressed air sprayers. Mix 3/8 pound (6 ounces) of this product in 5 gallons of spray. Use 1 gallon per 100 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

Direct mist spraying for chicken mite and fleas, as a supplement to spraying roosts and buildings for control of these pests.

For bedbugs, chicken mite, fleas, fowl tick, lesser mealworms, lice and northern fowl mite; spray roosts and buildings with conventional power spray or knapsack equipment. For chicken mite, fleas and bedbugs, use 2 pounds per 25 gallons of water. For fowl ticks, use 8 pounds per 25 gallons of water. Spray 1 to 2 gallons per 1000 square feet of wall, bedding, litter or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Ventilate while spraying. Avoid spraying of nests, eggs and feeding and watering troughs.

Do not apply to poultry and game birds or to their premises within 7 days of slaughter.

## STORAGE AND DISPOSAL

### STORAGE

Store unused SEVIN® brand 50-W Carbaryl Insecticide in original container only, in cool, dry areas out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

### PESTICIDE DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If bag is burned, stay out of smoke.

**BEST AVAILABLE COPY**

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

100-1  
PM 12  
Reg, # 264-316

MAR 29 1985

**BEST AVAILABLE COPY**

Mr. J. S. Lovell  
Union Carbide Agricultural Products Company, Inc.  
P.O. Box 12014  
Research Triangle Park, NC 27709

Dear Mr. Lovell:

Subject: General Revision of Label in Response to the  
Carbaryl Registration Standard  
Sevin 80S Carbaryl Insecticide  
EPA Registration No. 264-316  
Your Application Dated November 13, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Submit five copies of finished printed labeling incorporating the comments listed below before you release the product for shipment:

1. In your Environmental Hazards paragraph delete the statement "Do not apply when weather conditions favor drift from area treated."
2. In your Bee Caution paragraph delete "Do not use when value of bees as pollinators is more important than insect control.  
Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions."

Sincerely yours,

Jay S. Ellenberger  
Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosure		CONCURRENCES					
SYMBOL							
SURNAME	RD-IRB/JOB; 85057; RD45; Ellenberger; Kendrick; 898-1270; 2/27/85; Del. 3/7/85; dac						
DATE							

2 2 2 2 4-314 20 15

**BEST AVAILABLE COPY**

**ACCEPTED**  
with **COMMENTS**  
in EPA Letter Dated:

**SEVIN® brand 50-W**  
**Carbaryl Insecticide**

**MAR 29 1985**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
264-314

For Agricultural Or Commercial Use Only

	Percent by wt.
ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-methylcarbamate) . . . . .	50% by wt.
INERT INGREDIENTS: . . . . .	50% by wt.
E.P.A. Reg. No. 264-314	E.P.A. Est. No. 477-MD-01

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See Page 2 for additional PRECAUTIONARY STATEMENTS.

IN CASE OF EMERGENCY TELEPHONE COLLECT (24 HOURS A DAY) IN THE U.S.A.  
(304) 744-3487.  
FOR GENERAL INFORMATION CALL 800-334-9745.

NET CONTENTS. \_\_\_\_\_ Made in U.S.A.

©UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.  
P. O. Box 12014, T. W. Alexander Drive  
Research Triangle Park, NC 27709

SEVIN is a registered trademark of Union Carbide Corporation for carbaryl  
insecticides.

APC Form No. \_\_\_\_\_



3 of 22 # 264-314 PM12

## PRECAUTIONARY STATEMENTS

CAUTION

**BEST AVAILABLE COPY**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**MAY BE HARMFUL IF SWALLOWED.** Avoid breathing of dust and spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

STATEMENT OF PRACTICAL TREATMENT

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

**IF SWALLOWED:** Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF INHALED:** Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (79-567)  
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

128-125

DATE OF ISSUANCE

3/21/85

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

Spraypak Wax & Cornet  
Spray

NAME AND ADDRESS OF REGISTRANT (include ZIP code)

White Products Company  
11 & Gardner Road  
Bedford, N.J. 07009

WEST LAMBERT, MISSOURI

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(7)(A) when the Agency requires all registrants of similar products to submit such data.

2. Add the phrase "EPA Registration No. 128-125" to your label before you release the product for shipment.

3. Submit five (5) copies of your final printed label no before you release the product for shipment. Refer to the label specifications for further description of final printed labeling.

\* If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 3(c)(7)(A). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

For further information  
contact the Office of  
Inspection and Compliance  
Registration Division (79-567)

REG. DIVISION

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

*[Signature]*

DATE

3/21/85