

24/2
264-314 Y2

ACCEPTED
JUL 27 1987
Under the Insecticide Fungicide and Rodenticide Act
EPA Reg. No. 227/107

GRASSHOPPERS
Apply 1 to 3 pounds as often as necessary to control grasshoppers on the forage field and vegetable crops, follow PRE-HARVEST APPLICATIONS GIVEN AFTER EACH CROP A 1 to 2 pounds rate is suggested for nymphs or small plants in sparse vegetation in residential, rangeland, ditchbanks and borders. A 2 to 3 pounds dosage should be used when grasshoppers are mature or mature & applied to crops requiring greater coverage.

SMALL FRUIT INSECT CONTROL
Recommended dosage rates in pounds of SEVIN[®] brand 50-W Carbaryl insecticide per acre. Keep spray mixture well agitated. Apply when insects or the damage appear. Repeat at 7 to 10 day intervals or as necessary.

BLUEBERRIES, RASPBERRIES AND STRAWBERRIES (including Raspberry and Loganberries) Use 2 to 4 pounds for Japanese beetles, weevilers, leafhoppers, European raspberry aphid, rose chaffer, and/or rose cricket. In California use 4 pounds for omnivorous weeviler and raspberry weevil. Do not apply within 7 days of harvest of these crops.

BLUEBERRIES: Use 3 pounds for blueberry maggot, cherry and cranberry fruitworms, European fruit lecanium and Japanese beetle. Apply 3 weeks before harvest and repeat as necessary.

CRANBERRIES: Use 3 to 6 pounds for cutworms, cranberry fruitworms, cranberry fruitworm, Japanese beetle, leafhoppers, and rose chaffer. Apply in late bloom and as needed at 7 to 10 day intervals. Do not apply within 1 day of harvest.

CAUTION: May kill shrimp and crabs. Do not use in areas where these are important resources.

GRAPES: Use 2 to 4 pounds for European fruit lecanium, grape leafhopper, grape leafhopper, eastern grape leafhopper and western grape leafhopper. Apply to leafhoppers and broad leafhoppers larvae emerging from roots and as needed for leafhoppers. Use 4 pounds for grape berry moth, Japanese beetle, June beetle, orange foot omnivorous weeviler, ribworted leafhopper, cutworms, eight spotted leafhopper and saltmarsh caterpillar.

STRAWBERRIES: Use 2 to 4 pounds for flea beetle, meadow spittlebug, omnivorous weeviler, strawberry fruitworm, strawberry leafhopper and strawberry weevil. Carbaryl may harm Earth Dawn and Sunrise varieties on Delmarva Peninsula. Do not apply within 1 day of harvest.

TREES AND ORNAMENTAL INSECT CONTROL
For dilute-spray ground applications to trees (including shade trees, sugar maples, ashlar beetle, forest tent caterpillar, parula, and neotropical forest caterpillar) (including roses, flowers and other herbaceous plants), spray plants and shrubs, apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in full-canopy spray. June to provide thorough coverage. Use 2 pounds for control of:

Ants	European pine shoot moth	Mealybugs	Phyllorhiza
Apple aphid	Fall armyworm	Mineral woolworm	Saltmarsh caterpillar
Armyworm	Fall armyworm	Nantuxial pine tip moth	Specific leafminer
Basswood leafminer	Flea beetles	Oil spittlebugs	Scale insects
Basswood	Flea beetles	Oil spittlebugs	Scale insects
Bark weeviler	Gall midges	Osage weeviler	Scythid
Bark weevil	Gall wasps	Osage weeviler	Scythid
Borers	Green spruce aphid	Oswald complex	Springtails
Borers	Green spruce aphid	Oswald complex	Springtails
Bronze leaf miner	Mapleworm	Oswald complex	Springtails
Brown leaf miner	Mapleworm	Oswald complex	Springtails
Caterpillars	Mapleworm	Oswald complex	Springtails
Colorado spruce aphid	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails
Corn	Mapleworm	Oswald complex	Springtails

RECREATIONAL AREA INSECT CONTROL
Use 1/2 to 1/4 pound of SEVIN[®] brand 50-W Carbaryl insecticide per 1000 square feet (1/2 to 1/4 pound per acre).

RECREATIONAL AREA INSECT CONTROL
Use 1/2 to 1/4 pound of SEVIN[®] brand 50-W Carbaryl insecticide per 1000 square feet (1/2 to 1/4 pound per acre).

Ants	Corn	Japanese beetle	Scale insects
Armyworm	Eastern spruce aphid	Leafhoppers	Springtails
Bark weeviler	European leafminer	Lycophotus	Springtails
Bark weevil	Fall armyworm	Lycophotus	Springtails
Borers	Fern weeviler	Mistlebeetle	Springtails
Bronze leaf miner	Flies	Ring-necked pupa	Springtails
Brown leaf miner	Grasshoppers	Tree toad	Springtails

SUPPORTED FIRE ART CONTROL

SEVIN[®] 50W
100 gal. water

INSTRUCTIONS
Apply at rate of 2 gallons of the 50-W Carbaryl insecticide per acre. Repeat application if necessary. Do not apply more than 3 gallons per acre per season.

PEST CONTROL IN AND AROUND BUILDINGS
LIMITED TO PEST CONTROL OPERATORS ONLY
Use in and around buildings such as warehouses, barns, and nurseries for control of insects and other pests. Do not use for pest control in and around buildings.



CARBARYL INSECTICIDE
FOR AGRICULTURAL OR COMMERCIAL USE ONLY
ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate)
INERT INGREDIENTS:
EPA Reg No 264-314
50% by wt
50% by wt

E.P.A. EST. NO. 477-MD-01-34704-MS-1
Refer to first letter of lot number for packaging location

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
CAUTION
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
MAY BE HARMFUL IF SWALLOWED. Avoid breathing dust and spray. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long sleeved work clothing. Wash face and hands daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT
ANTIDOTE IS GLENEPHEPRINE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

STATEMENT OF PRACTICAL TREATMENT
See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN
Carbaryl is a carbamate insecticide. It is highly effective against a wide range of insects. It is not effective against sucking insects such as aphids, cicadas, and psyllids. It is not effective against fleas, ticks, and mites. It is not effective against termites.

STORAGE AND DISPOSAL
STORAGE
Store in a cool, dry place. Do not store in areas where children and animals can access. Do not store in areas where food or feed is stored.

PESTICIDE DISPOSAL
Do not dump or pour into water or soil. Do not burn. Do not bury. Do not incinerate.

(24 HOURS A DAY) IN U.S.A. 1-800-UCC-HELP

FOR GENERAL PRODUCT INFORMATION CALL 1-800-334-6768

Direct Mail Spray on Birds: Control northern leaf-neck, chicken mite, lice and fleas by

- Misting with Direct Mail. Max. 5.0 pound (10 ounces) of SEVIN® brand 50-w Carbaryl Insecticide in 1 gallon of spray. Use 1 1/2 gallons per 1000 hens in cages on tier or stalled floor. Repeat in 4 weeks if necessary.
- Spraying with knapsack or cylinder type compressed air sprayer. Max. 5.0 pound (10 ounces) of this product in 5 gallons of spray. Use 1 gallon per 100 hens in cages on tier or on stalled floor. Repeat in 4 weeks if necessary.

Direct Mail is only for chicken mite and lice. Use as a supplement to spraying roosts and buildings for control of these pests.

For bedbugs, chicken mite, flea, tick, head louse, earwig, and northern leaf-neck, spray roosts and buildings with conventional power spray or knapsack equipment. For chicken mite, flea and bedbugs, use 2 ounces per 25 gallons of water. For ticks, use 8 pounds per 25 gallons of water. Spray 1 to 2 gallons per 1000 square feet of wall, bedding, nest or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Vibrate while spraying. Avoid spraying of nests, eggs and feeding and nesting troughs.

Do not apply to poultry and game birds or to their premises within 7 days of slaughter.

BEST AVAILABLE COPY

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label, (b) that the product is reasonably fit for the uses and purposes set forth in the directions for use when it is used in accordance with such directions, and (c) that the directions, warnings and other statements on the label are based upon responsible expert evaluation of residue, its rate of effectiveness, or toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER, NEITHER MAKES NOR INTENDS NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAIRMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

© UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
 P.O. Box 12014, T.W. Alexander Drive, Research Triangle Park, NC 27709

SEVIN is a registered trademark of Union Carbide Corporation for carbaryl insecticides.

NET WT. 5 LBS. APC-1702-002-17
 Made in U.S.A.

10/2

CHINESE CABBAGE, COLLARDS, HARROW SALAD, NORSEADISH, KALE, MUSTARD GREENS, RADICCHIO, RUTABAGA AND TURNIPS

Use 1 to 2 pounds for flea beetles, flea hopper, and leafhopper.
Use 2 to 3 pounds for flea beetles, flea hopper and leafhopper.
Use 2 to 4 pounds for armyworm, fall armyworm, imported cabbageworm, corn earworm, tarnished plant bug and tortoise beetle.

Do not apply within 3 days of harvest of headcabbage, collards, rutabagas and turnips/roots.
Do not apply within 14 days of harvest of Chinese cabbage, collards, harrow salad, kale, mustard greens and turnip/roots.

CARROTS, CELERY, PARSNIPS AND PARSLEY (DO NOT USE ON CELERY IN CALIFORNIA)

Use 1 to 2 pounds for flea beetles and leafhoppers.
Use 2 to 3 pounds for flea beetles and leafhoppers (not for use on leafhoppers).
Use 2 to 4 pounds for armyworm, fall armyworm, corn earworm, tarnished plant bug and tortoise beetle.

No insect application on carrots.
Do not apply within 3 days of harvest of carrots.
Do not apply within 14 days of harvest of parsley and celery.

CORN (FIELD, SWEET AND POP)

Use 2 to 4 pounds for armyworm, corn leafhopper, corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetle, Japanese beetle, silk borer, Southwestern corn borer, and leafhoppers. For insects in wheat and foliage, apply in entire plant.
Repeat as necessary. For insects in wheat and foliage, apply in entire plant.
Repeat as necessary. For insects in wheat and foliage, apply in entire plant.
Repeat as necessary. For insects in wheat and foliage, apply in entire plant.
Repeat as necessary. For insects in wheat and foliage, apply in entire plant.

Use 4 pounds for Western bean cutworm. Test when infestation averages 10% and at 90% to 100% stand emergence. Treatment after 90% stand emergence will reduce effectiveness.
Use 4 to 13 pounds in at least 15 gallons of water per acre for cutworm control. Spray in 2-inch band using 348 pounds of ounces per 1000 linear feet of row. For broadcast ground application use up to 13 pounds in at least 20 gallons of water per acre.

COTTON

Use 1 to 2 pounds for early season control of cotton leafhopper, cotton leafworm, New Mexican striped leafhopper, and tortoise beetle.
Use 2 to 4 pounds for mid and late season control of boll weevil, budworm, cotton leafhopper, fall armyworm, leafhoppers, tarnished plant bug, and leafminer (fall armyworm/cotton worm). Use on a 5 to 7 day schedule for as long as weathers. Make 2 applications after boll open.

Use 2 to 4 pounds for light to moderate populations of high flight Western flower fly.
Use 3 to 5 pounds for pine borer.
Use 4 pounds for stink bugs and silverleaf caterpillar.

Aphid populations will be suppressed by repeated applications of this mixture.
Do not use SEVIN® brand 50W Carbaryl insecticide within 7 days of harvest.

CUCUMBERS, MELONS, PUMPKIN AND SQUASH

Use 1 to 2 pounds for pepper and melonworm. 2 pounds for cucumber beetle, flea beetle, leafhoppers and squash bug. Avoid excessive applications.
To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

DANDELION, BROKE ESCAROLE, LETTUCE AND SALAD

Use 1 to 2 pounds for flea beetles, flea hopper and leafhoppers.
Use 2 to 4 pounds in 15 to 7 day schedule for flea leafhopper, lygids bugs, and spiderbugs.
Use 2 to 4 pounds for armyworm, fall armyworm, imported cabbageworm, corn earworm, tarnished plant bug and tortoise beetle. For optimum control of armyworms after their heads begin to form, use 4 pounds in a 5 to 7 day schedule.

To avoid possible injury on lettuce, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.
Do not apply within 3 days of harvest of head lettuce and leaf lettuce.
Do not apply within 14 days of harvest of dandelion and escarole.

POPPING CRABAPPLES AND PISTACHE

Use 2 to 3 pounds for armyworm, caterpillar, leafhopper, and tortoise beetle. For optimum control of armyworms after their heads begin to form, use 4 pounds in a 5 to 7 day schedule.
Use 3 to 4 pounds for flea beetle, flea hopper and leafhopper.
Use 3 to 4 pounds for flea beetle, flea hopper and leafhopper.
Use 3 to 4 pounds for flea beetle, flea hopper and leafhopper.

GARDEN BEET, SPRING AND SUMMER CHARD

Use 1 to 2 pounds for flea beetle, flea hopper, and leafhoppers.
Use 2 to 3 pounds for flea beetle, flea hopper, and leafhoppers.
Use 2 to 3 pounds for flea beetle, flea hopper, and leafhoppers.

ORZEA

Use 2 to 4 pounds for corn earworm and stink bugs. Repeat as necessary.
Do not apply within 3 days of harvest of garden beets, chard, and spinach.
Do not apply within 14 days of harvest of garden beets, chard, and spinach.

PEANUTS

Use 1 to 2 pounds for flea beetles and Mexican leafhopper.
Use 2 to 3 pounds for flea beetles, leaf miner, leafhoppers, and tortoise beetle.
Use 2 to 3 pounds for flea beetles, leaf miner, leafhoppers, and tortoise beetle.

PEAS

Use 2 to 3 pounds for armyworm, fall armyworm, corn earworm, stink bug and leafhoppers.
Use 4 pounds for white winged beetle adults and caterpillars.
To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

POTATO, TOMATO, BEANS, PEAS AND PEPPER

Use 1 to 2 pounds for Colorado potato beetle, flea beetles and leafhoppers.
Use 2 to 4 pounds for European corn borer, fall armyworm, leafhoppers, stink bugs, corn earworm, imported cabbageworm, tarnished plant bug and tortoise beetle.
Use 4 pounds for cutworm control.

RICE

CAUTION: May kill shrimp and crabs. Do not use in areas where these shellfish are harvested. Do not use on rice fields in which shrimp and crabs are being raised for human consumption.
In the Mississippi Delta and Texas, use 2 to 3 pounds of armyworm, fall armyworm, stink bugs and stink bugs.
Do not apply PROPRINIL, HERBICIDES within 15 days before or after application of this product or plant injury will result.

ROSEHIPPUS (into grain sorghum, small sorghum, and hybrids)

Use 2 to 4 pounds for armyworm, stink bugs, corn earworm, fall armyworm, stink bugs and leafhoppers. For stink bugs, use high coverage ground application (broadcast) in 2-inch band using 348 pounds of ounces per 1000 linear feet of row.
Use 3 pounds for stink bugs, tortoise beetle and leafhoppers.

DIRECTIONS FOR USE

THIS IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION

SEVIN® brand 50W Carbaryl insecticide is a dry powder for dispersion in water and application as an insecticide spray. READ THE LABEL BEFORE USE. STRICTLY OBSERVE LABEL DIRECTIONS AND CAUTIONS, AND APPLICABLE FEDERAL AND STATE REGULATIONS.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unobscured persons.
Do not enter treated areas without protective clothing until sprays have dried.
Because certain states may require more restrictive worker protection for various crops, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be involved in the treated area or in an area about to be treated with this product. Advise workers to stay out of these areas during application and until sprays have dried. Repeat warnings after each clothing item is changed when working in treated fields. See PRECAUTIONARY STATEMENT OF PRACTICAL TREATMENT and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: Applicable signal word (CAUTION), crop area treated with SEVIN® brand 50W Carbaryl insecticide, date of application, appropriate clothing and safety material (i.e., until sprays have dried).

PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Intolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crop bearing probable carbaryl residues when carbaryl is used in accordance with label directions. Pre-harvest and grazing use limitations are listed at the end of this direction for each crop use. Where no pre-harvest or grazing use limitations are stated, application can be made to the day of harvest and crops can be grazed without five day delay if directions for use on the crops are followed. If SEVIN® brand 50W Carbaryl insecticide is used in accordance with label directions, forage and feed crops, including ground hulls, bean vicia, corn silage, citrus pulp, coffee, hay, peanut hay, pea vicia, rice straw, soybean hay and sugar beet tops, may be grazed or harvested for use as feed for dairy and meat animals without resulting in residue in milk or meat.
Do not place ripened feed and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

Do not use ripened grain or feed crops treated with carbaryl or other insecticides for which carbaryl residues are not established.

PLANT RESPONSE PRECAUTION

To avoid possible injury on tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.
Do not use on Boston ivy, Virginia creeper and man-of-war fern, as injury will result. During early season, if they are used, Virginia creeper and man-of-war fern. Observe label directions for apple thinning and/or combination with certain herbicides on rice and soybeans.

SPRAY PREPARATION

To ASSURE A UNIFORM SUSPENSION, AGITATE SPRAY CONTAINER THOROUGHLY BEFORE EACH USE. SEVIN® BRAND 50W CARBARYL INSECTICIDE SPRAY MIXTURES PRIOR TO AND DURING USE. Remove all dust scale, pesticide residues and water. Fill spray or mister tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of SEVIN® brand 50W Carbaryl insecticide and then the remaining volume of water. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURE FOR PROLONGED PERIODS.

COMPATIBILITY

SEVIN® brand 50W Carbaryl insecticide has been effectively used without plant injury with most common insecticides, miticides, fungicides, herbicides, and aquatic and air use products. If compatibility with another product and the resulting crop response is unknown, it should be tested on a small area of the growing crop before making a larger application. Observe all directions and limitations on labeling of all products used in mixture. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD SEVIN® BRAND 50W CARBARYL INSECTICIDE TO A THE DESIRED AMOUNT OF WATER, MIX THOROUGHLY, AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE. DO NOT APPLY MIX MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS, EXCESSIVE RESIDUES OR PLANT INJURY.

Unsafe under highly alkaline conditions. Not effective with certain alkaline materials such as Bordeaux, lime sulfur and calcium arsenate. Do not use with summer oil on deciduous fruits.

APPLICATION

Apply to crops in a uniform coverage to reach throughout. Use 1/2 inch nozzle for general spray application to deliver the required volume. Use 90 mesh screen on tanks of spray system and 25 mesh screen on nozzle. Adjust spray volume to deliver desired coverage. Do not spray when wind is blowing toward you. Do not spray when wind is blowing toward you.

INSECT CONTROL

Apply when there is no other damage system. The spray should be applied at 1/2 to 3/4 the desired amount of water. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary. Repeat as necessary.

FORAGE, FIELD AND VEGETABLE CROP INSECT CONTROL

For corn and sorghum, use 2 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre.

For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre.

For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre. For sorghum, use 3 to 4 pounds of SEVIN® brand 50W Carbaryl insecticide per acre.

