

Reg # 264-316 PM-19 1915

SEVIN[®] brand 80S Carbaryl Insecticide

FOR AGRICULTURAL OR COMMERCIAL USE ONLY
NOT FOR RESIDENTIAL USE

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate)80% by wt. ✓
INERT INGREDIENTS:.....20% by wt.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

E.P.A. Reg. No. 264-316

E.P.A. Est. No.

For **PRODUCT USE** information Call 1-800-334-9745

For **MEDICAL And TRANSPORTATION Emergencies ONLY** Call 24 Hours A Day 1-800-334-7577

PRECAUTIONARY STATEMENTS

WARNING

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

MAY BE FATAL IF SWALLOWED. Avoid breathing of dust or spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE.
See NOTE TO PHYSICIAN below

STATEMENT OF PRACTICAL TREATMENT

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.**

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent over atropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children

should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling Rhône-Poulenc emergency number 1-800-334-7577.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water and wetlands, except under the forest canopy and use on rice. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service or your local Rhône-Poulenc Ag Company representative for further information.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION

SEVIN[®] Brand 80S Carbaryl Insecticide is a dry powder for dispersion in water and application as an insecticidal spray in hydraulic sprayers, mist blowers, low draftage ground equipment and aircraft. **READ THIS LABEL BEFORE USE.** STRICTLY OBSERVE LABEL DIRECTIONS AND CAUTIONS, AND APPLICABLE FEDERAL AND STATE REGULATIONS.

ACCEPTED

AUG 16 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered under EPA Reg. No. 264-316

BEST AVAILABLE COPY

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENTS, STATEMENT OF PRACTICAL TREATMENT and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (WARNING), area treated with SEVIN® brand 80S Carbaryl Insecticide, date of application, appropriate clothing, and re-entry interval (i.e., until sprays have dried).

PREHARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when this product is used in accordance with label directions. If used as directed, treated forage may be grazed or used as feed for dairy and meat animals without causing illegal residues in meat or milk. This product may be applied up to and including the day of harvest or grazing of forage crops. Application may be made without removing livestock from area being treated.

Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops for which carbaryl tolerances are not established.

PLANT RESPONSE PRECAUTIONS

Avoid possible injury to tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.

Do not use on Boston Ivy, Virginia creeper and maidenhair fern as injury will result. During early season, it may also injure Virginia and sand pines.

Observe label instructions on apple thinning and on combinations with certain herbicides on rice and soybeans.

SPRAY PREPARATION

TO ASSURE A UNIFORM SUSPENSION, AGITATE, STIR OR RECIRCULATE ALL SEVIN® BRAND 80S CARBARYL INSECTICIDE SPRAY MIXTURES PRIOR TO AND DURING USE. Remove oil, rust, scale, pesticide residues and other foreign matter from

mix tanks and entire spray system. Flush with clean water. Fill spray or mix tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of SEVIN® brand 80S Carbaryl Insecticide, and then the remaining volume of water. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURES FOR PROLONGED PERIODS.

COMPATIBILITY

SEVIN® brand 80S Carbaryl Insecticide has been used without plant injury with most common insecticides, miticides, fungicides, nutrients, adjuvants and petroleum oil as used on citrus. If compatibility of SEVIN® brand 80S Carbaryl Insecticide with another product and the resulting crop response is unknown, it should be tested on a small scale. Curdling, precipitation, greasing, layer formation or increased viscosity are symptoms of incompatibility. Incompatibility will reduce insect control and may cause mechanical difficulties or plant injury. Observe all cautions and limitations on labeling of all products used in mixtures. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD SEVIN® 80S CARBARYL INSECTICIDE TO AT LEAST 1/2 THE DESIRED AMOUNT OF WATER, MIX THOROUGHLY, AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE, AND THEN THE REMAINING VOLUME OF WATER. DO NOT APPLY TANK MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS, EXCESSIVE RESIDUES OR PLANT INJURY.

Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime-sulfur and casein-lime spreaders.

APPLICATION

On all crops, use sufficient gallonage to obtain thorough and uniform coverage.

Calibrate spray equipment to deliver the required volume. Use 50 mesh slotted strainers in spray system and 25 mesh slotted strainers behind nozzles. Avoid applications just before rainfall as poor insect control may result. To clean spray system after use, drain and flush with water. Refer to the STORAGE AND DISPOSAL directions for proper disposal of rinse water.

INSECT CONTROL

Apply when insects or their damage appear. To maintain control repeat at 7 to 14 day intervals or as necessary unless a shorter interval is specified below. Where a dosage range is indicated, use lower rate on young plants and early instars and higher rate on mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

SEVIN® brand 80S Carbaryl Insecticide does not control spider mites. If spider mites are a problem, use a registered miticide.

Regional differences have been noted in the susceptibility of certain strains of fall armyworm to SEVIN® brand 80S Carbaryl Insecticide. If your experience indicates inadequate control, use an alternative pesticide.

FORAGE, FIELD AND VEGETABLE CROPS

Use at least 1 gallon of finished spray per acre for aerial application and at least 3 gallons of finished spray per acre for concentrate ground application.

GRASSHOPPERS

CROP	POUNDS OF SEVIN® BRAND 80S/ACRE	PREHARVEST INTERVAL (DAYS)	SPECIFIC DIRECTIONS
All Forage, Field and Vegetable Crops	2/3 to 1 7/8 Unless Noted Otherwise	See specific Forage, Field or Vegetable Crop	Use the Lower rate for nymphs on small plants or sparse vegetation in wasteland, rangeland, pastures, rights-of-way, ditchbanks, hedge rows and roadsides. Use the higher rate for adults or applications to dense vegetation.


**POUNDS OF PREHARVEST
SEVIN® BRAND INTERVAL
80S/ACRE (DAYS)**

CROP	INSECT	80S/ACRE	INTERVAL (DAYS)	SPECIFIC DIRECTIONS	
Alfalfa Clovers	Blister beetles Mexican bean beetle	2/3 to 1 1/4		Observe plant response precautions.	
	Alfalfa caterpillar Bean leaf beetle Cucumber beetles Green cloverworm Japanese beetle	Leafhoppers Three cornered alfalfa hopper Thrips Velvetbean caterpillar	1 1/4	7 (Alfalfa)	On dense growth use 25 to 40 gallons per acre with ground equipment to ensure adequate coverage. ✓
	Armyworm Cloverhead weevil Corn earworm Cutworm Egyptian alfalfa weevil larvae Essex skipper	European alfalfa beetle Fall armyworm Lygus bugs Webworms Yellowstriped armyworm	1 1/4 to 1 7/8	0 (Clovers)	For alfalfa weevil larvae, if pretreatment damage is extensive, cut alfalfa and treat the stubble. Apply only once per cutting. ✓
	Alfalfa weevil larvae West of the Rocky Mountains East of the Rocky Mountains	1/4 to 1 7/8 1 7/8			
Asparagus	Asparagus beetle	1 1/4 to 2 1/2	1	✓	
	Apache cicada Asparagus beetle	2 1/2 to 5	Post harvest application only	Treat ferns or brush growth. Do not treat more than once every 3 days. ✓	
Beans (including blackeyed peas, cowpeas, crowder or southern peas, dry beans, green beans, lima beans, navy beans and pinto beans)	Blister beetles Mexican bean beetle	2/3 to 1 1/4		Observe plant response precautions. ✓	
	Alfalfa caterpillar Bean leaf beetle Cucumber beetles Flea beetles Green cloverworm Japanese beetle Leafhoppers	Three cornered alfalfa hopper Thrips Velvetbean caterpillar Western bean cutworm	1 1/4	0 (except cowpeas) 3 (cowpeas)	
	Armyworm Corn earworm Cutworms European corn borer Fall armyworm	Stink bugs Tarnished plant bug Webworms	1 1/4 to 1 7/8		
	Cowpea curculio	2 1/2			
	Corn earworm Limbean pod borer	Lygus bugs Stink bugs	2 1/2	CALIFORNIA ONLY	
Cabbage Broccoli Brussels Sprouts Cauliflower Kohlrabi	Flea beetles	Harlequin bug	2/3 to 1 1/4	✓	
	Armyworm Corn earworm	Fall armyworm Imported cabbageworm	1 1/4 to 2 1/2	3	

**POUNDS OF PREHARVEST
SEVIN® BRAND
80S/ACRE INTERVAL
(DAYS)**

CROP	INSECT				SPECIFIC DIRECTIONS
Chinese cabbage Collards Hanover salad Horse radish Kale Mustard greens Radishes Rutabagas Turnips	Flea beetles Harlequin bug	Leafhoppers	2/3 to 1 1/4	3 (Horse-radish, radishes, rutabagas and turnip roots)	
	Aster leafhopper		1 1/4 to 1 7/8		
	Armyworm Corn earworm Fall armyworm Imported cabbageworm	Stink bugs Tarnished plant bug	1 1/4 to 2 1/2	14 (Chinese cabbage, collards, Hanover salad, kale, mustard greens, and turnip tops)	✓
Carrots Celery Parsnips Parsley	Flea beetles	Leafhoppers	2/3 to 1 1/4	0 (carrots)	DO NOT USE ON CELERY IN CALIFORNIA.
	Aster leafhopper Lygus bugs	Spittlebugs	1 1/4 to 1 7/8	3 (parsnips)	✓
	Armyworm Corn earworm Fall armyworm	Stink bugs Tarnished plant bugs	1 1/4 to 2 1/2	14 (celery & parsley)	
Corn (field, sweet, pop)	Armyworm Chinch bugs Corn earworm Corn rootworm adults European corn borer Fall armyworm Flea beetles	Japanese beetle Sap beetles Southwestern corn borer Leafhoppers	1 1/4 to 2 1/2	0	OBSERVE BEE CAUTION: For insects attacking silks and ears apply at 1 to 6 day intervals starting when first silks appear and continuing until silks begin to dry. For larvae in whorl and apply foliage feeders, as necessary. For chinch bugs use high gallonage ground application directed at the base of plants. Optimum timing and good coverage are essential for effective control. ✓
	Western bean cutworm		2 1/2		Treat when infestation averages 15% and at 90% to 100% tassel emergence. Treatment after 100% silk emergence will reduce effectiveness.
	Cutworms		2 1/2 to 8		Apply in a 12 inch band, using 1/4 pound (4 ounces) per 1000 linear feet of row, in at least 15 gallons of water per acre. For broadcast application use up to 8 pounds in at least 20 gallons (ground) or 5 gallons (air) of water per acre.
Cotton	Cotton fleahopper Cotton leafworm Flea beetles	Striped blister beetle Thrips	2/3 to 1 1/4	7	Early season insect control.
	Boll weevil Bollworms Cotton leaf- perforator Fall armyworm Leafrollers	Leafhoppers Tarnished plant bug Yellowstriped armyworm (cotton cutworm)	1 1/4 to 2 1/2		Treat on a 5 to 7 day schedule for as long as control is necessary. Mid and late season insect control. May be applied after bolls open. ✓
	Lygus bugs		1 1/4 to 2 1/2		For light to moderate populations in Western irrigated cotton.
	Pink bollworm		1 7/8 to 3		Aphid populations will be suppressed by repeated applications of this insecticide.
	Stink bugs Saltmarsh caterpillar		2 1/2		

POUNDS OF PREHARVEST SEVIN® BRAND 80S/ACRE INTERVAL (DAYS)

CROP	INSECT		POUNDS OF PREHARVEST SEVIN® BRAND 80S/ACRE	INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Cucumber Melons Pumpkins Squash	Pickleworm	Melonworm	2/3 to 1 1/4	0	Observe plant response precautions.
	Cucumber beetles Flea beetles	Leafhoppers Squash bugs	1 1/4		Avoid excessive applications.
Dandelion Endive (Escarole) Lettuce Salad	Flea beetles Harlequin bug	Leafhoppers	2/3 to 1 1/4	3 (head lettuce & salad roots) 14 (dandelion, endive (escarole), leaf lettuce & salad tops)	Observe plant response precautions. Treat on a 5 to 7 day schedule after heads begin to form. ✓
	Aster leafhopper Lygus bugs	Spittlebugs	1 1/4 to 1 7/8		
	Amyworm Corn earworm Fall armyworm	Imported cabbage worm Stink bugs Tarnished plant bug	1 1/4 to 2 1/2		
Pasture	Amyworm Black grass bug Chinch bugs Essex skipper Fall armyworm	Range caterpillars Range crane fly Striped grass looper Thrips	1 1/4 to 1 7/8	0 (aerial application) 14 (ground application)	To control thrips in grasses grown for seed, use high spray pressure to improve penetration into boot. Apply a maximum of 2 applications per year. Allow at least 14 days between applications. Do not allow foraging or cut for hay within 14 days of last application by ground. Aerially treated pastures may be grazed or cut for hay on day of treatment. ✓ Carefully mark swaths to avoid over-application.
	Grasshoppers				Refer to general GRASSHOPPER heading above.
Rangeland	Black grass bug Mormon cricket Range caterpillars Range crane fly		2/3 to 1 1/4	0	FOR AERIAL APPLICATION ONLY. Apply a maximum of 2 applications per year. Allow at least 14 days between applications. Carefully mark swaths to avoid over application. ✓
	Grasshoppers				Refer to general GRASSHOPPER heading above.
Wasteland, Rights-of-way, Hedgerows, Ditchbanks Roadsides	Black grass bug Mormon cricket Range caterpillar Range crane fly		2/3 to 1 1/4	0 (aerial application) 14 (ground application)	Apply a maximum of 2 applications per year. Allow at least 14 days between applications. Do not allow foraging or cut for hay within 14 days of last application by ground. Aerially treated areas may be grazed or cut for hay on day of treatment. ✓ Carefully mark swaths to avoid over-applications.
	Grasshoppers				Refer to general GRASSHOPPER heading above.
Garden beet Spinach Swiss chard	Flea beetles Harlequin bug	Leafhoppers	2/3 to 1 1/4	3 (garden beet roots)	
	Aster leafhopper		1 1/4 to 1 7/8		
	Amyworm Corn earworm Fall armyworm	Stink bugs Tarnished plant bug	1 1/4 to 2 1/2	14 (garden beet tops, spinach, Swiss chard)	

POUNDS OF PREHARVEST
SEVIN® BRAND INTERVAL
80S/ACRE (DAYS)

CROP	INSECT		POUNDS OF PREHARVEST SEVIN® BRAND 80S/ACRE	INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Okra	Corn earworm	Stink bugs	1 1/4 to 2 1/2	0	Treat on a 5 to 7 day schedule. ✓
Peanuts	Blister beetles	Mexican bean beetle	2/3 to 1 1/4	0	Observe plant response precautions. ✓
	Alfalfa caterpillar Bean leaf beetle Cucumber beetles Green cloverworm Japanese beetle Leafhoppers	Red-necked peanut worm Three cornered alfalfa hopper Thrips Velvetbean caterpillar	1 1/4		
	Amyworm Corn earworm Fall armyworm	Stinkbugs Webworms	1 1/4 to 1 7/8		
	Whitefringed beetle (adults) Cutworms		2 1/2		
Peas	Colorado potato beetle Leafhoppers		1 1/4	3	✓
	Amyworm		1 1/4 to 1 7/8		
	Alfalfa caterpillar Cutworms Pea leaf weevil	Pea weevil Yellowstriped armyworm	1 7/8		
Potato Tomato Eggplant Pepper	Colorado potato beetle Flea beetles Leafhoppers		2/3 to 1 1/4	0	✓
	European corn borer Fall armyworm Lace bugs Stink bugs	Tarnished plant bug Tomato fruit worm Tomato hornworm Tomato pinworm	1 1/2 to 2 1/2		
	Cutworms		2 1/2		
Rice	Amyworm Chinch bugs	Fall armyworm Stink bugs	1 1/4 to 1 7/8	14	CAUTION! May kill shrimp and crabs. Do not use in areas where these are important resources. Do not use on rice fields in which crayfish and/or catfish farming are included in the cultural practice. ✓
	Amyworm Leafhoppers	Tadpole shrimp	2 1/2		DO NOT APPLY PROPANIL HERBICIDES WITHIN 15 DAYS BEFORE OR AFTER APPLICATION OF THIS PRODUCT OR PLANT INJURY WILL RESULT. ✓ MISSISSIPPI DELTA & TEXAS CALIFORNIA ONLY: For optimum tadpole shrimp control, apply to water when pest first appears.

BEST AVAILABLE COPY

**POUNDS OF PREHARVEST
SEVIN® BRAND INTERVAL
80S/ACRE (DAYS)**

CROP	INSECT		POUNDS OF PREHARVEST SEVIN® BRAND INTERVAL 80S/ACRE	(DAYS)	SPECIFIC DIRECTIONS
Sorghums (milo, grain sorghum, sweet sorghum and hybrid)	Amyworm Chinch bugs Corn earworm	Fall armyworm Stink bugs Webworms	1 1/4 to 2 1/2	21 (grain) 0 (forage)	Direct spray into forming heads for optimum insect control. Treat for sorghum midge when 25 to 30 percent of heads have emerged from boot and are in bloom. Repeat application 3 to 5 days later if adults are still active. A third application may be necessary in late planted sorghum or if midge are abundant. For chinch bugs use high gallonage ground application directed at base of plants.
	Sorghum midge Southwestern corn borer		1 7/8		
	Cutworms		2 1/2		
Soybeans	Bean leaf beetle Cucumber beetles Green cloverworm	Mexican bean beetle Velvetbean caterpillar	2/3	0	DO NOT APPLY A COMBINATION OF THIS PRODUCT AND 2,4-DB HERBICIDES TO SOYBEANS AS CROP INJURY MAY RESULT. Use lower rates for light to moderate populations and smaller instars. Use the higher rates for heavy populations and larger instars.
	Corn earworm		2/3 to 1 1 1/4 to 1 7/8		
	Blister beetles Grape colaspis Mexican bean beetle		2/3 to 1 1/4		
	Alfalfa caterpillar Bean leaf beetle Cucumber beetles Green cloverworm Japanese beetle	Leafhoppers Three cornered alfalfa hopper Thrips Velvetbean caterpillar	1 1/4		
	Amyworm Cutworms	Fall armyworm Stink bugs Webworms	1 1/4 to 1 7/8		
	Painted lady (Thistle caterpillar)	Saltmarsh caterpillar Yellowstriped armyworm	2 1/2 to 3		
Sugar beets	Amyworm Beet leaf beetle Fall armyworm	Flea beetles Leafhoppers Webworms	1 1/4 to 1 7/8	14	
	Cutworms		1 7/8		
Sunflower	Cutworms		1 7/8	60	DO NOT USE IN CALIFORNIA. Do not allow animals to graze on treated crops.
	Amyworm Fall armyworm		1 7/8 to 2 1/2		
	Stem Weevil Sunflower beetle		1 1/4 to 2 1/2		
Sweet Potatoes	Sweet potato weevil		9 tablespoons/ gallon.	0	DO NOT USE IN CALIFORNIA Prelant Dip: Just prior to planting, dip sweet potato cuttings in a suspension of this product (9 tablespoons/gallon of water). Foliar Application: Full coverage of plants is essential. Use lower rate on young plants and higher rate on mature plants. Apply as a foliar spray as needed.
			1 1/4 to 2 1/2		
	Corn earworm Cucumber beetles Flea beetles	Sweet potato hornworm Tortoise beetles	1 1/4 to 2 1/2		

BEST AVAILABLE COPY

8915
7

CROP	INSECT	POUNDS OF PREHARVEST SEVIN® BRAND		SPECIFIC DIRECTIONS
		80S/ACRE	INTERVAL (DAYS)	
Tobacco (In Plant Beds)	Tobacco flea beetle	2 1/2 lbs/gal. or 1 1/4 lbs / 50 gallons and apply 6 gal / 100 sq. yards	3	To prevent plant injury, avoid excessive applications.
	Green June beetle grubs	2/3 lb./200 gal.		Apply only to areas that larvae have uprooted by sprinkling as a drench with 50 to 100 gallons per 100 square yards.
(In Fields)	Budworms Fall armyworm Flea beetles Homworms	Japanese June beetles Suckfly	1 1/4 to 2 1/2	Use lower rate on young plants (up to knee height). Use at least 10 gallons of prepared spray per acre. Begin treatments when worms are small. ✓ If late season applications are necessary, allow three days before priming or cutting.
Wheat	Cereal leaf beetle	1 1/4	0 (Forage)	DO NOT USE IN CALIFORNIA. Do not make more than two applications after grain heads emerge from boot. ✓
	Armyworm Fall armyworm	1 1/4 to 1 7/8	21 (grain)	Application is effective against eggs larvae and adults.

TREE FRUIT AND NUT CROPS

For dilute sprays apply the specified dosage per 100 gallons of water. For concentrate and aerial sprays increase the concentration of SEVIN® brand 80S Carbaryl Insecticide in the spray mixture to apply an amount per acre equivalent to that in a dilute spray. The optimum spray gallonage will depend on tree size, density and stage of growth. Typical spray gallonages per acre range from 200-300 gallons for dilute sprays, 30-100 gallons for concentrate sprays and 10-25 gallons for aerial sprays. Do not exceed maximum label rate per acre per application.

CROP	POUNDS OF SEVIN® BRAND 80S/100 GAL	PREHARVEST INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Apples only	1/3 to 2/3	1	Apply 1 full coverage dilute spray between 10 and 25 days after full bloom. If factors such as tree age, variety, nutrition, previous crop, pruning, bloom and degree of set favor excessive fruit thinning with this product, exercise caution to avoid possible yield reduction. Consult with your County Extension Service or other experts for advice on the proper use of this product. For easily thinned varieties including Cortland, Grimes, Jonathan, McIntosh, Orleans, Rome Beauty, Puritan, Red Delicious, Winesap, Yellow Newton.
	2/3 to 1 1/4		For difficult to thin varieties including Baldwin, Ben Davis, Duchess, Early McIntosh, Golden Delicious, Lady Apple, Northern Spy, Rhode Island Greening, Stark Red, Tiffin, Wealthy, Yellow Transparent and York Imperial.

CROP	INSECT	POUNDS OF PREHARVEST SEVIN® BRAND		SPECIFIC DIRECTIONS
		80S/100 GAL	INTERVAL (DAYS)	
Almond	Peach twig borer San Jose scale Fruitree leafroller	1 1/4	28	OBSERVE BEE CAUTION For peach twig borer, apply in "popcorn" or petal fall stages when the May brood begins to hatch or there after as needed. For dormant or delayed dormant timing, apply in combination with a recommended dormant oil. For Navel orangeworm, time early and mid season applications to correspond to moth flight peaks. Make a late season application at initiation of hull split up to 10% hull split. Do not apply more than 6 1/4 pounds per acre.
	Naval orangeworm			

* Refer to the dormant oil label for additional use directions and restrictions.

**POUNDS OF PREHARVEST
SEVIN® BRAND INTERVAL
80S/100 GAL. (DAYS)**

CROP	INSECT			SPECIFIC DIRECTIONS
Apples Pears	Apple aphid Apple rust mite Apple sucker Bagworms California pearslug (pear sawfly) Codling moth Eyespotted bud moth Green fruitworm Lecanium scales Lesser appleworm	Lygus bugs Orange tortrix Oystershell scale Pear leaf blister mite Pear psylla Pear rust mite San Jose scale Tentiform leafminers Woolly apple aphid	1 to 1 1/4	WEST OF THE ROCKY MOUNTAINS. To avoid undesired apple thinning, delay use until at least 30 days after full bloom. For psylla control, apply when eggs hatch or young nymphs are present.
	Apple mealybug Apple aphid Codling moth	White apple leafhopper	2/3	EAST OF THE ROCKY MOUNTAINS. To avoid undesired apple thinning, delay use until at least 30 days after full bloom. For optimum scale control, apply when crawlers are present.
	Apple maggot Apple rust mite Bagworms Eastern tent caterpillar European apple sawfly Eyespotted bud moth Forbes scale Fruitree leafroller Green fruitworm Japanese beetle Lecanium scales Lesser appleworm Oystershell scale	Pear psylla Pear leaf blister mite Pear rust mite Periodical cicada Plum curculio Redbanded leafroller Rosy apple aphid San Jose scale Tarnished plant bug Tentiform leafminers Woolly apple aphid Yellowheaded fireworm	1 1/4	
Chestnuts	Chestnut weevil		2 1/2 to 3 3/4	DO NOT USE IN CALIFORNIA Make 4 applications at weekly intervals beginning in late July for adult chestnut weevil control. Last application should be made prior to shuck split.
Citrus Fruits (such as grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats and hybrids)	Avocado leafroller California orangedog Citrus cutworm Citrus root weevil Fruitree leafroller	Orange tortrix Western tussock moth West Indian sugarcane borer (adults)	1 1/4	5 Do not apply more than 25 pounds of this product per acre per application. May be mixed with petroleum oils commonly used on citrus.
	Black scale Brown soft scale California red scale	Citricola scale Citrus snow scale Yellow scale	1 to 1 1/4	
Filbert	Filbert aphid Filbert leafroller Filbertworm		1 1/4	0 Apply when leafroller eggs are hatching. Repeat on first appearance of adult filbert moths and again 3 to 4 weeks later.
Olives	Olive scale		1 to 1 1/4	0 For optimum scale control, add 1 1/2 gallons of summer oil and apply mixture when crawlers are present. Do not exceed 2 applications per year. Do not apply more than 18 3/4 pounds of this product per acre per application.

**POUNDS OF PREHARVEST
SEVIN® BRAND INTERVAL
80S/100 GAL. (DAYS)**

CROP	INSECT		POUNDS OF PREHARVEST SEVIN® BRAND INTERVAL 80S/100 GAL. (DAYS)		SPECIFIC DIRECTIONS
Peaches Apricots Nectarines	Apple pandemis	Orange tortrix	1 1/4	1 (peaches)	Do not apply more than 7 1/2 pounds of this product per acre per application to apricots. For optimum scale control, apply when crawlers are present. Spray limbs and trunk thoroughly, weekly during moth flight.
	Codling moth	Oriental fruit moth		3 (apricots & nectarines)	
Pecans	Cucumber beetles	Peach twig borer	1 1/4 to 3	0	Do not apply more than 9 pounds of this product per acre per application.
	European earwig	Periodical cicada			
Pistachios	Fruitree leafroller	Plum curculio	2/3 to 2 1/2	14	Apply dilute volumes of 150 to 300 gallons of mixed spray per acre for full coverage.
	Japanese beetle	Redbanded leafroller			
Plums Prunes cherries	June beetles	San Jose scale	1 1/4	1	Do not apply more than 7 1/2 pounds of this product per acre per application. For optimum scale control, apply when crawlers are present. For lesser peachtree borer control, spray limbs and tree trunks thoroughly, weekly during moth flight.
	Lecanium scales	Tarnished plant bug			
Walnut	Lesser peachtree borer	Tussock moths	1	0	Apply 1000 gallons of dilute spray per acre for mature trees. For codling moth apply first spray when average cross sectional diameters of developing nuts are 1/2 to 3/4 inch. Repeat during middle or late June as needed.
	Olive scale	Variegated leafroller			
Walnut	Black margined aphid	Pecan nut casebearer	2/3	0	Spray tree trunks to point of run-off.
	Fall webworm	Pecan spittlebug			
Walnut	Hickory shuckworm	Pecan stem phylloxera	2 1/2		
	Lesser webworm	Pecan weevil			
Walnut	Pecan leaf phylloxera	Twig girdler			
		Walnut caterpillar			
	Navel orangeworm				
	Black cherry aphid	Mealy plum aphid			
	Brown soft scale	Oystershell scale			
	Cherry fruitworm	Peach twig borer			
	Cherry maggot	Plum curculio			
	European earwig	Prune leafhopper			
	Eyespotted bud moth	Redbanded leafroller			
	Forbes scale	Rose chafer			
	Fruitree leafroller	San Jose scale			
	Green fruitworm	Variegated leafroller			
	Japanese beetle				
	Lecanium scales				
	Lesser peach-tree borer				
	Codling moth	Orange tortrix			
	Eastern tent caterpillar	Tussock moth			
	Calico scale	Fibertworm			
	European fruit lecanium	Fruitree leafroller			
	Codling moth	Frosted scale			


BEST AVAILABLE COPY

SMALL FRUIT CROPS

11915

Recommended dosages refer to pounds of SEVIN® Brand 80S Carbaryl insecticide per acre. The optimum spray gallonage will depend on plant size, density and stage of growth. Typical spray gallonages per acre range from 100-300 gallons for dilute sprays, 30-100 gallons for concentrate sprays and 10-25 gallons for aerial sprays. Do not exceed maximum label rate per acre per application.

CROP	INSECT	POUNDS OF PREHARVEST SEVIN® BRAND 80S/ACRE		INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Blackberries Raspberries Dewberries (including boysenberries and loganberries)	European raspberry aphid Japanese beetle	Leafhoppers Leafrollers Rose chafer Snowy tree cricket	1 1/4 to 2 1/2	7	CALIFORNIA ONLY
	Omnivorous leafroller Raspberry sawfly		2 1/2		
Blueberries	Blueberry maggot Cherry fruitworm Cranberry fruitworm	European fruit lecanium Japanese beetle	1 7/8	0	Apply 3 weeks before harvest and repeat as necessary.
Cranberries	Cutworms Cranberry fireworms Cranberry fruitworms	Japanese beetle Leafhoppers Rose chafer	1 7/8 to 3 3/5	1	CAUTION! May kill shrimp and crabs. Do not use in areas where these are important resources. Apply in late bloom and as needed at 7 to 10 day intervals.
Grapes	European fruit lecanium Grape leafroller Grape leafhopper	Western grapeleaf skeletonizer Western yellow-striped armyworm	1 1/4 to 2 1/2	7	Apply before first brood leafroller larvae emerge from rolls.
	Cutworms Eight-spotted forester Grape berry moth Japanese beetle June beetles	Orange tortrix Omnivorous leafroller Redbanded leafroller Saltmarsh caterpillar	2 1/2		
Strawberries	Flea beetles Meadow spittlebug Omnivorous leaflier (strawberry fruitworm)	Strawberry leafroller Strawberry weevil	1 1/2 to 2 1/2	1	Carbaryl may injure Early Dawn and Sunrise varieties on the Delmarva Peninsula.



BEST AVAILABLE COPY

NON-RESIDENTIAL TREES AND ORNAMENTALS

12/15

For dilute-spray ground applications to non-residential trees (including shade trees, shelter belts, forests, plantations, parks, and recreational areas), non-residential ornamentals, woody plants and shrubs, apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage.

For aerial applications to forest trees (including shade trees, shelter belts, plantations, parks, and recreational areas) and commercially grown ornamentals, woody plants and shrubs, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage. Avoid direct application to lakes, streams and ponds.

CROP APPLICATION:	INSECT			POUNDS OF SEVIN® BRAND 80S/ 100 GALLONS	SPECIFIC DIRECTIONS
	GROUND APPLICATION: Trees (including shade trees, shelter belts, forests, plantations, parks and recreational areas). Ornamentals including roses, flowers and other herbaceous plants). Woody Plants and Shrubs.	Ants Apple aphid Armyworm Azalea leafminer Bagworms Birch leafminer Blister beetle Boxelder bug Boxwood leafminer Brown tail moth Cankerworms Catalpa sphinx Chiggers Cooley spruce gall aphid Cutworms Cypress tip moth Douglas-fir tussock moth Eastern spruce gall aphid Elm leaf aphid Elm leaf beetle Elm spanworm Eriophyid mites European pine shoot moth Fall armyworm Flea beetles Fuller rose beetle Gall midges Gall wasps Green striped mapleworm Grasshoppers Gypsey moth Hackberry nipplegall maker	Holly bud moth Holly leafminer Jackpine budworm Japanese beetle Jeffrey pine needleminer June beetles Lace bugs Leafhoppers Leafrollers Locust borer Maple leafcutter Mealy bugs Mimosa webworm Nantucket pine tip moth Oak leafminers Oak leaf skeletonizer Oakworm complex Oleander caterpillar Olive ash borer Orange-striped oakworm Orange tortrix Periodical cicada Pine sawfly Pine spittlebug Pitch pine tip moth Plant bugs Poinsettia hornworm Psyllids Puss caterpillar Redhumped oakworm	Rose aphid Rose chafar Roselug Saddled prominent Sawflies (exposed) Scale insects Sowbugs Spiny elm caterpillar Springtails Spruce budworm Spruce needleminer Subtropical tip moth Tent caterpillars Thorn bug Thrips (exposed) Ticks Walnut caterpillar Webworms Western hemlock looper Western spruce budworm Willow leaf beetles Yellow poplar weevil	1 1/4
AERIAL APPLICATION Forest Trees (including shade trees, shelter belts, forests, plantations, parks and recreational areas) and Commercially Grown Ornamentals including roses, flowers and other herbaceous plants). Woody Plants and Shrubs	Ips engraver beetles Mountain pine beetle Roundheaded pine beetle Western pine beetle			25	Effective as a preventive treatment only. Repeat annually as required to prevent beetle attacks. Apply 1 gallon of spray per 50 square feet of bark in late May to early June or prior to beetle attack. Treat tree trunks from ground level up, until trunk diameter is less than 5 inches.
	Elm bark beetle				Apply approximately 20 to 30 gallons of spray mixture for each 50 feet of elm tree for thorough coverage of all bark surfaces on trunk, limbs and twigs.

NON-RESIDENTIAL TURF AND RECREATIONAL AREAS

13 15

CROP	INSECT		SPECIFIC DIRECTIONS
Turf grasses	Ants Armyworm Bluegrass billbug Centipedes Chiggers Chinch bugs Cutworms Earwigs Essex skipper European chafer Fall armyworm Fiery skipper Fleas	Grasshoppers June beetles Leafhoppers Lucerne moth Millipedes Mosquitoes Sod webworm (lawn moths) Sowbugs Springtails Ticks White grubs Yellowstriped armyworm	Use 4 ounces of SEVIN® brand 80S Carbaryl insecticide per 1000 square feet (10 3/4 pounds per acre) of turf grass. Make applications in sufficient spray volume for thorough coverage and turf thatch penetration. Repeat treatment as necessary. <u>For Armyworm, Cutworms:</u> Fall Armyworm and Sod Webworm Control: Do not irrigate treated areas following insecticide application. <u>For Chinch Bug Control:</u> Treat entire turf grass area rather than just damaged areas. Irrigation of turf grass area before insecticide application will aid in penetration into turf grass. <u>For White Grub Control:</u> Applications should be made when grubs are feeding near the soil surface, usually during late March through May, or July to early September, or as recommended by local Agricultural Extension Service agents. Water or irrigate turf grasses thoroughly soon after treatment.

IMPORTED FIRE ANT CONTROL

SITE	POUNDS OF SEVIN® BRAND 80S/100 GALLONS OF WATER		SPECIFIC DIRECTIONS
Cemeteries, and non-residential Recreational Areas (including turf, golf courses, and parks) Pastures, Rangeland, Forested Lands, Wastelands		2	Apply a total of 2 gallons of the diluted solution over the surface of each mound or at least 1 quart per 6 inches of mound diameter using a bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). Do not disturb mounds prior to treatment. Pour solution from a height of about three feet to give sufficient force to break mound apex and flow into ant tunnels. For best results, apply in cool weather 65-80° F, or in early morning or late evening hours. Repeat application if mound activity resumes after 10 days. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reduction in product effectiveness.

PEST CONTROL IN AND AROUND NON-RESIDENTIAL BUILDINGS LIMITED TO PEST CONTROL OPERATORS ONLY

INSECT	POUNDS OF SEVIN® BRAND 80S/100 GAL.		SPECIFIC DIRECTIONS
Ants Cockroaches		25 (25 tablespoonfuls/gallon)	For use in and around non-residential buildings such as warehouses, barns, and municipal and recreational areas.
Brown dog tick, Earwigs Fleas, Millipedes, Silverfish, Spiders		12 1/2 (12 1/2 tablespoonfuls/gallon)	Apply as a coarse wet spray or with a paint brush, and thoroughly treat cracks and crevices, baseboards, corners, walls, door sills, window frames, ant trails, and other areas where insects tend to congregate. Do not use as a space spray. Repeat when necessary but not more often than twice per week. Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed. Do not use in serving areas while food is exposed. Avoid application to surfaces where visible spray residues are objectionable.

BEST AVAILABLE COPY

14 9 15

POULTRY INSECT CONTROL

POULTRY	INSECT	SPECIFIC DIRECTIONS
Chickens, Ducks, Geese, Game birds, Pigeons, and Turkeys	Chicken mites Fleas Lice Northern fowl mite	For use as a direct spray on birds by: 1. Misting with Electric Fog Machine: Mix 3/8 pounds (6 ounces) of this product in 1 gallon of spray. Use 1 1/2 gallons per 1000 hens in cages, or litter or on slatted floors. Repeat in 4 weeks if necessary. or 2. Spraying with knapsack or cylinder type compressed air sprayers: Mix 1/4 pound (4 ounces) of this product in 5 gallons of spray. Use 1 gallon per 100 hens in cages, on litter or on slatted floors. Repeat in 4 weeks if necessary. Direct mist spraying for chicken mite and fleas in a supplement to spraying roosts and buildings for control of these pests. Do not apply to poultry and game birds within 7 days of slaughter.

POULTRY QUARTERS	INSECT	SPECIFIC DIRECTIONS
Roosts and Buildings	Bedbugs Chicken mites Fleas Fowl tick Darkling Beetle (Lesser mealworms) Lice Northern fowl mite	Spray roosts and buildings with conventional power spray or knapsack equipment. For chicken mite, fleas and bedbugs, use 5 pounds of this product per 100 gallons of water. For fowl ticks, use 20 pounds of this product per 100 gallons of water. For lesser mealworms, use 62 1/2 pounds per 100 gallons. Spray 1 to 2 gallons per 1000 square feet of wall, bedding, litter or roost surface. Force spray into cracks. Repeat as needed. Avoid spraying of nests, eggs and feeding and watering troughs. Ventilate while spraying. Do not treat premises within 7 days of slaughter.

DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including center pivot and solid set. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of SEVIN® brand 80S Carbaryl Insecticide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of SEVIN® brand 80S, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of SEVIN® brand 80S per 1 to 4 gallons of water are recommended). Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of SEVIN® brand 80S into the irrigation water line so as to deliver the desired rate per acre. The suspension of SEVIN® brand 80S should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with SEVIN® brand 80S has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief

valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines, and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

STORAGE AND DISPOSAL

STORAGE

Store unused SEVIN® brand 80S Carbaryl insecticide in original container only, in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F.

PESTICIDE DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If bag is burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER

MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

THIS SPECIMEN LABEL IS INTENDED FOR USE ONLY AS A GUIDE IN PROVIDING GENERAL INFORMATION REGARDING THE DIRECTIONS, WARNINGS AND CAUTIONS ASSOCIATED WITH THE USE OF THIS PRODUCT. AS WITH ANY AGRICULTURAL CHEMICAL, ALWAYS FOLLOW THE LABEL INSTRUCTIONS ON THE PACKAGE BEFORE USING.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

SEVIN is the registered trademark of RHONE-POULENC for carbaryl insecticides.
©1989 Rhône-Poulenc Ag Company
Made in USA.

Form No. AS90055
Printed USA 9/89

EPA Approval: 8/22/88