

RESTRICTED USE PESTICIDE: For retail sale to and use by certified applicators only, and only under their direct supervision and only for those uses covered by the certification of the applicator.

PRODUCT BULLETIN




UNION CARBIDE AGRICULTURAL
PRODUCTS COMPANY, INC.
T. W. ALEXANDER DRIVE
P.O. BOX 12014
RESEARCH TRIANGLE PARK, NC 27709

TEMJK 15G Aldicarb Pesticide
EPA Reg. No. 264-330

Page 1 of 2

Supplemental Labeling For Control of Insects and Nematodes on Potatoes

Crop & Time of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet Of Row	Recommended Application
34-Inch Row Spacing				
POTATOES AT PLANTING Do Not Apply In Suffolk County, Long Island, New York, At Anytime.	Aphids	7 (Maine Only)	7	Apply Granules With Seed Pieces In Planting Furrow and Cover With Soil.
	Flea Beetles	20	21	
Do Not Apply At Planting In Wisconsin or Maine (Except 7lb Rate for Aphids in Maine Only)	Aphids, Colorado Potato Beetle, Leafhoppers	14 to 20	14.5 to 21	Apply As Above Or, Drill Granules 2 To 4 Inches On Both Sides of Seed Row And 3 To 8 Inches Deep (Usually 1 To 2 Inches Below Seed Pieces).
	Psyllids	14 (Arizona Only)	14.5	Apply As Above.
	Nematodes (Except Golden Nematode) (Root-Knot Nematodes Are Suppressed) (Verticillium Wilt Has Been Suppressed Where Nematode Populations Have Been Reduced)	20	21	Apply Granules With Seed Pieces In Planting Furrow And Cover With Soil Or, Apply Granules In An 8-Inch Band And Work Into The Soil Or Cover With Soil To A Depth Of 4 Inches. Plant Seed Pieces In Treated Zone.
AFTER PLANTING In Maine and Wisconsin, Apply 4-6 Weeks After Planting.	Aphids, Colorado Potato Beetle, Leafhoppers	14 (Maine And Wisconsin Only)	14.5	Apply Granules in 4-6 Inch Band Over Row On Top Of Hill. Cover Granules With Soil By Cultivators Or Hilling Discs. DO NOT APPLY THIS PRODUCT TO ANY FIELD MORE THAN ONCE EVERY TWO YEARS.



RESTRICTED USE PESTICIDE

TEMI- 15G Aldicarb Pesticide
EPA Reg. No. 264-330

Page 2 of 2

Supplemental Labeling
For Control of Insects and Nematodes on Potatoes

Crop & Time of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
POTATOES				
AFTER PLANTING (continued)				
Other Than Maine And Wisconsin, Apply At 75 Emergence Or Before Plants Are 6 Inches High	Aphids, Colorado Potato Beetle	14 to 20	14.5 to 21	Side-Dress Granules 4 To 5 Inches On Both Sides Of Plant Row And 4 Inches Deep. Apply Where Pest Populations Are Expected 2-3 Weeks After Plant Emergence. Do Not Apply If At Planting Treatment Was Made.

PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS:

- Do not make more than one application per crop.
- In Maine and Wisconsin, do not apply to any field more than once every two years.
- Do not harvest within 90 days of an at planting application.
- Do not apply an "after planting" application within 50 days of harvest.
- Do not apply this product in Suffolk County, Long Island, New York.

DECOMPOSITION AND MOVEMENT IN SOIL

This product is readily decomposed into harmless residues under most use conditions: However, a combination of sandy and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F at application time, tend to reduce degradation and promote movement of residues to groundwater. If this describes your local use conditions and groundwater in your area is used for drinking, do not use this product without first contacting Union Carbide at 1 (800) 334-9769.

IMPORTANT:

Before using TEMIK 15G Aldicarb Pesticide, read and carefully observe the precautionary statements and all other information appearing on the product label. This bulletin contains new and supplemental directions for use of this product which may not appear on the package label. Follow the directions carefully.

3705

0404