

564-230 PM12 1/12

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATORS CERTIFICATION.

TEMIK® 15G
ALDICARB PESTICIDE

For Control of Certain Insects, Mites and Nematodes

Active ingredient: Aldicarb [2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl)oxime]	15%
Inert ingredients	85%

EPA Reg. No. 264-330

EPA Est. No. 10352-GA-01

©UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
P.O. Box 12014, T.W. Alexander Drive
Research Triangle Park, NC 27709

TEMIK is the registered trademark of Union Carbide Corporation for aldicarb pesticide.


MADE IN U.S.A.

APC-2448-473051-12²L

ACCEPTED

MAY 31 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under
EPA Reg. No. 264-330



KEEP OUT OF REACH OF CHILDREN

D A N G E R P O I S O N

PELIGRO

PRECAUCION AL USARIO: Si usted no lee ingles no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION TO THE USER: IF YOU CANNOT READ ENGLISH DO NOT USE THIS PRODUCT UNTIL THE LABEL HAS BEEN FULLY EXPLAINED TO YOU.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

POISONOUS IF SWALLOWED. MAY BE FATAL OR HARMFUL BY SKIN OR EYE CONTACT OR BY BREATHING DUST. RAPIDLY ABSORBED THROUGH SKIN OR EYES. DO NOT GET ON SKIN OR EYES. DO NOT BREATHE DUST.

WEAR LONG-SLEEVED CLOTHING AND PROTECTIVE GLOVES WHEN HANDLING. WASH HANDS AND FACE BEFORE EATING OR SMOKING. BATHE AT THE END OF WORK DAY, WASHING ENTIRE BODY AND HAIR WITH SOAP AND WATER. CHANGE CLOTHING DAILY. WASH CONTAMINATED CLOTHING IN STRONG WASHING SODA SOLUTION AND RINSE THOROUGHLY BEFORE REUSING.

SIGNS AND SYMPTOMS OF OVEREXPOSURE:

Salivation	Muscle tremor	Nausea
Watery eyes	Difficult breathing	Vomiting
Pinpoint eye pupils	Excessive sweating	Diarrhea
Blurred vision	Abdominal cramps	Weakness
	Headache	

In severe cases, convulsions, unconsciousness and respiratory failure may occur.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

STATEMENT OF PRACTICAL TREATMENT

GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately, and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never give anything by mouth to an unconscious or convulsing person. Call a physician and follow General advice listed above.

IF IN EYES: Flush eyes with plenty of water and get medical attention.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Call a physician and follow General advice listed above.

NOTE TO PHYSICIAN

TEMIK® aldicarb is a methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Union Carbide Agricultural Products Company.

Consultation on therapy can be obtained at all hours by calling the Union Carbide emergency number: (304) 744-3487.

ENVIRONMENTAL HAZARDS

DECOMPOSITION AND MOVEMENT IN SOIL

This product is readily decomposed into harmless residues under most use conditions. However, a combination of permeable and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F at application time, tend to reduce degradation and promote movement of residues to groundwater. If this describes your local use conditions and groundwater in your area is used for drinking, do not use this product without first contacting Union Carbide at 1(800) 334-8577.

TOXIC TO FISH, BIRDS, AND WILDLIFE

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

11/14

NOTICE: Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

This Act protects Attwater's Greater Prairie Chicken in the Texas counties of Aransas, Austin, Brazoria, Colorado, Galveston, Goliad, Harris, Refugio, and Victoria.

Prior to making applications in these counties, the user must determine that this species is not located in or immediately adjacent to the area to be treated. If the user is in doubt whether or not the above named endangered species may be affected, he should contact either the regional U.S. Fish & Wildlife Service office (Endangered Species Specialist) or personnel of the State Fish and Game office.

IN CASE OF EMERGENCY TELEPHONE COLLECT
(24 HOURS A DAY) IN U.S.A. (304) 744-3487

GENERAL DIRECTIONS FOR USE

GENERAL INFORMATION


READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

TEMIK® 15G Aldicarb Pesticide controls certain insects, mites and nematodes. When applied into moist soil at-planting and/or postemergence the active ingredient is rapidly absorbed by roots and translocated throughout all parts of the plant. Control often lasts more than six weeks varying with growing conditions, rate of use and pests. Crop yields are usually increased with treatments of TEMIK®.

GENERAL CAUTIONS

Use TEMIK® 15G Aldicarb Pesticide only in accordance with label directions, warnings and cautions. Do not use on any crop not listed on this label or supplemental labeling as any residues remaining may be illegal or harmful. DO NOT STORE OR USE IN OR AROUND THE HOME OR HOME GARDEN. Use high rate on heavy organic or clay soils. Do not over-apply. Treatments in excess of 7 pounds per acre made directly in the seed furrow of cotton, sugar beets, soybeans or drybeans may delay emergence and reduce plant stand.

Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, the TEMIK® and water must be on the same side of the plant row.



Calibrate and adjust application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. Bury any excess material in soil (see instructions for STORAGE AND DISPOSAL in this booklet).

Deep disc any spills at row ends immediately to prevent birds from feeding on exposed granules.

DO NOT APPLY THIS PRODUCT IN SUFFOLK COUNTY, LONG ISLAND, NEW YORK.


COMPATIBILITY

Pesticidal activity of TEMIK® 15G Aldicarb Pesticide is not affected by normal applications of fertilizers or other pesticides. It may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix TEMIK® Aldicarb Pesticide with other materials before application.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated area without protective footgear. Persons coming in direct contact with wet treated soil after the initial irrigation or rainfall following treatment shall wear coveralls and rubber or neoprene boots and gloves. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application. Regular long-sleeved work clothing should be worn when working in treated fields. See Precautionary Statement, Statement of Practical Treatment and Note To Physician for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (DANGER-POISON), area treated with TEMIK® 15G Aldicarb Pesticide, date of application, appropriate protective clothing, and re-entry requirements.



17/10

STORAGE AND DISPOSAL

STORAGE

STORE UNUSED TEMIK® ALDICARB PESTICIDE IN ORIGINAL CONTAINER ONLY, IN WELL VENTILATED CLEAN DRY AREA OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT STORE IN AREAS WHERE TEMPERATURE AVERAGES 115°F OR GREATER. DO NOT STORE IN OR AROUND THE HOME OR HOME GARDEN.

IF CONTAINER IS BROKEN, HANDLE WITH RUBBER GLOVES. DO NOT GET DUST OR GRANULES ON SKIN OR EYES. DO NOT BREATHE DUST. SWEEP UP AND BURY ANY SMALL SPILLS OR EXCESS TEMIK® AT LEAST 18 INCHES DEEP IN SOIL ISOLATED FROM WATER SUPPLIES AND FOOD CROPS.

PESTICIDE DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.


OPEN DUMPING IS PROHIBITED.

PESTICIDE WASTES ARE ACUTELY HAZARDOUS. IMPROPER DISPOSAL OF EXCESS PESTICIDE IS A VIOLATION OF FEDERAL LAW. IF THESE WASTES CANNOT BE DISPOSED OF BY USE ACCORDING TO LABEL INSTRUCTIONS, CONTACT YOUR STATE PESTICIDE OR ENVIRONMENTAL CONTROL AGENCY OR THE HAZARDOUS WASTE REPRESENTATIVE AT THE NEAREST EPA REGIONAL OFFICE FOR GUIDANCE.

CONTAINER DISPOSAL

COMPLETELY EMPTY CONTAINER INTO APPLICATION EQUIPMENT. THEN DISPOSE OF EMPTY CONTAINER IN A SANITARY LANDFILL OR BY INCINERATION, OR IF ALLOWED BY STATE AND LOCAL AUTHORITIES, BY OPEN BURNING. IF CONTAINER IS BURNED, STAY AWAY FROM AND DO NOT BREATHE OR CONTACT SMOKE.

IN CASE OF TRANSPORTATION OR WAREHOUSE EMERGENCY INVOLVING A SPILL, FIRE, OR EXPOSURE, CALL COLLECT (304) 744-3487 TWENTY-FOUR HOURS A DAY IN THE USA.



PRE-HARVEST, GRAZING, AND ENVIRONMENTAL LIMITATIONS

To avoid excessive residues in or on:

COTTONSEED

- Do not make more than one at-planting application and one postemergence application per crop.
- Do not apply within 90 days of harvest.

POTATOES

- Do not make more than one application per crop.
- Do not harvest within 90 days of an at-planting application or within 50 days of an 'after-planting' application.

PEANUTS


- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not hog off treated fields.
- Do not feed hay or vines to livestock.

SUGAR BEETS

- Do not make more than one at-planting application and two postemergence applications per crop.
- Do not exceed a total of 40 pounds per acre.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.

SOYBEANS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay or straw to livestock.



DRYBEANS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay or straw to livestock.
- Do not use green pods as food for humans.

SUGARCANE AND SWEET POTATOES

- Do not make more than one application per crop.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.

PECANS

- Do not make more than one application per year.
- Do not allow livestock to graze in treated areas.
- Do not harvest foliage or hay from treated areas.

SORGHUM

- Do not make more than one application per year.
- Do not harvest within 90 days of application.
- Do not feed green forage to livestock.

OTHER CROPS


- Do not plant any crops not listed on this label in soil treated with TEMIK® 15G Aldicarb pesticide within six months after last application.
- Do not apply this product in Suffolk County, Long Island, New York.

MEAT AND MILK

- Do not allow livestock to graze in treated areas before harvest.

GROUNDWATER

- Observe Environmental Hazard precautions regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- Do Not apply in Suffolk County, Long Island, New York.




DIRECTIONS FOR USE AND RECOMMENDED APPLICATIONS

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THE LABELING. To provide maximum benefit and to minimize hazard to birds, work granules into or cover with soil to a depth of 2 inches or more. Deep disc a row ends immediately to prevent birds from feeding on granules.


Crop & Time

Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
COTTON				
At-planting	Aphids, Thrips	2 to 3.5	2.5 to 4.5	Drill granules just below seed place in seed furrow and cover. If seeds and TEMIK are hill-planted, rates may be reduced by one-half.
	Fleahoppers, Leafminers, Mites, Overwintering boll weevil (adults feeding on foliage), Plant bugs, including Lygus	4 to 7	5 to 8.5	
	Nematodes	3.5 to 10 (Except far West)	4.5 to 12	Drill granules just below seed place in seed furrow and cover. If rate exceeds 7 pounds per acre, drill granules in a 4 to 6-inch band into the soil or cover with soil in or above treated zone.
	Nematodes	14 to 27 (Far West Only)	17 to 33	Apply granules in a 4 to 6-inch band below seed line or cover with 2 inches of soil. Plant seed zone.
At first squaring	Leafhoppers, Fleahoppers, Mites, Boll weevil, Plant bugs, including Lygus	7 to 14 14 to 20	8.5 to 17 17 to 24.5	Side-dress granules 8 to 16 inches side of the plant row and 2 inches deep (usually at or below bottom of furrow).
From squaring through early bloom	Cotton leaf perforator, Whiteflies	14 14 to 20	17 17 to 24.5	
GRAIN SORGHUM				
	Nematodes	36 - inch row spacing		
		7	7.5	
		3.5 to 7	4 to 7.5	
At Planting	Greenbug And Chinch Bug (early season control)	[Texas and Oklahoma only]		Apply granules in seed furrow with soil.
		7	7.5	




Crop & Time

Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
POTATOES				
34 - inch row spacing				
<p>DO NOT APPLY IN SUFFOLK COUNTY LONG ISLAND, NEW YORK</p> <p>In the following areas, apply at emergence only:</p>				
States of:	Aphids	7	(Maine Only) 7	<p>Apply granules in 4-6 inch band over row on top of hill. Cover granules with soil by cultivators or hilling discs. DO NOT APPLY THIS PRODUCT TO ANY FIELD MORE THAN ONCE EVERY TWO YEARS.</p>
Maine and Wisconsin				
Counties of:	Colorado potato beetle	14	14.5	
Hartford, CT	Leafhoppers			
Kent and New Castle, DE	Aphids (except in Maine)			
Franklin and Hampshire, MA				
Worcester, MD				
Atlantic, Burlington, Cumberland, Monmouth and Salem, NJ				
Newport and Washington, RI				
Accomack and Northampton, VA				
In all other areas:	Flea beetles	20	21	<p>Apply granules with seed pieces in planting furrow and cover with soil.</p> <p>Apply as above, OR drill granules 2 to 4 inches on both sides of seed row and 3 to 8 inches deep (usually 1 to 2 inches below seed pieces.)</p> <p>Apply as above.</p> <p>Apply granules with seed pieces in planting furrow and cover with soil, OR apply granules in an 8-inch band and work into the soil or cover with soil to a depth of 4 inches. Plant seed pieces in treated zone.</p>
<u>At Planting</u>	Aphids, Colorado potato beetle, leafhoppers	14 to 20	14.5 to 21	
	Psyllids	14 (Arizona Only)	14.5	
	Nematodes (Except Golden Nematode Root-knot nematodes are suppressed. Verticillium wilt has been suppressed where nematode populations have been reduced.)	20	21	
<u>After Planting</u>				<p>Side-dress granules 4 to 5 inches on both sides of plant row and 4 inches deep where pest populations are expected 2-3 weeks after plant emergence. Do not apply if at planting treatment was made.</p>
At 75% emergence or before plants are 6 inches high.	Aphids, Colorado potato beetle	14 to 20	14.5 to 21	





0266G/PENDING/03-14-85



Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 feet of Row	Recommended Application
PEANUTS				
35 - inch row spacing				
At planting	Thrips	7 to 14	7.5 to 15	Apply granules in seed furrow and cover with soil. In Southwest use high rate only.
	Nematodes	14 to 20	15 to 22	Apply granules in a 6 to 12-inch band and work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in treated zone.
SUGAR BEETS				
22 - inch row spacing				
At-planting or within one week before planting.	Nematodes	27 to 33	18 to 22	Apply granules in a 4 to 6-inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in or above treated zone, OR drill granules 1.5 to 3 inches to one side of seed row and 2 to 4 inches deep.
At-planting	Aphids	7 to 14	4.5 to 9.5	Drill granules 1 to 3 inches below seed line. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre. Apply granules in a 2 to 4-inch band over seed row and immediately work into the soil or cover with soil, OR where furrow irrigation is employed for seed germination, drill granules 2 inches deep and 2 inches from seed row on water furrow side. for the 7 lb. rate, apply granules in a 1 to 2 inch band in front of the press wheel as the furrow is closing.
	leafminers, Leafhoppers	14 to 20	9.5 to 13.5	
	Sugar beet root maggot	7 to 14	6.5 to 9.5	
At-planting plus postemergence (split application)	Nematodes	14 at-planting and 14 postemergence	9.5 at-planting and 9.5 post-emergence	AT PLANTING: Apply granules in a 4 to 6-inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in or above treated zone, OR drill granules 1.5 to 3 inches to one side of seed row and 2 to 4 inches deep. POSTEMERGENCE: Side-dress granules 4 to 8 inches to one side of plant row and 3 to 6 inches deep (furrow depth). Apply 40 to 60 days after planting.
Postemergence (Do not make any postemergence application if 27 to 33 lbs./acre were used for nematode control at planting or 1 week before planting).	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil.
	Aphids Leafminers, Leafhoppers	7 to 14 14 to 20	4.5 to 9.5 9.5 to 13.5	Side-dress granules 4 to 8 inches to one side of plant row and 2 to 6 inches deep. A repeat application may be required for continued protection against virus vectors (aphids, leafhoppers).
	Nematodes	27	18	Side-dress granules 4 to 8 inches to one side of plant row and 3 to 6 inches deep (furrow depth). Apply 40 to 60 days after planting.


Crop & Time	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
SOYBEANS				
At-planting	Mexican bean beetle	30 to 48 inch row spacing		Drill granules 2 to 3 inches below seed line, OR 2 to 3 inches to the side of the seed row and 2 to 3 inches deep. Granules can be applied in seed furrow if rate does not exceed 5 pounds per acre.
	Thrips Suppression of three cornered alfalfa hopper	5 to 10 (Based on 36 inch row spacing)	5.5 to 11	
	Nematodes	10 to 20 (Based on 36 inch row spacing)	11 to 22	Apply granules in a 6 to 8 inch band and immediately work into the soil or cover with soil. Plant seed in treated zone.
BEANS				
(Dried only)	Seedcorn maggot	22 to 48 inch row spacing		Apply granules in seed furrow and cover with soil.
At-Planting		3.5 to 5 (Based on 36 inch row spacing)	4.0 to 5.5 (New York Only)	
	Aphids	5 to 7 (Based on 36 inch row spacing)	5.5 to 7.5	Drill granules 2 to 3 inches below seed line OR, 2 to 3 inches to side of seed row and 2 to 3 inches deep,
	Leafhoppers, Mexican bean beetle, Mites	7 to 14 (Based on 36 inch row spacing)	7.5 to 15	OR Granules can be placed in seed furrow if rate does not exceed 5 pounds per acre.
	Nematodes	7 to 14 (Based on 36 inch row spacing)	7.5 to 15	Apply granules in a 6 to 8 inch band and immediately work into the soil or cover with soil. Plant seed in treated zone.
SWEET POTATOES				
At-planting	Nematodes	10 - 20 (Based on 48 inch row spacing)	15 to 30	Apply granules in a 12-inch band in opened row. Cover immediately with soil by hilling up 8 to 10 inches. Plant in center of treated zone.



7/1/85

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 feet of Row	Recommended Application
SUGAR CANE				
At-planting	Nematodes	60 - inch row spacing 14 to 20	26 to 37 (Louisiana Only)	Apply granules in open row on top of newly planted cane and cover immediately with at least 6 inches of soil.
PECANS (Southeast Only) Producing trees				
During period from bud break to nut set, or about April 15 to May 15	Aphids, Mites	33 - 67	Not Applicable	Apply as a 4-6 foot band along dripline on both sides of tree row by spreading granules uniformly and immediately working into soil, OR by shanking 2-3 inches into soil on 12 inch centers.
One to two weeks before bud break.	Suppression of pecan leaf phylloxera			
<u>Newly transplanted trees (1-5 years old)</u> During period of bud break, or about April 15 to May 15	Aphids Mites	Dosage Ounces TEMIK 15G/tree		
Or to two weeks before bud break.	Bud moth, Suppression of pecan leaf phylloxera	5 - 20	Not Applicable	Apply as a side-dress to individual trees by spreading the granules uniformly around the tree and immediately working into the soil to a depth of 2-3 inches.

SEE PREHARVEST, GRAZING, AND ENVIRONMENTAL LIMITATIONS



2/1

statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER OR NOT SUCH LIABILITY IS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

