

Yellow Copy  
PM 19  
264-330

JUL 9 1991

pg. 1 of 29

Mr. Warren A. Davis  
Rhone-Poulenc Ag Company  
2 T.W. Alexander Drive  
P.O. Box 12014  
Research Triangle Park, NC 27709

Dear Mr. Davis:

Subject: Revised Confidential Statements of Formula  
Temik 15G Aldicarb Pesticide  
EPA Reg. No. 264-330  
Your submission of 11/26/90

We have completed review of the revised confidential statements of formula (CSF) and our comments are listed below:

1. The explanation about incorporation of fines into the finished product is acceptable.

2. Use of nominal concentration is an acceptable label claim, per PR Notice 91-2.

3. Inerts listed in revised formulations "A" (basic, all grits formulation) and "B" (alternate, gypsum formulation) are cleared under 40 CFR 180.1001, para (c) or para (d) and are therefore acceptable.

4. Confidential statements of formula A and B are acceptable and this information has been added to our files for this product. It is our understanding that these two are the only formulations that are actively being sold, and that you wish to retain these and drop all other formulations currently on our files.

Sincerely yours,

Dennis H. Edwards, Jr.  
Product Manager (19)  
Insecticide Rodenticide Branch  
Registration Division (H7505C)

**BEST AVAILABLE COPY**

2079

# RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

## TEMIK® brand 15G ALDICARB PESTICIDE

For Control of Certain Insects, Mites, and Nematodes

ACTIVE INGREDIENT: Aldicarb [2-methyl-2-(methylthio)propionaldehyde O-(methylcarbamoyl)oxime].....15%

INERT INGREDIENTS: .....85%

EPA Reg. No. 264-330

EPA Est. No. 264-GA-01



**KEEP OUT OF REACH OF CHILDREN**  
**DANGER POISON**  
**PELIGRO**



PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

TRANSLATION TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.

For **PRODUCT USE** Information Call 1-800-334-9745

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

### STATEMENT OF PRACTICAL TREATMENT

#### GENERAL

Contact a physician immediately in all cases of suspected poisoning. Illness may be produced rapidly following overexposure to TEMIK® Aldicarb. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN**. If poisoning is suspected in animals, contact a veterinarian.

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything orally to an unconscious or convulsing person. Call a physician and follow General advice listed above.

**IF IN EYES:** Flush eyes with plenty of water and get medical attention.

**IF ON SKIN:** Wash thoroughly with soap and water. Remove contaminated clothing and wash before reuse.

**IF INHALED:** Call a physician and follow General advice listed above.

### PRECAUTIONARY STATEMENTS

#### DANGER

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**FATAL IF SWALLOWED.** Causes cholinesterase inhibition. May be fatal or harmful by skin or eye contact or by breathing dust. Rapidly absorbed through skin or eyes. Do not get on skin or in eyes. Do not breathe dust. Keep away from domestic animals.

Wear long-sleeved clothing, full-length trousers, head coverings, and protective gloves when handling. Always stand up-wind from hopper when loading. Wash hands and face before eating, drinking, or using tobacco. Bathe at the end of work day, washing entire body and hair with soap and water. Change clothing daily. Wash contaminated clothing in strong washing soda solution and rinse thoroughly before reusing.

#### SIGNS AND SYMPTOMS OF OVEREXPOSURE

Abdominal cramps

Blurred vision

Diarrhea

Difficult breathing

Excessive sweating

Headache

Muscle tremor

Nausea

Pinpoint eye pupils

Salivation

Vomiting

Watery eyes

Weakness

In severe cases, convulsions, unconsciousness, and respiratory failure may occur.

#### ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN EPA Letter 11-1-77

ACCEPTED  
with COMMENTS

JUL 9 1991

**BEST AVAILABLE COPY**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

264-330

## GENERAL CAUTIONS

Use TEMIK® brand 15G Aldicarb Pesticide only in accordance with label directions, warnings, and cautions. **DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING AS ANY RESIDUES REMAINING MAY BE ILLEGAL OR HARMFUL. DO NOT STORE OR USE IN OR AROUND THE HOME OR HOME GARDEN.** Use high rate on heavy organic or clay soils. Do not exceed the maximum label rate. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crop, meat, and milk. Treatments in excess of 7 pounds per acre made directly in the seed furrow of cotton or sugar beets, or treatments in excess of 5 pounds per acre made directly to the seed furrow of soybeans or dry beans may delay plant emergence and reduce plant stand.

Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, TEMIK® brand 15G Aldicarb Pesticide and water must be on the same side of the plant row.

Calibrate and adjust application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. For any left over material, see instructions for STORAGE AND DISPOSAL in this booklet.

Deep disc any spills at row ends immediately to prevent birds from feeding on exposed granules.

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination.

Do not apply in Del Norte or Humboldt counties in California; Nassau or Suffolk counties in New York; or Curry County, Oregon.

## COMPATIBILITY

Pesticidal activity of TEMIK® brand 15G Aldicarb Pesticide is not affected by normal applications of fertilizers or other pesticides. Its effectiveness may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix TEMIK® aldicarb with other materials before application.

## GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated area without protective footwear. Persons coming in direct contact with wet treated soil after the initial irrigation or rainfall following treatment shall wear coveralls, rubber or neoprene boots, and gloves. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application. Regular long-sleeved shirt, full-length trousers, and head covering should be worn when working in treated fields. See Precautionary Statement, Statement of Practical Treatment, and Note To Physician for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (DANGER-POISON), area treated with TEMIK® brand 15G Aldicarb Pesticide, date of application, appropriate protective clothing, and re-entry requirements.

**BEST AVAILABLE COPY**

# **PRE-HARVEST AND GRAZING LIMITATIONS**

TO AVOID EXCESSIVE RESIDUES IN OR ON:

## **COTTONSEED**

- Do not make more than one 'at planting' application and one postemergence application per crop.
- Do not apply within 90 days of harvest.
- Do not feed cotton forage to livestock or allow livestock to graze in treated area.

## **POTATOES**

- Do not make more than one application per crop.
- Do not harvest within 90 days of an 'at planting' application or within 50 days of an 'after planting' application.

## **PEANUTS**

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not hog-off treated fields.
- Do not feed hay or vines to livestock.

## **SUGAR BEETS**

- Do not make more than one 'at planting' application and two postemergence applications per crop.
- Do not exceed a total of 40 pounds per acre.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.

## **SOYBEANS**

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay, or straw to livestock.

## **DRY BEANS**

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay or straw to livestock.
- Do not use green pods as food for humans.

## **SUGARCANE AND SWEET POTATOES**

- Do not make more than one application per crop.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.

## **PECANS**

- Do not make more than one application per year.
- Do not allow livestock to graze in treated areas.
- Do not harvest foliage or hay from treated areas.

## **SORGHUM**

- Do not make more than one application per year.
- Do not harvest within 90 days of application.
- Do not feed green forage to livestock.

## **OTHER CROPS**

- Do not plant any crop not listed on this label in soil treated with TEMIK® brand 15G Aldicarb Pesticide within 12 months after last application.

## **MEAT AND MILK**

- Do not allow livestock to graze in treated areas before harvest.

## **GROUNDWATER**

- Observe Environmental Hazard precautions regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- Do not wash, load, or empty application equipment near any well, as this practice is a potential source of groundwater contamination.
- Do not apply this product in Del Norte, and Humboldt counties in California; Nassau or Suffolk counties in New York; or in Curry County, Oregon.
- Do not apply more than 33 lbs/A in the state of Florida.
- Do not apply TEMIK® brand 15G Aldicarb Pesticide (EPA Reg. No. 264-330) to citrus. For citrus apply TEMIK® brand 15G Aldicarb Pesticide for Citrus (EPA Reg. No. 264-417).
- Do not apply to any field more than once every two years in Maine and Wisconsin.

**BEST AVAILABLE COPY**

## DIRECTIONS FOR USE AND RECOMMENDED APPLICATIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read entire label before using this product.

To provide maximum performance and to minimize hazard to birds, work granules into or cover with soil to a depth of 2 inches or more. Deep disc any spills at row ends immediately to prevent birds from feeding on granules. When a range or rates are recommended, use the higher rate if pest infestations are expected to be severe.

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>COTTON</b>				
			<u>40" row spacing</u>	
At planting	Aphids, Thrips	3.5 to 5	4.5 to 6	Drill granules just below seed line or place in seed furrow and cover with soil.
		(Except Texas Oklahoma and New Mexico)		
		2 to 5	2.5 to 6	If seeds and TEMIK® are hill-dropped, TEMIK® rates may be reduced by one-half
		(Texas, Oklahoma, and New Mexico Only)		
	Fleahoppers, Leafminers, Mites, Overwintering boll weevil (adults feeding on foliage), Plant bugs including Lygus	4 to 7	5 to 8.5	
	Nematodes	3.5 to 10 (Except Far West)	4.5 to 12	Drill granules just below seed line or place in seed furrow and cover with soil. If rate exceeds 7 pounds per acre apply granules in a 4 to 6-inch band and work into the soil or cover with soil; plant seed in or above treated zone.
	Nematodes	14 to 27 (Far West Only)	17 to 33	Apply granules in a 4 to 6-inch band 3 to 6 inches below seed line or cover with 3 to 6 inches of soil. Plant seed above treated zone.
	Leafhoppers, Fleahoppers	7 to 14	8.5 to 17	
At first squaring	Mites, Boll weevil, Plant bugs, including Lygus	14 to 20	17 to 24.5	Side-dress granules 8 to 16 inches to one side of the plant row and 2 to 6 inches deep (usually at or below bottom of water furrow).
From squaring through early bloom	Cotton leaf perforator	14	17	
	Whiteflies	14 to 20	17 to 24.5	
<b>GRAIN SORGHUM</b>				
			<u>36" row spacing</u>	
At Planting	Nematodes	7	7.5	
		3.5 to 7 (Texas and Oklahoma only)	4 to 7.5	Apply granules in seed furrow and cover with soil.
	Greenbug And Chinch Bug (early season control)	7	7.5	

BEST AVAILABLE COPY

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>POTATOES</b>  IN THE FOLLOWING STATES, APPLY AT EMERGENCE ONLY: Connecticut Delaware Maine Maryland Massachusetts New Hampshire New Jersey Pennsylvania Vermont Virginia Wisconsin  IN THE FOLLOWING STATES/COUNTIES, DO NOT APPLY THIS PRODUCT TO ANY FIELD MORE THAN ONCE EVERY TWO YEARS: Maine Wisconsin Hartford Co., CT Kent and New Castle Co., DE Franklin and Hampshire Co., MA Worcester Co., MD Atlantic, Burlington, Cumberland, Monmouth and Salem Co., NJ Accomack and Northampton Co., VA  DO NOT APPLY IN SUFFOLK OR NASSAU COUNTIES IN NEW YORK.	Aphids (Maine only)	7	<u>34" row spacing</u>  7	Apply granules in 4 to 6-inch band over row on top of hill. Cover granules with soil by cultivators or hilling discs. Mechanical injury may occur during incorporation following applications to plants more than three inches tall. There must be a 50-day interval between application and harvest.
	Colorado potato beetle Leafhoppers  Aphids (except in Maine)	14	14.5	Apply as above.

**BEST AVAILABLE COPY**

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>POTATOES (continued)</b>  <b>IN ALL OTHER AREAS</b>  <b>At Planting</b>  Do not exceed a total of 20 lbs. per acre.	Flea Beetles	20	21	Apply granules with seed pieces in planting furrow and cover with soil.
	Aphids Colorado potato beetle Leafhoppers	14 to 20	14.5 to 21	Apply as above, OR for furrow irrigation drill granules 2 to 4 inches on both sides of seed row and 1 to 2 inches below seed piece depth.
	Psyllids	(Arizona Only)		Apply as above.
	Nematodes (Except Golden Nematode and Root-knot nematodes). Verticillium wilt has been suppressed where nematode populations have been controlled. Root-knot nematodes are suppressed. Despite suppression of Root-knot nematode, tuber quality may not be adequately protected. If Root-knot nematode is a severe economic pest, the use of other control measures should also be employed.	14 20	14.5 21	Apply granules with seed pieces in planting furrow and cover with soil, OR apply granules in a 6-inch band and work into the soil or cover with soil to a depth of 4 inches. Plant seed pieces in treated zone.
	Aphids Colorado potato beetle	14 to 20	14.5 to 21	Apply granules in a 4 to 6 inch band over row on top of hill. Cover granules with soil by cultivator or hilling discs. Mechanical injury may occur during incorporation following applications to plants more than three inches tall. Where furrow irrigation is used, side-dress granules 5 to 8 inches on both sides of plant row and 4 inches deep where pest populations are expected 2 to 3 weeks after plant emergence. Do not apply if at- planting treatment was made. There must be a 50-day interval between application and harvest.
<b>After Planting</b>  <b>At Emergence</b>  Do not exceed a total of 20 lbs. per acre.				

**BEST AVAILABLE COPY**

8729

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>PEANUTS</b>				
			<u>36" row spacing</u>	
At planting	Thrips	7 to 14	7.5 to 15	Apply granules in seed furrow and cover with soil. In Southwest use high rate only.
Do not exceed a total of 20 lbs. per acre.	Nematodes	14 to 20	15 to 22	Apply granules in a 6 to 12 inch band and work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in treated zone.
<b>SUGAR BEETS</b>				
			<u>22" row spacing</u>	
At planting or within one week before planting.	Nematodes	27 to 33	18 to 22	Apply granules in a 4 to 6 inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in or above treated zone, OR where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
At planting	Aphids Leafminers Leafhoppers	7 to 14 14 to 20	4.5 to 9.5 9.5 to 13.5	Drill granules 1 to 3 inches below seed line. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre.  Repeat applications will be required for continued protection against virus vectors (aphid and leafhoppers).
	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules in a 2 to 3 inch band over seed row and immediately work into the soil or cover with soil, OR where furrow irrigation is employed for seed germination, drill granules 2 inches deep and 2 inches from seed row on water furrow side.  For the 7 lb. rate, apply granules in a 1 to 2 inch band in front of the press wheel as the furrow is closing.
At planting plus postemergence (split applications)	Nematodes	14 to 20 at planting and 14 to 20 postemergence	9.5 to 13.5 at planting and 9.5 to 13.5 postemergence	AT PLANTING: Apply granules in a 4 to 6-inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in or above treated zone, OR where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side. POSTEMERGENCE: Apply granules to both sides of plant row and immediately work into the soil or cover with soil, or for furrow irrigation side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.

BEST AVAILABLE COPY



Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SUGAR BEETS (cont.)</b> Postemergence	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil, or for furrow irrigation, side-dress granules 4 to 8 inches to water furrow side of plant row and at furrow depth. Irrigate soon after application.
Do not make any post- emergence application if 27 to 33 lbs/acre were applied at planting or 1 week before planting. Do not make more than 1 'at-planting' application and 2 postemergence applications per crop.	Achids	7 to 14	4.5 to 9.5	Apply as above. A repeat application may be required for continued protection against virus vectors (aphid, leafhoppers).
	Leafminers, Leafhoppers	14 to 20	9.5 to 13.5	
	Nematodes	27	18	Apply as above. Apply within 60 days after planting.

**DO NOT EXCEED A TOTAL OF 40 LBS. PER ACRE**

<b>SOYBEANS</b> At planting	Mexican bean beetle Thrips Suppression of three cornered alfalfa hopper	30" to 48" row spacing		Drill granules 2 to 3 inches below seed line, OR 2 to 3 inches to the side of the seed row and 2 to 3 inches deep. Granules can be applied in seed furrow if rate does not exceed 5 pounds per acre.
		5 to 10	5.5 to 11	
	Nematodes	10 to 20	11 to 22	Apply granules in a 6 to 8-inch band and immediately work into the soil or cover with soil. Plant seed in treated zone.

<b>BEANS</b> (Dried only)	Seedcorn maggot		<u>22" to 48" row spacing</u>	Apply granules in seed furrow and cover with soil.
At Planting		3.5 to 5	4.0 to 5.5	
(Michigan and New York Only)				
	Aphids	5 to 7	5.5 to 7.5	Drill granules 2 to 3 inches below seed line OR, 2 to 3 inches to side of seed row and 2 to 3 inches deep. OR Granules can be placed in seed furrow if the rate does not exceed 5 pounds per acre.
	Leafhoppers Mexican bean beetle Mites	7 to 14		
Do not exceed a total of 14 lbs. per acre.	Nematodes	7 to 14	7.5 to 15	Apply granules in a 6-inch band and im- mediately work into the soil or cover with soil. Plant seed in treated zone. Where furrow irrigation is employed soon after planting, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.

**BEST AVAILABLE COPY**

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
<b>SWEET POTATOES</b> At planting	Nematodes	10 to 20	<u>48" row spacing</u> 15 to 30	Apply granules in a 12-inch band in opened row. Cover immediately with soil by hilling up 8 to 10 inches. Plant in center of treated zone.
<b>SUGARCANE</b> At planting	Nematodes	14 to 20 (Louisiana Only)	<u>60" row spacing</u> 26 to 37	Apply granules in open row on top of newly planted cane and cover immediately with at least 6 inches of soil.
<b>PECANS</b> <u>Producing trees</u> During period from bud break to nut set,	Aphids	33 to 47 (Arizona Only)	Not Applicable	Apply as a 4 to 6-foot band along dripline on both sides of tree row by spreading granules uniformly and immediately working into soil, OR by shanking 2-3 inches into soil on 12 inch centers.
	Aphids, Mites	33 to 67 (except Florida)		
	Suppression of pecan leaf phylloxera	33 (Florida only)		
One to two weeks before bud break.				In Florida, apply between January 1 and April 30 only.
<b>Newly transplanted trees</b> (1-5 years old) During period of bud break, (except Florida)	Aphids Mites	Dosage Ounces <u>TEMIK® 15G/Tree</u>	Not Applicable	Apply as a side-dress to individual trees by spreading the granules uniformly around the tree and immediately working into the soil to a depth of 2-3 inches.
		5 to 20		
One to two weeks before bud break.	Bud Moth Suppression of pecan leaf phylloxera	5 to 20	Not Applicable	In Florida, do not exceed 20 lbs/A and apply between January 1 and April 30 only.

**SEE PREHARVEST AND GRAZING LIMITATIONS**

**PLANT GROWTH STIMULATION:** In the absence of recognizable target pests and under certain growing conditions, stimulation of plant growth by TEMIK® aldicarb on certain crops has been demonstrated under laboratory, greenhouse, and field conditions. Plants may be visibly taller, greener, and denser. Faster grow-off and an increase in fruiting rate and size often are results of the growth enhancement by TEMIK® brand aldicarb. However, such effects do not occur with all crops or under all conditions. Therefore, this product should not be used solely as a plant growth regulator, and its effect as one, should be considered a side benefit which may or may not occur as a consequence of its use.

**BEST AVAILABLE COPY**

# TEMIK® BRAND 15G (GYPSUM) ALDICARB PESTICIDE CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS\* FOR TEMIK® BRAND 15G ALDICARB PESTICIDE (GYPSUM) AT 4, 5 & 6 MPH

NOTE. This calibration chart is applicable only to TEMIK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a gypsum carrier. All settings are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Guide provided with tube). Calibration tubes are available from your TEMIK® brand 15G Aldicarb supplier.

POUNDS TEMIK® 15G PER ACRE FOR VARIOUS ROW SPACINGS					TYPE OF GRANULAR APPLICATOR											
					GANDY GAUGE SETTING			JOHN DEERE (7100) SETTING			NOBLE E-Z FLOW GAUGE SETTING			CLAMP CO 3/8" FEEDER DIAL RPM		
					MPH			MPH			MPH			MPH		
ROW SPACING	22"	34"	36"	40"	48"	4	5	6	4	5	6	4	5	6	4	5
1.8	1.2	1.1	1	.8	7	8	9	3**	3**	4**	2	2	3	3		
3.6	2.4	2.2	2	1.7	10	11	12	4**	5**	6**	3	4	5	6		
5.5	3.5	3.3	3	2.5	12	13	14	6**	7**	8	5	6	7	8		
7.3	4.7	4.4	4	3.3	13	15	16	7**	9	10	6	7	8	11		
9.1	5.9	5.5	5	4.2	14	16	18	9	11	13	7	9	10	14		
10.9	7.1	6.7	6	5.0	16	18	19	11	13	15	8	10	12	16		
12.7	8.2	7.8	7	5.8	17	19	21	12	15	18	9	11	13	18		
14.5	9.4	8.9	8	6.7	18	20	22	14	17	20	10	12	14	21		
16.4	10.6	10.0	9	7.5	19	21	23	16	19	23	11	13	15	23		
18.2	11.8	11.1	10	8.3	20	22	24	17	21	25	12	15	16	26		
21.8	14.1	13.3	12	10.0	22	24	26	20	25	28	14	16	19	31		
25.5	16.5	15.6	14	11.7	23	26	29	24	28	30	16	18	21	37		
29.1	18.8	17.8	16	13.3	25	28	31	26	29	32	17	20	23	42		
32.7	21.2	20.0	18	15.0	26	30	33	28	31	33	19	21	24	48		
36.4	23.5	22.2	20	16.7	28	31	34	30	32	35	20	23	26	54		
43.6	28.2	26.7	24	20.0	31	34	37	32	34	37	22	26	29	65		
49.1	31.8	30.0	27	22.5	33	36	40	33	36	38	24	28	31	74		

\*All settings for one hopper box outlet per row

\*\*For John Deere gauge setting below 8 there will be great variations in flow rate. A gauge setting of 6 is the lowest recommended setting.

NOTE: THIS CALIBRATION CHART WILL APPEAR ON THE 30 LB. POUR BOX.

BEST AVAILABLE COPY

# TEMIK® BRAND 15G (CORNCOB) ALDICARB PESTICIDE CALIBRATION GUIDE

## APPROXIMATE APPLICATOR SETTINGS\* FOR TEMIK® BRAND 15G ALDICARB PESTICIDE (CORNCOB) AT 4, 5 & 6 MPH

NOTE: This calibration chart is applicable only to TEMIK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a corncob carrier. All rates are APPROXIMATIONS must be confirmed by using a calibration tube (See Calibration Guide provided with tube). Calibration tubes are available from your TEMIK® 15G Aldicarb supplier.

POUNDS TEMIK® 15G PER ACR. FOR VARIOUS ROW SPACINGS					TYPE OF GRANULAR APPLICATOR									
					GANDY GAUGE SETTING			JOHN DEERE MAX EMERGE GAUGE SETTING**			NOBLE E-Z FLOW GAUGE SETTING			CLAMPCO 1/2" FEEDER DIAL RPM
					MPH			MPH			MPH			MPH
22"	34"	36"	40"	48"	4	5	6	4	5	6	4	5	6	5
1.8	1.2	1.1	1	.8	14	16	17	8	9	10	5	6	7	4
3.6	2.4	2.2	2	1.7	19	21	22	11	13	14	10	11	13	8
5.5	3.5	3.3	3	2.5	22	24	27	14	16	18	13	15	17	11
7.3	4.7	4.4	4	3.3	25	28	30	17	19	21	16	18	21	15
9.1	5.9	5.5	5	4.2	28	31	33	19	21	23	18	22	24	19
10.9	7.1	6.7	6	5.0	30	33	35	21	23	25	21	24	26	22
12.7	8.2	7.8	7	5.8	32	35	37	22	25	26	23	26	29	26
14.5	9.4	8.9	8	6.7	34	37	39	24	26	28	25	28	31	30
16.4	10.6	10.0	9	7.5	35	38	41	25	27	29	26	30	33	33
18.2	11.8	11.1	10	8.3	37	40	43	26	28	30	28	31	34	37
21.8	14.1	13.3	12	10.0	39	43	47	28	30	33	31	34	37	44
25.5	16.5	15.6	14	11.7	42	46	50	29	32	35	33	36	40	52
29.1	18.8	17.8	16	13.3	44	49	54	31	34	37	35	39	44	59
32.7	21.2	20.0	18	15.0	47	52	59	33	36	38	37	42		67
36.4	23.5	22.2	20	16.7	49	56	66	34	37	40	39	45		76
43.6	28.2	26.7	24	20.0	54	66		37	40	44	44			95
49.1	31.8	30.0	27	22.5	59			38	42					

\*All settings for one hopper box outlet per row

\*\*Old Max Emerge Housing

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER OR NOT SUCH LIABILITY IS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

### NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company  
Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

TEMIK is the registered trademark of Rhône-Poulenc for aldicarb pesticide.  
© 1991 Rhône-Poulenc Ag Company

TEMIK brand 15G Aldicarb Pesticide (PENDING) submitted 6/13/91.

BEST AVAILABLE COPY

130629

## **RESTRICTED USE PESTICIDE**

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

# **TEMIK® Brand 15G Aldicarb Pesticide**

EPA Reg. No. 264-330

## **Environmental Precautions and Soil Type Restriction Tables**



Refer to the container label for additional use precautions and directions.



Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T. W. Alexander Drive  
Research Triangle Park, North Carolina

TEMIK is the registered trademark of Rhône-Poulenc for aldicarb pesticide  
© 1991 Rhône-Poulenc Ag Company

**BEST AVAILABLE COPY**

140829

**Contents** **Page**

ENVIRONMENTAL PRECAUTIONS & GENERAL USE RESTRICTIONS ..... 3

**SOIL TYPE TABLES BY STA. \***

ALABAMA.....	4
COLORADO.....	4
CONNECTICUT.....	4
DELAWARE.....	5
GEORGIA.....	5
ILLINOIS.....	5
INDIANA.....	5
IOWA.....	6
KANSAS.....	6
KENTUCKY.....	6
LOUISIANA.....	6
MAINE.....	6
MARYLAND.....	7
MASSACHUSETTS.....	7
MICHIGAN.....	7
MINNESOTA.....	8
MISSISSIPPI.....	8
MISSOURI.....	8
MONTANA.....	8
NEBRASKA.....	9
NEW HAMPSHIRE.....	10
NEW JERSEY.....	10
NORTH CAROLINA.....	10
NORTH DAKOTA.....	11
OHIO.....	11
PENNSYLVANIA.....	11
SOUTH CAROLINA.....	11
SOUTH DAKOTA.....	12
TENNESSE.....	12
VERMONT.....	12
VIRGINIA.....	13
WEST VIRGINIA.....	13
WISCONSIN.....	13
WYOMING.....	14

**BEST AVAILABLE COPY**

## ENVIRONMENTAL PRECAUTIONS GENERAL USE RESTRICTIONS

**AGRICULTURAL CHEMICALS HAVE THE POTENTIAL TO MOVE INTO SHALLOW GROUNDWATER. THE FOLLOWING RESTRICTIONS HAVE BEEN DEVELOPED TO PROTECT DRINKING WATER SUPPLIES.**

**DO NOT APPLY WITHIN 50 FEET OF ANY DRINKING WATER WELL TO AVOID CONTAMINATION FROM SURFACE RUNOFF.  
MORE STRINGENT RESTRICTIONS MAY BE REQUIRED, AS DISCUSSED BELOW.**

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination.

In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.

Contact your state pesticide regulatory authority for further information on state requirements for the use of this product.

Some states, including Florida, Massachusetts, New York, Rhode Island, and Wisconsin, have or may be developing more restrictive regulations regarding the use and application of TEMIK®. Follow all state regulations restricting the use and application of this product, including limitations on applications near drinking water sources. In all cases, the more restrictive requirements must be followed. It is the responsibility of the applicator to document the construction of wells claimed not to be shallow.

### SPECIFIC USE RESTRICTIONS

**FOLLOW THE LISTED "ADDITIONAL RESTRICTIONS" if the following conditions are present:**

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS
AL, CO, DE, GA, KS, KY, LA, MD, MO, MS, NC, SC, TN, VA, WV	Loamy sand or sand surface soils <u>and</u> subsoils <u>with an</u> average organic matter in the upper 12 inches of less than 2% by weight.  See SOIL RESTRICTION TABLES for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within <u>300 feet</u> of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.
CT, IA, IL, IN, MA, MI, MN, MT, ND, NE, NJ, OH, PA, SD, WI, WY	Sandy loam, loamy sand, or sand surface soils, <u>and</u> loamy sand or sand subsoils, <u>with an</u> average organic matter in the upper 12 inches of less than 2% by weight.  See SOIL RESTRICTION TABLES for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet, do not apply within <u>500 feet</u> of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. In the following counties in Wisconsin: Portage, Langlade, Juneau, Marathon, Adams, Waushara, Waupaca, and Wood; the well set back distance is increased to 1000 feet.
ME, NH, VT	Sandy loam, loamy sand <u>or</u> sand surface soils.  See SOIL RESTRICTION TABLES for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within <u>500 feet</u> of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.
CA (Del Norte Co.) CA (Humboldt Co.) NY (Nassau Co.) NY (Suffolk Co.) OR (Curry Co.)	All Soils	Do not apply this product in the specified counties.
FL	All Soils	State regulations require that TEMIK® not be applied within <u>300 feet</u> of any drinking water well.

**SOIL RESTRICTION TABLE**
**FOLLOW THE LISTED ADDITIONAL RESTRICTIONS if the soil types are present in your state:**
**ALABAMA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	bigbee	crevasse	fripp	leon	rumford
alpin	bruno	duckston	gorgas	louisburg	saffell
americus	cartecay	eustis	jones	osier	scranton
bankhead	chipleay	flomaton	kalmia	pactolus	tarboro
bassfield	corolla	foxworth	lakeland	plummer	

**COLORADO**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

acasco	columbine	gilcrest	lonetree	pescar	tassel
adilis	comad	glenberg	luning	peyton	terry
alda	como	gracot	manhattan	platte	thiel
alex	corlett	graypoint	manter	pring	thoroughfare
amay	costilla	gredivid	marosa	quamon	tine
azeltine	cotopaxi	grimstone	maybell	ratake	tipper
bangston	coyet	gunbarrel	maysprings	redrob	tipperary
bankard	crestman	hiwan	medano	relsob	tivoli
barcus	curecanti	hobacker	mespun	riverwash	tolby
baroid	dahlquist	hooper	miracle	ruedloff	tomah
becks	dailey	howardsville	moenkopie	ruhe	trail
bernal	derrick	ildelfonso	montez	ryan park	tricara
biou	dix	inavale	moosed	ryark	truckton
blackston	driggs	ipage	mosca	sabe	trude
blakeland	dunday	ironsprings	mysten	saguache	tullock
blanca	dunul	jugot	naturita	san arcacio	uffens
bobtail	dwyer	julesburg	nederland	santana	uracca
boel	wachuston	kandaly	nelson	satanta	utica
bond	eckley	kassler	nesda	schamber	valent
bottle	ecklund	kerber	newcomb	schooner	valentine
bowdish	ellicott	kettle	newfork	sheppard	vasquez
bresser	els	kippen	newlin	silvercliff	vastine
broad canyon	elsmere	laird	niwot	southace	vernal
cabin	falcon	larim	norriston	space city	willwood
canlodore	farb	las	numa	spool	yetull
cascajo	florita	las animas	orsa	st. elmo	yoder
chappell	fourmile	layoint	osgood	stapleton	zeona
cheyenne	frisco	legault	ouray	stecum	
clark fork	fruitland	leighcan	pando	stoneham	
cliffdown	gas creek	lincoln	patna	stunner	
colby	gerrard	littlebear	peetz	tabernash	

**CONNECTICUT**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply with 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

agawam	canton	farmington	hollis	penwood	suncook
amenia	charlton	fredon	holyoke	pipestone	sutton
belgrade	cheshire	gloucester	manchester	pooratuck	wareham
birchwood	copake	groton	merimac	poquonock	windsor
branford	deerfield	hartford	montauk	saco	yalesville
brimfield	ellington	hero	nellis	scarboro	
brookfield	enfield	hinckley	occum	sudbury	



170829

**SOIL RESTRICTION TABLE (continued)**

**DELAWARE**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

atsion	evesboro	kenansville	osier
collington	kalmia	klej	rumford

**GEORGIA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

americus	centenary	fripp	kureb	molena	scranton
ashlar	chiple	galestown	lakeland	osier	sweetapple
bigbee	duckston	gorgas	lakewood	ousley	
buncombe	echaw	johns	leon	paola	
cainhoy	eustis	kalmia	louisburg	plummer	
cartecay	flomaton	kershaw	lynn haven	pottsborg	
cedarbluff	foxworth	klej	mandarin	ridgeland	

**ILLINOIS**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

ade	coyne	hayrie	lanier	parkville	udolpho
alvin	dakota	hesch	lawler	plainfield	ware
backbone	dickinson	hitt	lomar	ridgeville	warsaw
beardstown	disco	homer	lorenzo	robby	watseka
billet	dowagiac	hononegah	martinsville	rockton	wauke
bloomfield	downs	hoopston	matherton	rodman	waukegan
boone	dresden	iona	maumee	ruark	whalan
bowdre	el dara	kane	minneiska	sarpy	wheeling
boyer	eleva	kankakee	morocco	saude	wirt
burkhardt	flagler	kidder	muskingum	shoals	zumbro
burnside	fox	la hogue	oakville	sparta	
carmi	gale	la rose	ockley	st. charles	
casco	genesee	lacscent	onarga	stockland	
chelsea	gifford	lamont	orio	stonelick	
chute	haymond	landes	pana	trempealeau	

**INDIANA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

absco	carmi	fox	maumee	pipestone	st. charles
ade	casco	genesee	milgrove	plainfield	stockland
algansee	ceresco	gifford	monitor	pope	stonelick
alida	chatterton	gravelton	morocco	princeton	tedrow
alvin	chelsea	hanna	moundhaven	prochaska	tracy
ayrshire	chetwynd	haymond	mudlavia	quinn	tyner
beckville	cohocah	hillsdale	muskingum	riverdale	uniontown
belmore	coloma	homer	nesius	robby	warsaw
billet	conrad	hoopston	newton	rockton	wasepi
bloomfield	coupee	iona	nineveh	rodman	watseka
bobtown	craigmile	junius	oakville	ruark	weikert
bourbon	dickinson	kalamazoo	ockley	saugatuck	wheeling
boyer	door	kosciusko	onarga	seafield	wirt
brady	dowagiac	landes	ormas	shipshe	zaborosky
branch	dresden	linkville	orville	shoals	zadog
brems	ellean	lomar	oshtemo	sleeth	
bronson	elston	lydick	ouiatenon	sloan	
bruno	fabius	lyles	piankeshaw	sparta	
burnside	farmington	martinsville	pinhook	spinks	

**SOIL RESTRICTION TABLE (continued)**

**IOWA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

arkeny	chute	gale	lamont	perks	vore
aureola	coyne	gillford	landes	renova	wadena
backbone	cylinder	grable	lawler	ridgeport	wapsie
bertram	dickinson	hanlon	lilah	rockton	watseka
billet	dickman	hawick	linder	salida	waukegan
bolan	downs	hayfield	marlean	sarpy	whalan
boone	elrick	haynie	monona	saude	winneshiek
brady	estherville	hesch	montie'h	sparta	zenor
buckney	fairhaven	hixton	mottland	sunburg	
burkhardt	finchford	hoopeston	nevin	toolesboro	
carr	flagler	kanaranzi	oesterle	udolpho	
chelsea	fruitfield	kum	percival	ulster	

**KANSAS**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

albion	dix	jansen	meadin	schamber	valent
bankard	dorrance	kanza	munjor	simeon	valentine
boel	dwyer	las	platte	stephenville	
carr	els	las animas	pratt	sweetwater	
darr	elsmere	likes	quinlan	thurman	
dillwyn	inavale	lincoln	sarpy	tivoli	

**KENTUCKY**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

beulah	crevasse	lakin	potomac	saffell	
bruce	flomaton	molena	riverwash	steinsburg	

**LOUISIANA**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	darden	flo	kalmia	lotus	peveto
betis	eustis	hackberry	lakeland	osier	
crevasse	felicity	haggerty	latonia	palm beach	

**MAINE**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

abram	colton	fryeburg	medomak	potsdam	sunday
adams	cornish	groveton	mekrose	red hook	surplus
allagash	crary	halsey	monadnock	ricker	swanton
au gres	creasey	hermon	monarda	riverwash	telos
bangor	croghan	howland	moosilauke	rumney	thorndike
becket	danforth	kinsman	naskeag	saddleback	togus
berkshire	dixfield	lovewell	naumburg	schoodic	tunbridge
brayton	duane	lyman	nicholville	searsport	waumbek
canaan	eldridge	lym-	ondawa	sheepscot	westbury
caribou	elliottsville	m.	ossipee	shirley	winnecook
cathro	elmwood	ma	peacham	sisk	
charles	enchanted	mapleton	peru	skerry	
chesuncook	finch	marlow	plaisted	skowhegan	
colonel	fredon	masardis	podunk	stetson	

# SOIL RESTRICTION TABLE (continued)

## MARYLAND

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

atsion	downer	fort mott	lakeland	osier	st. johns
collington	dragston	galestown	leetonia	remlik	steinsburg
colts neck	evesboro	k'lej	leon	rumford	swansea

## MASSACHUSETTS

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

adams	canton	essex	hooksan	pipestone	shelburne
agawam	carver	evesboro	hoosic	pittsfield	sudbury
amenia	charlton	farmington	k'lej	plymouth	taconic
au gres	chattfield	fredon	lyman	pompton	unadilla
barnstable	cheshire	fullam	merrimac	pootatuck	walpole
belgrade	copake	gloucester	millis	poquonock	whitman
berkshire	deerfield	groton	montauk	quonset	
birchwood	dukes	hero	nellis	riverhead	
brimfield	eastchop	hinckley	norwell	saco	
broadbrook	eldridge	hollis	oakville	saugatuck	
brookfield	enfield	holyoke	occum	scituate	

## MICHIGAN

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abbaye	crosswell	guardlake	manitowish	oshtemo	sleeth
abscota	cunard	hagensville	martinsville	otisco	sloan
alcona	deer park	halfaday	matherton	ottokee	solona
algansee	deerton	hessel	maumee	padus	sparta
altonaz	dighton	hettinger	mcbride	pelkie	spinks
alpena	dixboro	hillsdale	mecosta	perene	springlake
arkona	dowagiac	hodenpyl	melita	pence	stauben
arkport	dresden	lonia	michigamme	pendleton	sturgeon
arnheim	dryburg	ishpeming	mitiwanga	pequaming	summerville
au gres	dryden	jesso	monico	perrin	sunfield
banat	duel	johnswood	monitor	peshekee	teasdale
beechwood	dunbridge	kalamazoo	montcalm	petticoat	tedrow
blue lake	east lake	kalkaska	moquah	pipestone	tekenink
borski	eastport	karlin	morocco	plainfield	thetford
boyer	eleva	kawkawlin	munising	ponozzo	tula
brady	elndale	kendallville	nadeau	posen	tyre
branch	elston	kibbie	nahma	remus	velvet
brems	emmet	kidder	newaygo	richter	vilas
bronson	ensign	kingsville	newton	riggsville	wainola
ceresco	ensley	kiva	nottawa	riverdale	waiska
champion	escanaba	klacking	nunica	rodman	wasepi
channing	evart	koontz	oakville	rousseau	whalan
chatham	fabius	landes	ockley	rubicon	winneshiek
cheboygan	fairport	lapeer	oconto	sanilac	winterfield
chelsea	finch	leelanau	okee	saugatuck	woodbeck
cohoctah	fox	leoni	omega	sayner	yalmer
coloma	genesee	liminga	omena	scalley	zimmerman
colonville	gilchrist	locke	onaway	schoolcraft	
coral	gifford	lode	onota	sheldrake	
coupee	gladwin	longrie	ontonagon	shoals	
covert	graycalm	mancelona	ormas	skanee	

BEST AVAILABLE COPY

# SOIL RESTRICTION TABLE (continued)

## MINNESOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abscoia	cromwell	goodland	langola	nowen	soderville
algonsee	cylinder	gotham	lasa	nymore	sparta
alvin	dakota	grayalm	lawler	omega	spinks
andrusia	demontreville	growton	lengby	osakis	sugarbush
ankeny	dickinson	halder	liah	oylen	sunburg
anoka	dickman	hamar	linder	pierz	sverdrup
arvilla	dorerton	hanlon	lino	plainfield	udolpho
automba	dorset	hawick	litchfield	pomroy	ulen
backbone	dowagiac	hecla	lohnes	poppleton	venlo
beavercreek	duelm	heimdal	lomax	prebish	verndale
becker	egeland	hermantown	maddock	quetico	wabedo
bellechester	elbaville	heyder	mahtomedi	rasset	wadena
billett	eleva	hiwood	malachy	redby	waderill
blowers	emmett	hixton	marcellon	redeye	waman
boone	estherville	hoopeston	marlean	renova	watseka
brainerd	etter	hubbard	marquette	renshaw	waukee
brodale	evart	huntersville	meehan	ridgeport	waukegan
burkhardt	eyota	insula	menahga	rockton	wawina
burnsville	fairhaven	jewett	meridian	rockwood	whalan
carmi	faunce	kalmerville	minneiska	rosewood	winterfield
chelsea	finchford	kanaranzi	minneopa	rosholt	wykeham
chetek	finland	karlstad	moquah	salida	wykoff
claire	flagler	kasota	mottland	santiago	zimmerman
clontarf	flak	kingsley	nemadji	sartell	zumbro
cloquet	flaming	koronis	newfound	saude	
conic	friendship	la forge	nokay	serden	
copaston	gale	lacscent	norden	sioux	
cowhorn	genesee	lamont	northfield	snellman	

## MISSISSIPPI

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 200 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	bigbee	eustis	osier	saffell
bassfield	bruno	lakeland	plummer	st. lucie
beulah	crevasse	latonia	riverwash	

## MISSOURI

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alvin	carr	crevasse	grable	riverwash	scotco
beulah	clana	diehlstadt	hodge	saffell	sheldrake
canabou	cooter	eustis	malden	sarpy	towosahgy

## MONTANA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alberton	assinniboine	biglake	breece	chinook	dayschool
alice	attewan	bigsheep	briggsdale	circleville	degrand
allentine	azaar	binna	brownlee	clark fork,	desmet
alona	babb	birney	bullock	coldcreek	dinnen
ambrant	badgerton	blackhall	busby	comad	dominic
amesha	bankard	blacklead	cabba	como	dooley
amor	banks	blaine	cabbart	courville	dryadine
apeldorn	baxendale	blanchard	canburn	cowood	dwyer
apron	bayerton	bonfri	castner	cozberg	earcree
ard	beaverton	borah	chamokane	craddock	elkner
armelle	beenom	bowbac	chanta	crago	elva
ashbon	beeskove	brandenburg	charlos	crittenden	ervide
ashio	belain	branharn	cheadle	cushman	ess
ashuelot	bigarm	brantford	cheyenne	dast	ethridge

SOIL RESTRICTION TABLE (continued)

MONTANA (continued)

evano	kalsted	musselshell	reian	stecum	utica
farnuf	kirby	naz	remmit	stryker	valentine
finleypoint	kiwanis	nelson	rencot	subwell	vastine
firth	krause	nemote	renohill	sudworth	vebar
flasher	labre	nesda	rentsac	tally	victor
flathead	lake creek	nevine	ridge	tamaneen	vona
fleak	lamedeer	niarada	ridgelawn	tamely	wabek
forelle	lantern	nihil	riedel	tenex	wages
frenchcreek	laporte	nippt	riverside	terry	waits
galbreth	larim	noonan	rivra	teton	waldbillig
garlet	lehr	norte	rochester	tevis	walstead
gas creek	libeg	nuley	rootel	thess	walters
geohrock	lihen	nunn	round butte	thiel	wemple
glaciercreek	lisk	oceanet	roxal	tigeron	windham
glenberg	token	olney	ryegate	tine	winfall
glendive	lolo	oraid	ryell	tineman	winkler
gunbarrel	lolopeak	oro fino	ryorp	tinsley	winnett
hagga	lone rock	outlet	sapphire	tolex	wishard
hamlet	lowercreek	ovando	sawcreek	tolman	woodgulch
hanly	macar	parchin	scravo	toluca	woodhall
happgood	macfarlane	parshall	seile	tomichi	woodrock
harvey	manhattan	passcreek	selway	travessilla	woodside
haskill	manning	pengilly	seroco	trembles	worland
haugan	mamarth	perma	shadow	trey	wormser
hedstrom	maurice	petty	shawmut	trimad	yegen
hilger	mccaffery	philcher	sheege	tropal	yellowbay
holloway	mccollum	pishkun	shingle	tullock	yetull
hopley	meadowcreek	poin	shook	turner	
hopsonville	mires	pring	shurley	tusler	
kim	mitten	randman	sixbeacon	twilight	
jocko	mocmont	redcreek	spang	upsata	
jugson	moiese	reeder	splitro	usine	

NEBRASKA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alda	calamus	fonner	laird	munjor	schamber
alice	campus	gibbon	las	nihil	selia
almeria	canyon	glenberg	las animas	norrest	shingle
altvan	carr	gospier	lawet	nunn	sibleyville
ascalon	cass	gothenburg	leshara	o'neill	simeon
bankard	chappell	grable	lex	ogilala	talmo
barney	cheyenne	haynie	lisco	olney	tassel
bayard	dailey	hedville	lockton	ord	thurman
bazile	darr	hemingford	loretto	ortello	trelna
blanche	dickinson	hennings	lute	orwet	tuthill
blendon	dix	hersh	mace	otero	valent
boel	doger	holt	manter	paka	valentine
boelus	duda	humbarger	mcgrew	percival	vetal
bolent	dunday	inavale	mckelvie	pivot	wann
boone	dunn	ipage	meadin	platte	wewela
brocksburg	dwyer	jansen	minatare	ronson	whitelake
brunswick	eckley	janude	minnequa	rosebud	wildhorse
busher	els	jayem	monona	rusco	woodly
bushman	elsmere	johnstown	muk	sarpy	

BEST AVAILABLE COPY

### NEW HAMPSHIRE

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

abernaki	deerfield	kearsarge	moosilauke	raynham	sudbury
acton	duane	kinsman	mundal	redstone	sunapee
adams	dutchess	leicester	naumburg	ricker	suncook
agawam	eldridge	lin	newfields	ridgebury	sunday
au gres	elmridge	londonderry	nicholville	rippowam	surplus
bangor	elmwood	lowell	ninigret	rumney	sutton
becket	fryeburg	lyman	occum	saddleback	swanton
belgrade	glebe	lyme	ondawa	saugatuck	thorndike
berkshire	gloucester	machias	ossipee	scarboro	tunbridge
bernardston	glover	madawaska	paxton	scio	unadilla
caesar	grange	marlow	peacham	scituate	walpole
canaan	groveton	masardis	pennichuck	searsport	wareham
canton	hadley	matunuck	peru	shaker	warwick
cardigan	haven	medomak	pillsbury	sheepscot	waumbek
charles	harmon	melrose	pipestone	sisk	whitman
charlton	hinckley	merimac	pittstown	skerry	wilmington
chaffield	hinesburg	metallak	plaisted	squamscott	windsor
cohas	hollis	millis	podunk	stetson	winnecook
colton	hoosic	monadnock	pootatuck	stissing	winooski
croghan	houghtonville	monarda	quonset	stratton	woodbridge
dartmouth	howland	montauk	rawsonville	success	worden

### NEW JERSEY

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

adelphia	ellington	hasbrouck	lakewood	pascack	shrewsbury
atsion	evesboro	hazen	lansdale	phalanx	st. johns
aura	fallsington	hazleton	leon	plummer	steinsburg
bowmansville	fort mott	hero	matapeake	pompton	swartswood
circumference	fredon	hibernia	matawan	popo	tinton
colington	freehold	holmdel	mullica	preakness	tunkhannock
colonie	tripp	holyoake	nassau	riverhead	wassaic
colts neck	galestown	hooksan	nixon	rockaway	woodstown
downer	gladstone	hoosic	otisville	rowland	wooster
dragston	haledon	klej	parker	sassafras	
dunellen	hammont	lakeland	parsippany	shrewsbory	

### NORTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	centenary	tripp	louisburg	pactolus	toccoa
alpin	charleston	galestown	lynn haven	potomac	tomahawk
baymeade	chipley	immokalee	mandarin	pottsburg	valhalla
biltmore	conetoe	johns	molena	rimini	wake
bojac	corolla	kalmia	newhan	rumford	wakulla
buncombe	dragston	kenansville	nimmo	seabrook	wando
cainhoy	duckston	klej	ona	st. johns	wateree
cartecay	echaw	kureb	osier	suncook	
carteret	eustis	lakeland	ousley	tarboro	

## NORTH DAKOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

amor	cabbart	lossum	lehr	parshall	tusler
arvilla	chantz	glendive	lemert	poppleton	ulen
ayimer	chinook	hamar	letcher	reeder	vebar
baahish	claire	hamlet	lhen	renshaw	venlo
banks	clontarf	harly	lohnes	rhone	vetal
bantry	coe	haskill	maddock	ruso	wabek
beisigl	cohagen	hecla	manning	serden	wakum
birford	desart	heimdal	marmarth	seroco	wamduska
bowdle	egeland	hidatsa	metigoshe	sioux	wanagan
brandenburg	ekalaka	inkster	minnewaukan	tally	warsing
breien	falsen	karlsruh	mott	teller	yegen
brisbane	flasher	ladner	noonan	toby	yetull
cabba	fleak	lefor	osakis	trembles	zeoria

## OHIO

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abscota	channahon	fitchville	kings	orrville	skidmore
algansee	chavies	fulton	laidig	oshtemo	sleeth
alvin	chenango	galen	lamson	otisville	sloan
arkport	clymer	gallipolis	landes	peoga	sparta
barkcamp	cohoctah	geeburg	licking	pike	st. clair
belmore	coloma	gessie	lordstown	piopolis	stafford
bogart	colonia	hackers	markland	plattville	stendal
boyer	conneaut	hanover	milford	pope	tawas
brady	defiance	haubstadt	minoia	princeton	tyler
bronson	del rey	holly	muskego	riddles	warners
carlisle	digby	homewood	nicholson	ritman	washtenaw
casco	dubois	ilion	nolin	rockyfork	wea
cavode	elba	jessup	oakville	shinrock	wharton
centerburg	ellsworth	jules	opequon	sisson	

## PENNSYLVANIA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

adams	conotton	hartsells	linden	phelps	swartwood
alton	cookport	hatboro	manlius	pocono	tunkhannock
bernudian	dekaib	haven	manor	pope	vanderlip
bowmansville	edgemont	hazleton	marr	rexford	weikert
chavies	eliber	lakin	orrville	riverhead	wheeling
chenango	fallsington	lansdale	othello	riverwash	woodstown
chili	fredon	seetonia	otisville	rowland	wyalusing
colonia	gleneig	lehew	parker	sassafras	wyoming
comus	glenville	lewisberry	pequea	steinsburg	

## SOUTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	charleston	johns	molena	ridgeland	wakulla
alpin	chipley	kalmia	nansemond	rimini	wando
ashlar	coosaw	kenansville	newhan	riverwash	waterce
barstari	crevasse	kershaw	olanta	rosedhu	witherbee
beaches	echaw	lakeland	osler	scranton	
buncombe	edisto	leon	pectolus	seabrook	
cainhoy	eustis	louisburg	plummer	seewee	
cartecay	foxworth	lynn haven	quart-	tomahawk	
centenary	frupp	meggett	zipsamments	udipsamments	

# **SOUTH DAKOTA**

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

alice	cohen	hamar	mawer	rockoa	tuthill
akvan	cushman	hanly	meadin	ronson	twilight
akwida	dailey	haynie	minatare	rosebud	ulen
amor	delmont	heath	minnequa	sarpy	valent
anselmo	dix	hecla	moccasin	schamber	valentine
arvilla	doger	helmdal	munjor	serden	vebar
ascalon	dogiecreek	hilger	murdo	seroco	venlo
assinniboine	duda	holt	nihil	shingle	vetal
attewan	dunday	hopdraw	noonan	shirttail	wabek
bankard	dwyer	inavale	norrest	sioux	wann
banks	eckley	ipage	nunn	sorum	westover
blackhall	egeland	jansen	o'neil	storia	wewela
blendon	ekalaka	jayem	oake	sverdrup	whitelake
boei	els	kirby	ogala	tally	winetti
bowdle	elsmere	lefor	orton	talmo	woody
brocksburg	enet	lehr	orwet	tanna	woonsocket
bullock	evridge	letcher	parchin	tassel	wortman
butche	fedora	lihen	parshall	teller	yecross
cabba	flasher	lowry	percival	temvik	yegen
cabbart	flek	lute	platte	tetonka	zeona
canyon	glenberg	maddock	reeder	thurman	
cass	glendive	manning	renshaw	travessilla	
chappell	glenross	manter	reva	trembles	
chinook	grable	mathias	rhame	trey	

# **TENNESSE**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

harish	bruno	crevasse	eustis	saffell
bigbee	buncombe	dexter	riverwash	

# **VERMONT**

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

abram	dantorth	groveton	mekosa	quonset	tioga
adamant	deerfield	hadley	merrimac	rawsonville	tisbury
adams	dixfield	hero	middlebury	raynham	trow
agawam	duane	hinckley	missisquoi	ricker	tunbridge
allagash	dummerston	hinesburg	monadnock	rippowam	unadilla
amenia	dutchess	hogback	moosilauke	rumney	vershire
au gres	duxbury	houghtonville	mundal	scarboro	waitsfield
bangor	eldridge	hubbardton	nassau	scio	walkill
bartlett	elliottsville	kars	nellis	searsport	walpole
belgrade	elmridge	kendaia	nicholville	sheepscot	wappinger
berkshire	elmwood	killington	ninigret	sisk	wareham
bomoseen	enchanted	leicester	occum	st. albans	warwick
brayton	enosburg	londonderry	ondawa	statson	weider
buckland	farmington	lovewell	palatine	stissing	westbury
cabot	fredon	lyman	pawling	stockbridge	wilmington
castile	fryeburg	lyme	paxton	stowe	windsor
cathro	tullam	lyons	peacham	stratton	winooski
charles	galoo	machias	peru	sudbury	woodstock
claverack	galway	macomber	pittsfield	sunapee	worden
colonel	georgia	mansfield	pittstown	sunday	
colrain	glebe	markey	podunk	sunny	
colton	glover	marlow	potfret	swanton	
copake	grange	massena	potatuck	taconic	
croghan	groton	medomak	potadam	teel	



## VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

alaga	carteret	duckston	lakehurst	mcclena	seabrook
ashlar	catpoint	fisherman	lakeland	nansemond	steinsburg
assateague	chagrin	fripp	lakun	newhan	sweetapple
biltmore	chipley	galestown	leestonia	nimmo	tarboro
bojac	conetoe	johns	leon	osler	wakulla
buncombe	corolla	kalmia	louisburg	pacokus	wateree
camocca	dragston	kenansville	matneflat	riverwash	
cartecay	drail	klej	millrock	rumford	

## WEST VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.

drail	gauley	lander	potomac
fiveblock	takin	leestonia	sewell

## WISCONSIN

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. In the following counties in Wisconsin: Portage, Langlade, Juneau, Marathon, Adams, Waushara, Waupaca, and Wood; the well set back distance is increased to 1000 feet.

abskota	delton	hubbard	mahtomedi	ockley	shiffer
alban	dickinson	humbird	mancelona	oconto	soderville
alcona	dickman	impact	manitowish	oesterle	solona
alganssee	dornton	ionia	mann	okee	sparta
alpena	downs	ishpeming	marathon	omega	spinks
amery	dresden	jewett	marcellon	omena	st. charles
ankeny	duel	kalamazoo	markesan	onaway	sturgeon
arland	duelm	kane	matherton	oshtemo	summersville
arheim	dunnville	karlin	maumee	padus	sylvester
au gres	eaupleine	kartsburg	mecan	pardeeville	symco
banat	elderon	kennan	mecosta	partridge	tarr
bellevue	elmound	kert	meehan	pearl	tedrow
billet	elm lake	keweenaw	menahga	pelkie	tilleda
bonduel	emmet	kickapoo	meridian	pence	trempe
boone	emmet	kidder	merrillan	pequaming	trempealeau
borth	ensign	kingsville	michigamme	peshekee	tula
boyer	ensley	kiva	mifflin	plainbo	urne
brems	fabius	kolberg	military	plainfield	vilas
brodale	fairchild	kranski	monico	renova	wainola
burkhardt	fairport	la farge	moquah	richford	waman
caryville	fisk	lamont	morocco	richter	warsaw
casco	fox	lapeer	mosinee	rodman	wasapi
ceresco	friendship	lawler	moundville	rosholt	watseka
champion	gale	leola	nadeau	rotamer	waukegan
chelsea	gifford	leroy	nahma	rousseau	whalan
chetek	gotham	lino	namur	rozelville	winneshiek
cloquet	graycalm	locke	nebago	rubicon	winterfield
coloma	halder	longrie	nemadji	salter	worcester
cohwood	hatley	lorenzo	nenno	santiago	wyckoff
cromwell	hesch	lows	newton	sarona	wyocena
croswell	hixton	ludington	northfield	sayner	
cunard	hochheim	lutzke	nymore	scott lake	
dakota	hoopston	magnor	oakville	shawano	

BEST AVAILABLE COPY

# SOIL RESTRICTION TABLE (continued)

## WYOMING

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table.

abernone	butche	eckley	hoodle	mughut	reno
abston	byrnle	edlin	hooper	nathale	renohill
adilis	cabin	ek mountain	hoot	naturita	rentsac
agneston	calicott	ek hollow	huguston	nelber	ret
alcova	camberge	ellicott	indart	nelson	rimton
allack	canburn	emblem	ipson	nezda	riverwash
alice	canwall	embry	irgul	newfork	rivra
allens park	carbol	enos	jayem	nihill	rock river
alliance	carmony	farisita	jenkins	nizina	rockinchair
almy	camero	fanson	julesburg	norte	roger
altvan	cass	feltner	kandaly	nunn	rohonda
amesmont	castello	feltonia	keeline	oceanet	rootel
anasazi	castner	firth	keyner	olney	rosane
ansel	cathedral	fiveoh	kezar	onason	rosebud
anzelmo	cedak	flathead	koonich	orpha	roxal
apeldorn	cestnik	fluatsch	labou	otero	ryan park
apron	chaperton	fola	lake creek	otterson	ryark
ard	chappell	forle	lakehelen	outlet	saddle
ascalon	ch. rids	formor	lamarsh	owen creek	santana
ashbon	charlos	fournile	lambman	pahlow	sapphire
ashuelot	cheadle	foxton	lander	paoli	satanka
assinniboine	cheyenne	fraddle	laney	parmele	sawcreek
attewan	chincap	frisco	lantis	parmlead	schamber
avalon	chinook	frontier	laporte	passcreek	schooner
badgerton	chittum	garita	larim	pena	scout
bankard	chugcreek	garrett	larimer	pepal	scravo
banks	cific	gas creek	las animas	pepton	seedskadee
barcus	clarkelen	gateson	lazeat	perceton	sharland
baroid	clayburn	gellie	leighcan	perma	sheege
barren	cliffsand	gerrard	libeg	pescar	shingle
bateson	cloud	glenberg	lininger	pesmore	shoshone
bayard	comer	glenderson	littlebe	peyton	simmont
bayerton	conpeak	glendive	littsan	philerson	slipman
beaverton	cordes	glenton	lohsman	pilotpeak	sobson
becks	corlett	goslin	olak	pinedale	southace
beeno	cotha	granile	lona	pinequest	southfork
beenom	cotopaxi	granyon	lucky	pishkum	space city
bigwin	coutis	greenman	luhon	plite	spangler
bigwinder	crago	grenoble	lupinto	poin	splitro
billycreek	cragosen	greyback	lymanson	poker	spool
binton	crestman	greybull	macfarlane	pokey	stoneham
blackhall	crowheart	greypoint	manter	preatorson	stunner
blaine	crownest	grieves	mathers	pugsley	sublette
bluerim	cryluha	grifi	maysdorf	quander	sudworth
bodorump	curecanti	grimm	mccaffery	quealman	sunup
bondman	curtis	grimstone	mccort	quealy	taglake
bonjea	cushman	grobutte	meadowlake	rainbolt	taluce
bosler	cushool	gunbarrel	means	rairdent	tanna
bottle	dahlquist	hagga	mema	randman	tassel
boulder	dailey	hapiack	millburne	ratake	tasselman
bowbac	dast	hardhart	millpot	rauzy	teagulf
bowen	dawny	hargreave	milren	rawah	teapo
boyle	deaver	haterion	mine	rawlins	teeler
breece	debone	havertel	miracle	redbank	teemat
briggsdale	decolney	hawksprings	mirror	redcreek	teewinot
bross	derrick	hawkstone	mishak	reddale	telecan
brownsto	diamondville	hazton	moran	redfeather	tellman
bruja	dix	hechtman	morset	redrob	tenorio
buffork	dobrow	heinsaw	moskee	redwash	teradi
bullock	draknab	herbman	mosroc	reeder	terro
burgess	driggs	hesperus	moloqua	regat	terry
burnt	dunday	hiland	mudray	rekop	tetonville
busher	dunul	hilltop	muff	relsob	thayne
bushman	dwyer	hobacker	mughouse	rencot	theedle

## SOIL RESTRICTION TABLE (continued)

WYOMING (continued)					
thermopolis	travessilla	truckton	vona	whaley	worka
thiel	travson	tullock	vonason	wiggletton	workman
threestop	trelna	tumback	wages	willowman	workstone
tieside	trembles	turner	walcott	willwood	workland
tigon	treoff	turnercrest	walden	wilsonville	wormser
tine	treon	turson	wall	windham	worthenton
tineman	tresano	uffens	wallock	winnett	yetull
tipper	trimad	uinta	walson	wirt	youga
tipperary	tripk	valent	wardboro	wishard	youjay
tisworth	trivar	valentine	wellsville	wolf	zillman
tolman	trook	vetal	wendover	woodrock	
tongue river	troutville	vible	westvaco	worf	

BEST AVAILABLE COPY

**NOTES**

28029

**BEST AVAILABLE COPY**

**BEST AVAILABLE C**

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T. W. Alexander Drive  
Research Triangle Park, North Carolina

Form No. AG  
Printed USA -

TEMIK is the registered trademark of Rhône-Poulenc for aldicarb pesticide  
© 1991 Rhône-Poulenc Ag Company