

RESTRICTED USE PESTICIDE: For retail sale to and use by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

PRODUCT BULLETIN

**UNION
CARBIDE**

UNION CARBIDE AGRICULTURAL
PRODUCTS COMPANY, INC.
T. W. ALEXANDER DRIVE
P.O. BOX 12014
RESEARCH TRIANGLE PARK, NC 27709

TEMIK 10G Aldicarb Pesticide
For Agricultural Use

EPA Reg. No. 264-331

Page 1 of 2

Supplemental Labeling For Control of Insects and Nematodes on Potatoes

Crop & Time of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet Of Row	Recommended Application
POTATOES AT PLANTING Do Not Apply In Suffolk County, Long Island, New York, At Anytime. Do Not Apply At Planting In Wisconsin or Maine (Except 10 lb Rate For Aphids in Maine Only)		34-Inch Row Spacing		Apply Granules With Seed Pieces In Planting Furrow and Cover With Soil.
	Aphids	10 (Maine Only)	10.5	
	Flea Beetles	30	31	
	Aphids, Colorado Potato Beetle, Leafhoppers	20 to 30	21 to 31	Apply As Above Or, Drill Granules 2 To 4 Inches On Both Sides of Seed Row And 3 To 8 Inches Deep (Usually 1 To 2 Inches Below Seed Pieces).
	Psyllids	20 (Arizona Only)	21	Apply As Above.
	Nematodes (Except Golden Nematode) (Root-Knot Nematodes Are Suppressed) (Verticillium Wilt Has Been Suppressed where Nematode Populations Have Been Reduced)	30	31	Apply Granules With Seed Pieces In Planting Furrow And Cover With Soil Or, Apply Granules In An 8-Inch Band And Work Into The Soil Or Cover With Soil To A Depth Of 4 Inches. Plant Seed Pieces In Treated Zone.
AFTER PLANTING In Maine and Wisconsin, Apply 4-6 Weeks After Planting.	Aphids, Colorado Potato Beetle, Leafhoppers	20 (Maine And Wisconsin Only)	21	Apply Granules in 4-6 Inch Band Over Row On Top Of Hill. Cover Granules With Soil By Cultivators Or Hilling Discs. DO NOT APPLY THIS PRODUCT TO ANY FIELD MORE THAN ONCE EVERY TWO YEARS.

A
 MAR 15 1982
 264-331

RESTRICTED USE PESTICIDE

TEMIK 10G Aldicarb Pesticide
For Agricultural Use

EPA Reg. No. 264-331

Page 2 of 2

Supplemental Labeling
For Control of Insects and Nematodes on Potatoes

Crop & Time of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
POTATOES AFTER PLANTING (continued)				
Other Than Maine And Wisconsin, Apply At 75% Emergence Or Before Plants Are 6 Inches High	Aphids, Colorado Potato Beetle	20 to 30	21 to 31	Side-Dress Granules 4 To 5 Inches On Both Sides Of Plant Row And 4 Inches Deep. Apply Where Pest Populations Are Expected 2-3 Weeks After Plant Emergence. Do Not Apply If At Planting Treatment Was Made.

PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS:

- Do not make more than one application per crop.
- In Maine and Wisconsin, do not apply to any field more than once every two years.
- Do not harvest within 90 days of an at planting application.
- Do not apply an "after planting" application within 50 days of harvest.
- Do not apply this product in Suffolk County, Long Island, New York.

DECOMPOSITION AND MOVEMENT IN SOIL

This product is readily decomposed into harmless residues under most use conditions: However, a combination of sandy and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F at application time, tend to reduce degradation and promote movement of residues to groundwater. If this describes your local use conditions and groundwater in your area is used for drinking, do not use this product without first contacting Union Carbide at 1 (800) 334-9769.

IMPORTANT:

Before using TEMIK 10G Aldicarb Pesticide, read and carefully observe the precautionary statements and all other information appearing on the product label. This bulletin contains new and supplemental directions for use of this product which may not appear on the package label. Follow the directions carefully.