

264-334

11/28/2005

1/25

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0080, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20480

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 264-334	2. EPA Product Manager John Hebert	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SEVIN® brand RP2 Carbaryl Insecticide	PM# Team 7	
5. Name and Address of Applicant (Include ZIP Code) Bayer CropScience P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION NOV 28 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of minor labeling changes per PR Notice 98-10 - REGFEE - Not applicable
CERTIFICATION STATEMENT: This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Danielle A. Larochelle	Title Registration Product Manager	Telephone No. (Include Area Code) (919) 549-2714
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Product Manager	
4. Typed Name Danielle A. Larochelle	5. Date November 14, 2005	

2/25

NOTIFICATION

NOV 28 2005

SEVIN® brand RP2 Carbaryl Insecticide

Home and Garden Insecticide

ACTIVE INGREDIENTS: Carbaryl (1-naphthyl N-methylcarbamate)22.5% by wt.
INERT INGREDIENTS:77.5% by wt.
(Contains 2 Pounds Carbaryl Per Gallon)

EPA Reg. No. 264-334

E.P.A. Est. No. 264-MO-02

- Kills Japanese Beetles
- Kills over 100 Listed Insects on Vegetables, Fruits, Ornamentals, and Lawns
- 16 oz. makes up to 24 gallons
- 32 oz. makes up to 48 gallons.
- Ready to Spray
- Bug Killer

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridine-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage. To aid in the confirmation of diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analysis will be arranged by Bayer CropScience. Consultation on therapy can be obtained at all hours by calling the appropriate emergency number 1-800-334-7577 (U.S.A.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (& DOMESTIC ANIMALS) CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH THE SKIN OR INHALED. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

3/25

User Safety Requirements

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Avoid contamination of food and feed.
- Do not allow children or pets to contact treated area until sprays have dried.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops and weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service or your local Bayer CropScience representative for further information.

DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Strictly observe label directions and cautions. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. The area being treated must be vacated by unprotected persons. Keep unprotected persons out of treated areas until sprays have dried. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Shake all containers prior to use. Do not reuse empty containers or measuring devices for other purposes. Apply when insects or damage appear. Repeat as necessary to maintain control, unless spray interval is specified.

Apply when air is calm to avoid drift and contact with eyes and skin.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.

For trees taller than 10 feet, consider hiring a licensed professional.

Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.

Do not repeat applications more than once a week.

STORAGE AND DISPOSAL

STORAGE

Store unused SEVIN® brand RP2 Carbaryl Insecticide in original container only, in cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100°F.

After use, replace cap on insecticide container. Rinse hose end unit outdoors in clear water. Store insecticide and sprayer assembly in a cool place.

STORAGE FOR A SINGLE DOSE PACKAGE:

Store packages of SEVIN® brand RP2 Carbaryl Insecticide in cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100°F.

PESTICIDE DISPOSAL

To dispose of partially filled containers: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL:

Do not reuse empty container. Place in trash or offer for recycling if available. Do not reuse hose end sprayer assembly for other purposes.

CONTAINER DISPOSAL FOR A PLASTIC DRUM:

Triple rinse (or equivalent). Then offer container for recycling or reconditioning or puncture and dispose of container in a sanitary landfill, by incineration, or, if allowed by State and local authorities, by burning. If container is burned, stay out of smoke.

CONTAINER DISPOSAL FOR A SINGLE DOSE PACKAGE:

Do not reuse empty container. Place in trash.

4/25

GENERAL CAUTIONS AND RESTRICTIONS

PLANT RESPONSE PRECAUTIONS

Application to wet foliage or during periods of high humidity may cause injury to tender foliage.

Do not use on Boston Ivy, Virginia creeper and maidenhair fern as injury may result. Carbaryl may also injure Virginia and sand pines.

The use of adjuvants may increase the potential for crop injury to sensitive crops.

PREHARVEST AND GRAZING RESTRICTIONS AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of labeled crops bearing probable carbaryl residues when this product is used in accordance with the label directions. If used as directed, treated forage may be grazed or used as feed for dairy and meat animals without causing illegal residues in meat or milk. Do not apply at greater rates or at more frequent intervals than stated on the label. To do so may result in illegal residues in crops, meat, and milk.

Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits on painted or stained surfaces or finishes (i.e., cars, houses, trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid applications to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

SPECIFIC USE DIRECTIONS

CROP/SITE GROUPINGS:

Asparagus

Brassica Leafy Vegetable Crops

Sweet Corn

Cucurbit Vegetables

Fruiting Vegetables

Leafy Vegetables

Legume Vegetables

Root and Tuber Crops (Root and Tuber Crops except Sugar Beets and Sweet Potatoes; Sweet Potatoes)

Small Fruits and Berries

Tree Fruit Crops (Citrus Fruits; Pome Fruits; Stone Fruits)

Tree Nut Crops

Trees and Ornamentals

Turfgrass

Control of Specific Pests Across Multiple Sites

Grasshoppers

Ticks which Vector Lyme Disease

Imported Fire Ants

Adult Mosquito Control

Nuisance Pest Control Around Building Perimeters

Directions for Use with a Hose-End Applicator

Vegetables except Asparagus and Sweet Corn

Asparagus

Sweet Corn

Small Fruits and Berries

Tree Fruit Crops

Turfgrass

Trees and Ornamentals

Control of Specific Pests across Multiple Sites

Ticks which Vector Lyme Disease

Imported Fire Ants

Adult Mosquito Control

Nuisance Pest Control Around Building Perimeters

Specific Use Directions for a Single Dose Package

Vegetables

Fruits

Trees, Ornamentals, Shrubs and Flowers

Pest Control Around Buildings and Other Outdoor Areas

5/25

ASPARAGUS

For use on Asparagus to control Apache cicada, Asparagus beetle, and Cutworms, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Repeat applications as necessary up to a total of 3 times prior to harvest or a total of 5 times per crop but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant - **OR** apply 3 to 6 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: *Application to ferns or brush growth following harvest of spears*: Repeat applications as necessary but not more often than every 7 days. Do not make more than a total of 5 applications per year to spears and ferns combined. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 1 day of harvest. Do not apply more than a total of 9 fluid ounces per 1000 SQ. FT. before harvest of spears. Do not apply more than a total of 15 fluid ounces per 1000 sq. ft. per year.

BRASSICA LEAFY VEGETABLES CROPS

For use on Broccoli, Brussel Sprouts, Cauliflower, Cabbage, Chinese Cabbage, Collards, Kale, Kohlrabi, and Mustard Greens to control Flea beetles, Harlequin bugs, and Leafhoppers, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** Armyworm, Aster leafhopper, Corn earworm, Diamondback moth, Fall armyworm, Imported cabbageworm, Lygus bug, Spittle bugs, Stink bugs, and Tarnished plant bug apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Repeat applications as needed up to a total of 4 times but not more than once every 7 days. **RESTRICTIONS AND PRECAUTIONS**: For Broccoli, Brussel Sprouts, Cabbage Cauliflower, and Kohlrabi, do not apply within 3 days of harvest. For Chinese Cabbage, Collards, Kale, and Mustard Greens, do not apply within 14 days of harvest. Do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per crop.

SWEET CORN

For use on sweet corn to control Armyworm, Chinch bugs, Corn earworm, Corn rootworm adults, Fall armyworm, Flea beetles, Japanese beetles, Sap beetles, Southwestern corn borer, and Leafhoppers, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control European corn borer, apply 2 1/4 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Western bean cutworm and Cutworms, apply 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: **OBSERVE BEE CAUTION**. Repeat applications as necessary up to a total of 8 times but not more often than once every 7 days. Optimum timing and good coverage are essential for effective control. For insects attacking silks and ears, insecticide sprays should be applied starting when first silks appear and continuing until silks begin to dry. During silking, the minimum retreatment interval (7 days) may not provide adequate levels of protection under conditions of rapid growth or severe pest pressure. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 2 days of harvest of ears or within 14 days of harvest or grazing of forage or within 48 days of harvest of fodder. Do not apply more than a total of 24 fluid ounces per 1000 sq. ft. per crop.

CUCURBIT VEGETABLES

For use on Cucumbers, Melons, Pumpkins, and Squash to control Pickleworm and Melonworm, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Cucumber beetles, Flea beetles, Leafhoppers and Squash bugs, apply 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days. For optimum control of squash bugs, apply sufficient spray volume for thorough coverage and time sprays for early morning or late afternoon. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 3 days of harvest. Do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per crop. Observe plant response precautions.

FRUITING VEGETABLES

For use on Tomatoes, Peppers, and Eggplant to control Colorado potato beetle, European corn borer, Fall armyworm, Lace bugs, Stink bugs (suppression), Tarnished plant bug, Thrips (suppression), Tomato fruitworm, Tomato hornworm, and Tomato pinworm, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Flea beetles and Leafhoppers, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Cutworms, apply 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Repeat applications as necessary up to a total of 7 times but not more often than once every 7 days. Thorough coverage is essential to effectively suppress stink bugs. When disease transmission is suspected, monitor fields following application and retreat if reinfestation occurs but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 3 days of harvest. Do not apply more than a total of 12 fluid ounces per 1000 sq. ft. per crop.

LEAFY VEGETABLES

For use on Celery, Dandelion, Endive, Lettuce (head and leaf), Parsley, Spinach, and Swiss chard to control Flea beetles, Harlequin bugs, and Leafhoppers, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** for use on Celery, Dandelion, Endive, Lettuce (head and leaf), Parsley, Spinach, and Swiss Chard to control Armyworm, Aster leafhopper, Fall armyworm, Imported cabbageworm, Lygus bugs, Spittlebugs, Stink bugs, and Tarnished plant bug, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Repeat applications as necessary up to a total of 5 times but not more than once every 7 days. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 14 days of harvest. Do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per crop.

6/25

LEGUME VEGETABLES

For use on Fresh and Dried Beans (Phaseolus species including snap, navy, and kidney), Fresh and Dried Peas (Pisum species), Lentils, Cowpeas, and Southern Peas; (DO NOT USE ON LENTILS IN CALIFORNIA), to control Bean leaf beetle, Blister beetle, Cucumber beetles, Grape colaspis, Green cloverworm, Japanese beetle, Mexican bean beetle, and Velvetbean caterpillar, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Corn earworm, apply 3/4 to 2 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Alfalfa caterpillar, Colorado potato beetle, Flea beetles, Leafhoppers, Three cornered alfalfa hopper, Thrips and Western bean cutworm, apply 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Armyworm, Cutworms, European corn borer, Fall armyworm, Stink bugs, Tarnished plant bug, and Webworms, apply 1 1/2 to 2 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Alfalfa looper (suppression), Cowpea curculio (suppression), Painted lady (Thistle caterpillar), Pea leaf weevil, Pea weevil, Saltmarsh caterpillar, Woollybean caterpillar, and Yellowstriped armyworm, apply 2 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR (California Only)** Corn earworm (suppression), Limabean podborer (suppression), Lygus bugs (suppression), and Stink bugs (suppression), apply 2 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** Repeat applications as necessary up to a total of 4 times but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects and spiders. Use the higher rates for heavy populations and larger instars. **RESTRICTIONS AND PRECAUTIONS:** Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of fresh beans or peas or within 21 days of harvest of dried beans or peas, seed, or hay. Do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per crop. Observe plant response precautions.

ROOT AND TUBER CROPS: ROOT AND TUBER CROPS EXCEPT SUGAR BEETS & SWEET POTATOES

For use on Garden Beets, Carrots, Horseradish, Parsnips, Radishes, Rutabagas, Salsify, and Potatoes to control Flea beetles and Leafhoppers, apply 3/4 to 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Armyworm, Aster leafhopper, Colorado potato beetle, Corn earworm, Cutworms, European corn borer, Fall armyworm, Lace bugs, Lygus bugs, Spittlebugs, Stink bugs, and Tarnished plant bug, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** Repeat applications as necessary up to a total of 6 times but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. **RESTRICTIONS AND PRECAUTIONS:** Do not apply within 7 days of harvest. Do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per crop.

SWEET POTATOES

For use on Sweet Potatoes to control Corn earworm, Cucumber beetles, Flea beetles, Sweet potato hornworm, Sweet potato weevil, Tortoise beetles, and Whitefringed beetle, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Yellowstriped armyworm, apply 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** Preplant dip for control of sweet potato weevil: Just prior to planting, dip sweet potato cuttings in a suspension containing 4 gallons of this product in 100 gallons of water (5.2 fluid ounces of this product per gallon of water). For foliar sprays, repeat applications as necessary up to a total of 8 times but not more often than once every 7 days. **RESTRICTIONS AND PRECAUTIONS:** Do not apply within 7 days of harvest. Do not apply more than a total of 12 fluid ounces per 1000 sq. ft. per crop with in-season sprays. Do not apply more than a total of 3 3/5 fluid ounces per 1000 sq. ft. as a preplant dip treatment.

SMALL FRUITS AND BERRIES

For use on Canberries, Blueberries, Cranberries, Grapes and Strawberries to control European fruit lecanium, European raspberry aphid, Flea beetles, Grape leafhopper, Grape leafroller, Japanese beetle, Leafhoppers, Leafrollers, Meadow spittlebug, Omnivorous leafhopper, Rose chafer, Snowy tree cricket, Strawberry bud weevil, Strawberry clipper, Strawberry fruitworm, Strawberry leafroller, Strawberry weevil, Western grapeleaf skeletonizer, and Western yellowstriped armyworm, apply 1 1/2 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Blueberry maggot, Cherry fruitworm, Cranberry fireworm, Cranberry fruitworms, Cranberry twig girdler, Elm spanworm, Gypsy moth, Spaganothus worm, and Tarnished plant bug, apply 2 1/4 to 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Eight-spotted forester, Cutworms, Grape berry moth, June beetles, Omnivorous leafroller, Orange tortrix, Raspberry fruitworm, Raspberry sawfly, Redbanded leafroller, and Saltmarsh caterpillar, apply 3 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. In grapes, do not concentrate spray on the bunch or visible residues may result. **RESTRICTIONS AND PRECAUTIONS:** Do not apply within 7 days of harvest. Do not apply more than a total of 15 fluid ounces per 1000 sq. ft. per crop. CAUTION: Use in cranberries may kill shrimp and crabs. Do not use in areas where these are important resources. Carbaryl may injure Early Dawn and Sunrise varieties of strawberries.

TREE FRUIT CROPS

On all tree fruit crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors.

CITRUS FRUITS:

For use on Citrus Fruits to control Avocado leafroller, California orangedog, Citrus cutworm, Fruittree leafroller, Orange tortrix, and Western tussock moth, apply 3 to 4 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 8 times but not more often than once every 14 days - **OR** to control Citrus rust mite, Eriophyid mites, Plant bugs, and Scale insects (Black scale, brown soft scale, California red scale (except in California), citrus snow scale, yellow scale (except in California)), apply 4 1/2 to 7 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Apopka weevil (adult), Citrus root weevils (adults), Fuller rose beetle, and Little leaf notcher (adult), apply 7 1/2 to 11 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: For scale control, apply when crawlers are present. For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low. - **OR** (for use in California Only): California red scale and Yellow Scale, apply 7 1/2 to 24 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: Do not make more than 1 application per season for California red scale. Apply when crawlers are present. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 5 days of harvest. Do not apply more than a total of 30 fluid ounces per 1000 sq. ft. per crop. For trees taller than 10 feet, consider hiring a licensed professional.

POME FRUITS:

For use on Apples, Pears, Loquats, Crabapples, and Oriental Pears to control White apple leafhopper, apply 3/4 to 2 1/4 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Apple aphid and Codling moth, apply 1 1/2 to 4 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Apple aphid, Apple maggot, Apple mealybug, Apple rust mite, Bagworms, California pearslug (pear sawfly), European apple sawfly, Eyespotted bug moth, Fruittree leafroller, Gypsy moth, Japanese beetle, Lesser appleworm, Lygus bugs, Orange tortrix, Pearleaf blister mite; Pear psylla, Pear rust mite, Periodical cicada, Plum curculio, Redbanded leafroller, Rcsy apply aphid, Scale insects (Forbes scale, Lecanium scale, San Jose scale), Tarnished plant bug, Tentiform leafminers, Woolly apple aphid, and Yellowheaded fireworm, apply 2 1/4 to 4 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: OBSERVE BEE CAUTION. On apples, avoid use during the period from full bloom until 30 days after full bloom unless fruit thinning is desired. Use for pest control during this period also may result in fruit removal. Repeat applications as necessary up to a total of 8 times per crop (including thinning sprays on apples) but not more often than once every 14 days. For psylla control, apply when eggs hatch or young nymphs are present. For scale control, apply when crawlers are present. - **OR** for use on Apples Only - For Fruit Thinning, apply 1 1/2 to 4 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: OBSERVE BEE CAUTION. For easily thinned varieties: apply 1/5 to 1/3 fluid ounce per 1000 sq. ft. For difficult to thin varieties: apply 1/3 to 2/3 fluid ounce per 1000 sq. ft. Apply between 10 and 25 days after full bloom. Factors such as tree age, variety, nutrition, previous crop, pruning, bloom and degree of set favor excessive fruit thinning with this product. Exercise caution to avoid possible yield reduction. Rates may vary depending on variety and local orchard conditions. Consult with your County Extension Service or other experts for advise on the proper use of this product. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 3 days of harvest. Do not apply more than a total of 22 1/2 fluid ounces per 1000 sq. ft. per crop. Do not make more than a total of 8 applications per crop. For trees taller than 10 feet, consider hiring a licensed professional.

STONE FRUITS

For use on Apricots, Cherries, Nectarines, Peaches, Plums and Prunes to control Apple pandemis, Black cherry aphid, Cherry fruitworm, Cherry maggot (Cherry fruit fly), Codling moth, Cucumber beetles, Eastern tent caterpillar, Eyespotted bug moth, European earwig, Fruittree leafroller, Green fruitworm, Gypsy moth, Japanese beetle, June beetle, Lesser peachtree borer, Mealy plum aphid, Orange tortrix, Oriental fruit moth, Peach twig borer, Periodical cicada, Plum curculio, Prune leafhopper, Redbanded leafroller, Rose chafer, Scale insects (Brown soft scale, Forbes scale, Lecanium scale, Olive scale, Oystershell scale, San Jose scale), Tarnished plant bug, Tussock moth, and Variegated leafroller, apply 3 to 4 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** (for California Only) Black cherry aphid, Cherry fruitworm, Cherry maggot (Cherry fruit fly), Codling moth, Cucumber beetles, Eyespotted bug moth, European earwig, Fruittree leafroller, Green fruitworm, Mealy plum aphid, Orange tortrix, Oriental fruit moth, Peach twig borer, Scale insects (Brown soft scale, Forbes scale, Lecanium scale, Olive scale, Oystershell scale, San Jose scale), Tarnished plant bug, and Tussock moth, apply 4 1/2 to 6 fluid ounces per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 3 times per crop but not more often than once every 14 days. An additional application at the dormant or delayed dormant timing may be made. For optimum scale control, apply when crawlers are present. - **OR** to control Peach twig borer, Scale insects (Brown soft scale, Forbes scale, Lecanium scale, Olive scale, Oystershell scale, San Jose scale), apply 6 to 7 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS**: For dormant or delayed dormant timing, apply in combination with a recommended dormant oil. Refer to the dormant oil product label for additional use directions and restrictions. **RESTRICTIONS AND PRECAUTIONS**: Do not apply within 3 days of harvest, except in California. In California, do not apply within 1 day of harvest. Do not apply more than a total of 21 fluid ounces per 1000 sq. ft. per crop. Do not apply more than a total of 7 1/2 fluid ounces per 1000 sq. ft. at the dormant or delayed dormant timing. Do not apply more than a total of 13 1/2 fluid ounces per 1000 sq. ft. during the production season. For trees taller than 10 feet, consider hiring a licensed professional.

TREE NUT CROPS

On all tree nut crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors.

TREE NUTS

For use on Almonds, Chestnuts, Filberts, Pecans, and Walnuts to control Black margined aphid, Calico scale, Codling moth, European fruit lecanium, Fall webworm, Filbert aphid, Filbert leafroller, Filbertworm, Frosted scale, Fruittree leafroller, Hickory shuckworm, Lesser webworm, Navel orangeworm, Peach twig borer, Pecan leaf phylloxera, Pecan stem phylloxera, Pecan nut casebearer, Pecan spittlebug, Pecan weevil, San Jose scale, Twig girdler, and Walnut caterpillar apply 3 to 7 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS: OBSERVE BEE CAUTION.** Repeat applications as necessary up to a total of 4 times per crop (except at the dormant or delayed dormant timing) but not more often than once every 7 days. Use lower rates for pests attacking leaves. Use higher rates for pests attacking fruit and for higher infestations. For scale control, apply when crawlers are present. For peach twig borer, best results with foliar applications have been found by making applications in "popcorn" or petal fall stages when the May brood begins to hatch. For navel orangeworm in almonds and walnuts, best results have been found by timing early and midseason applications to correspond with moth flight peaks. For filbert leafroller, best results have been found by making applications when eggs are hatching, repeating application on first appearance of moths and again 3 to 4 weeks later. For codling moth in walnuts, best results have been found by making applications when average cross-sectional diameters of developing nuts are 0.5 to 0.75 inches and again during middle or late June as needed. - **OR** to control Chestnut weevil and European earwig, apply 6 to 7 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** For chestnut weevil, best results have been found with 4 applications at weekly intervals beginning in late July. The last application should be made prior to shuck split. For European earwig, thorough coverage of trunks, branches, and nuts is needed for best results. For use on Almonds ONLY, to control Peach Twig Borer and Scale insects apply 6 to 7 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** For dormant or delayed dormant timing, apply in combination with a recommended dormant oil. Refer to the dormant oil product label for additional use directions and restrictions. **RESTRICTIONS AND PRECAUTIONS:** Do not apply within 14 days of harvest. Do not apply more than a total of 22 1/2 fluid ounces per 1000 sq. ft. per crop, including any application at the dormant or delayed dormant timing. For trees taller than 10 feet, consider hiring a licensed professional.

TREES AND ORNAMENTALS

Apply in sufficient volume for adequate coverage. This will vary depending on the plant or tree size, density, and stage of growth. For use on Roses, Flowers, Other Herbaceous Plants, Woody Plants and Shrubs to control Ants, Apple aphid, Armyworm, Ash whitefly, Azalea leafminer, Bagworms, Balsam twig aphid, Birch leafminer, Blister beetle, Boxelder bug, Boxwood leafminer, Brown tail moth, Cankerworms, Catalpa sphinx, Chiggers, Cooley spruce gall adelgid, Cutworms, Cypress tip moth, Douglas-fir tussock moth, Eastern spruce gall adelgid, Elm leaf aphid, Elm leaf beetle, Elm spanworm, Eriophyid mites, European pine shoot moth, Fall armyworm, Flea beetle, Fuchsia gall mite, Fuller rose beetle, Gall midges, Gall wasps, Greenstriped mapleworm, Hackberry nipplegall maker, Holly bud moth, Holly leafminer, Jackpine budworm, Japanese beetle, Jeffrey pine needleminer, June bugs, Lace bugs, Leafhoppers, Leafrollers, Locust borer, Maple leafcutter, Mealy bugs, Mimosa webworm, Nantucket pine tip moth, Oak leafminers, Oak moth, Oak skeletonizer, Oakworm complex, Oleander caterpillar, Olive ash borer, Orange-striped oakworm, Periodical cicada, Pine looper, Pine sawfly, Pine spittlebug, Pitch pine tip moth, Plant bugs, Poinsettia hornworm, Psyllids, Puss caterpillar, Redhumped oakworm, Rose aphid, Rose chafer, Rose slug, Sadd'ed prominent, Sawflies (exposed), Scale insects (crawlers), Sowbugs, Spiney elm caterpillar, Springtails, Spruce budworm, Spruce needleminer, Subtropical pine tip moth, Tent caterpillars, Thorn bug, Thrips (exposed), Ticks, Walnut caterpillar, Webworms, Western hemlock looper, Western spruce budworm, Willow leaf beetles, Woolly gall aphid, and Yellow poplar weevil using 1 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** for control of Gypsy Moth, apply 1 to 1/2 fluid ounces of this product per gallon of water to cover 1000 sq. ft using the following **SPECIFIC DIRECTIONS:** Observe plant response precautions. Obtain thorough coverage of upper and lower leaf surfaces. The addition of a sticker may improve residual control. To control scale insects, treat trunks, stems, and twigs in addition to plant foliage. For optimum worm control, treat when pests are small. Do not use on syrup-producing sugar maples where sap is harvested. Applications for control of maple leafcutters on sugar maple should be made when larvae are in 2nd instar after mining and as cases are being formed. Repeat treatments as necessary up to a total of 6 times per year but not more often than once every 7 days. For gypsy moth control, use the higher rate for heavy infestations. **RESTRICTIONS AND PRECAUTIONS:** Do not make more than a total of 6 foliar treatments per year. **DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.** For trees taller than 10 feet, consider hiring a licensed professional.

TURFGRASS

Apply the recommended rates using pressure-type equipment in sufficient spray volume for thorough coverage and thatch penetration. For use on Domestic lawns and Recreational areas to control Ants, Armyworms, Centipedes, Chiggers, Cutworms, Earwigs, Essex skipper, European chafer, Fall armyworm, Fiery skipper, Grasshoppers, Green june beetle larvae, Leafhoppers, Lucerne moth, Millipedes, Mosquitoes (adults), Scarab beetle adults (May beetle, June beetle, Japanese beetle, Green june beetle), Sowbugs, Spittlebugs, Springtails, Ticks and Yellowstriped armyworm, apply 3 to 6 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** For armyworm, cutworm and fall armyworm control: Do not irrigate treated areas within 24 hours following insecticide application. For green june beetle grub (larvae) control: Make applications when grubs are feeding near the soil surface. Water or irrigate turfgrass soon after treatment. Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days. - **OR** to control Chinch bugs, and Sod webworms (lawn moths), apply 12 to 16 fluid ounces of this product per gallon of water to cover 1000 sq. ft. - **OR** to control Bluegrass billbug larvae, European crane fly, Fleas, White grubs (Japanese beetle, Chafer beetle, and Phyllophaga spp. larvae), Black turfgrass ataenius beetle larvae, and Hyperoides weevil (bluegrass weevil) larvae, apply 12 fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** For chinch bug control: Treat entire grass area rather than just damaged areas. Irrigation prior to treatment will aid in penetration of insecticide into turfgrass. Do not irrigate treated areas within 24 hours following insecticide application. For european crane fly control: Treatment should be applied in early spring, April 1 to April 15, or as recommended by local Agricultural Extension Service agents. Water or irrigate turfgrass soon after treatment. For White Grub Control: Applications should be made when grubs are feeding near the soil surfaces, usually during late March through May, or July through early September,

9/25

or as recommended by local Agricultural Extension Service agents. Water or irrigate turfgrass soon after treatment. Repeat applications as necessary up to 2 times per year but not more often than once every 7 days. **RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES

GRASSHOPPERS:

For use on all crops and sites on this label to control Grasshoppers, apply 3/4 to 2 1/4* fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** Apply 3/4 to 1 1/8 fluid ounces per 1000 sq. ft. of this product for nymphs on small plants or sparse vegetation. Apply 1 1/2 to 2 1/4 fluid ounces per 1000 sq. ft. for mature grasshoppers or applications to dense foliage or if extended residual control is desired. Be certain spray volumes are appropriate to assure adequate coverage. **RESTRICTIONS AND PRECAUTIONS:** *NOTE: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions.

CONTROL OF TICKS WHICH VECTOR LYME DISEASE

For control of juvenile and adult ticks which vector Lyme Disease, apply the recommended amount in sufficient volume for thorough coverage.

For use on all crops and sites on this label to control Ixodes spp. (Deer tick, Bear tick, Black legged tick) and Amblyomma spp. (Lone star tick) apply 1 1/2 to 3* fluid ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** *Use the high rate for heavy tick infestations. Use higher spray volumes for dense ground cover or heavy leaf litter. Target applications for nymphal control in late spring or early summer. Control of adult tick can be obtained with late summer and fall applications. Do not use spot treatments. Treat entire area and perimeter areas where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary to maintain adequate control levels.* **RESTRICTIONS AND PRECAUTIONS:** *NOTE: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED.

IMPORTED FIRE ANTS

For use as a Drench Application on all crops and sites on this label to control Imported fire ants, apply 3/4 fluid ounces of this product per gallon of water using the following **SPECIFIC DIRECTIONS:** Apply a total of 2 gallons of the diluted solution over the surface of each mound or at least 1 quart per 6 inches of mound diameter using a bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding areas to a 4 ft. diameter (12 sq. ft.). Do not disturb mound prior to treatment. Pour solution from a height of about three feet to give sufficient force to break mound apex and flow into ant tunnels. For best results apply in cool weather (65 - 80 degrees F) or in early morning or late evening hours. Repeat application if mound activity resumes after 7 days. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness. - **OR** for use as a Foliar Application, apply 1 1/2 fluid ounces (approximately 3 tablespoons) of this product per gallon of water using the following **SPECIFIC DIRECTIONS:** Avoid contact with foliage and treat only the growing media when using on bedding plants. Do not make more than one application, either as a root dip or drench treatments (applied to the point of saturation). **RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL DRENCH HAS DRIED. DO NOT USE IN GREENHOUSES. *NOTE: DO NOT USE ON ANY FOOD CROP NOT LISTED ON THIS LABEL. Refer to the specific crop section for additional restrictions and precautions.

ADULT MOSQUITO CONTROL

Apply in sufficient gallonage for thorough coverage. For use on all crops and sites on this label to control Mosquitoes(adults), apply 3/8 to 1 1/2* ounces of this product per gallon of water to cover 1000 sq. ft. using the following **SPECIFIC DIRECTIONS:** **OBSERVE BEE CAUTION.** Treat shrubbery and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early mornings or late evenings. Repeat applications as necessary.* **RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED. **CAUTION:** May kill shrimp and crabs. Do not use in areas where these are important resources. *NOTE: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions. To prepare small volumes of spray mixture, use 1/2 fl. oz. (approximately 4 teaspoons) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1 1/2 fluid ounces per 1000 sq. ft. are indicated.

NUISANCE PEST CONTROL

For use as a perimeter treatment to prevent infestations of buildings, apply the specified dosage in a band 6 to 10 feet wide around the outside perimeter of the structure.

For use on Ants, Bees, Brown dog tick, Centipedes, Cockroaches, Crickets, Darkling beetle (lesser mealworm or litter beetle), Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Spiders, and Wasps, apply 5 fluid ounces of this product per gallon using the following **SPECIFIC DIRECTIONS:** Observe plant response precautions. Apply as a coarse wet spray in a bank 6 to 10 feet wide around the outside perimeter of buildings. Confine applications to areas immediately adjacent to the building. Direct application to the structure should be minimal and restricted to cracks, crevices, and areas where insects tend to congregate. For use in and around flower beds and ornamental plantings, apply as a coarse wet spray in and around plant beds. May be applied to outdoor sleeping quarters of pets. Do not treat animals. Do not allow animals to use treated area until sprays have dried. **RESTRICTIONS AND PRECAUTIONS:** DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED. **NOTE:** Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits

10/25

on painted or stained surfaces of finishes (i.e., cars, house trailers, boats, etc.) should be removed immediately by washing to prevent discoloration. Avoid application to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

DIRECTIONS FOR USE WITH HOSE END APPLICATOR

- Kills Japanese Beetles
- Kills over 100 insects on Vegetables, Fruits, Ornamentials and Lawns
- Covers up to 1,000 square feet
- Ready To Spray
- Bug Killer

HOSE END SPRAYER INSTRUCTIONS

Shake before using.

How To Use:

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water".
4. Point nozzle toward plants and turn water control knob to "On". Product will automatically mix with water. Spray upper and lower leaf surfaces thoroughly."

W/85

VEGETABLES EXCEPT ASPARAGUS AND SWEET CORN

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Broccoli Brussel Sprouts Cauliflower Cabbage Celery Chinese Cabbage Collards Cowpeas Cucumbers Dandelion Eggplant Endive Fresh and Dried Beans (<i>Phaseolus</i> species including snap, navy and kidney) Fresh and Dried Peas (<i>Pisum</i> species) Kale Kohlrabi Lettuce (head and leaf) Lentils Melons Mustard Greens Parsley Peppers Pumpkins Tomatoes Root and Tuber Crops: Garden Beets, Carrots, Horseradish, Parsnips, Radishes, Rutabagas, Salsify, Potatoes Southern Peas Spinach Squash Sweet Potatoes Swiss Chard	Alfalfa caterpillar Armyworm Aster leafhopper Bean leaf beetle Blister beetle Colorado Potato beetle Cucumber beetles Cutworms Diamondback moth European corn borer Fall armyworm Flea beetles Grape colapsis Green cloverworm Harlequin bug Imported cabbageworm Japanese beetle Lace bugs Leafhoppers Melonworm Mexican bean beetle Pickleworm Rednecked peanutworm Spittlebugs Squash bugs Sweet potato hornworm Sweet potato weevil Tarnished plant bugs Three cornered alfalfa hopper Thrips Tomato fruitworm Tomato hornworm Tomato pinworm Tortoise beetles Velvetbean caterpillar Webworms Western bean cutworm Whitefringed beetle adults	1 1/2	For optimum control of squash bugs time sprays for early morning or late afternoon. When disease transmission is suspected, monitor plants following application and retreat if reinfestation occurs. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. For optimum control of thrips, ensure adequate coverage for the underside of leaves.

RESTRICTIONS AND PRECAUTIONS: VEGETABLES

- For Broccoli, Brussel Sprouts, Cauliflower, Cabbage, Cucumbers, Eggplant, Kohlrabi, Melons, Peppers, Pumpkins, Tomatoes and Squash do not apply within 3 days of harvest.
- For Fresh Beans and Peas, do not apply within 3 days of harvest. For Dried Beans and Peas, do not apply within 21 days of harvest.
- For Root and Tuber Crops and Sweet Potatoes do not apply within 7 days of harvest.
- For Celery, Chinese Cabbage, Collards, Dandelion, Endive, Kale, Lettuce (head and leaf), Mustard Greens, Parsley, Spinach and Swiss Chard do not apply within 14 days of harvest.
- For all vegetables do not make more than a total of 4 applications per year.
- For all vegetables do not apply more than a total of 9 fluid ounces per 1000 sq. ft. per year.
- Do not apply more often than once every 7 days.

12/25

DIRECTIONS FOR USE WITH HOSE END APPLICATOR, CONTINUED

ASPARAGUS

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Asparagus	Apache cicada Asparagus beetle Cutworms	3	Repeat applications as necessary up to a total of 3 times prior to harvest or a total of 5 times per crop but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant.
		6	Application to ferns or brush growth following harvest of spears: Repeat applications as necessary but not more often than once every 7 days. Do not make more than a total of 5 applications per year to spears and ferns combined.

RESTRICTIONS AND PRECAUTIONS: ASPARAGUS

- Do not apply within 1 day of harvest.
- Do not apply more than a total of 9 fluid ounces per 1000 SQ. FT. before harvest of spears.
- Do not apply more than a total of 15 fluid ounces per 1000 SQ. FT. per year.

SWEET CORN

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Sweet Corn	Armyworm Chinch bugs Corn earworm Corn rootworm adults Fall armyworm Flea beetles	Japanese beetle Sap beetles Southwestern corn borer Leafhoppers	OBSERVE BEE CAUTION Repeat applications as necessary up to a total of 8 times but not more often than once every 7 days. Optimum timing and good coverage are essential for effective control For insects attacking silks and ears, insecticide sprays should be applied starting when first silks appear and continuing until silks begin to dry. During silking, the minimum retreatment interval (7 days) may not provide adequate levels of protection under conditions of rapid growth or severe pest pressure. The use of an alternative product should be considered in conjunction with this product.
	European corn borer	3	
	Western bean cutworm	Cutworms	

RESTRICTIONS AND PRECAUTIONS: SWEET CORN

- Do not apply within 2 days of harvest of ears or within 14 days of harvest or grazing of forage.
- Do not apply more than a total of 24 fluid ounces per 1000 SQ. FT. per crop.

DIRECTIONS FOR USE WITH HOSE END APPLICATOR, CONTINUED

SMALL FRUITS AND BERRIES

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Small Fruits and Berries: Caneberries, Blueberries, Cranberries, Grapes, Strawberries	Blueberry maggot Cherry fruitworm Cranberry fireworm Cranberry fruitworms Cranberry twig girdler Cutworms Eight-spotted forester Elm spanworm European fruit lecanium European raspberry aphid Flea beetles Grape berry moth Grape leaf folder Grape leafroller Gypsy moth Japanese beetle June beetles Leafhoppers Leafrollers Meadow spittlebug Omnivorous leaf tier	Omnivorous leafroller Orange tortrix Raspberry fruitworm Raspberry sawfly Redbanded leafroller Rose chafer Saltmarsh caterpillar Snowy tree cricket Spaganothus worm Strawberry bud weevil Strawberry clipper Strawberry fruitworm Strawberry leafroller Strawberry weevil Tarnished plant bug Western grapeleaf skeletonizer Western yellowstriped armyworm	3 OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 5 times but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. In grapes for grape leaf folder control, apply before first brood larvae emerge from rolls. In grapes, do not concentrate spray on the bunch or visible residues may result.

RESTRICTIONS AND PRECAUTIONS: SMALL FRUITS AND BERRIES

- Do not apply within 7 days of harvest.
- Do not apply more than a total of 15 fluid ounces per 1000 SQ. FT. per crop.
- CAUTION: Use in cranberries may kill shrimp and crabs. Do not use in areas where these are important resources.
- Carbaryl may injure Early Dawn and Sunrise varieties of strawberries.

14/25

DIRECTIONS FOR USE WITH HOSE END APPLICATOR, CONTINUED

TREE FRUIT CROPS

On all tree fruit crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors.

CITRUS FRUITS

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Citrus Fruits	Apopka weevil (adult) Avocado leafroller California orangedog Citrus cutworm Citrus root weevils (adults) Citrus rust mite Eriophyid mites Fruittree leafroller Fuller Rose Beetle Little leaf notcher (adult) Orange Tortrix Plant bugs Scale insects (Black scale, brown soft scale, California red scale (except in California), citrus snow scale, yellow scale) Western tussock moth	7 1/2	OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 8 times but not more often than once every 14 days. Use lower rates for pests attacking leaves. Use higher rates for pests attacking fruit and for higher infestations. For scale control, apply when crawlers are present. For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low.
	California only: California red scale	7 1/2	Do not make more than 1 application per season. Apply when crawlers are present.

RESTRICTIONS AND PRECAUTIONS: CITRUS FRUITS

- For trees taller than 10 feet, consider hiring a licensed professional.
- Do not apply within 5 days of harvest.
- Do not apply more than a total of 30 fluid ounces per 1000 SQ. FT. per crop.

15/25

DIRECTIONS FOR USE WITH HOSE END APPLICATOR, CONTINUED

STONE FRUIT AND POME FRUIT

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
<p>Stone Fruits: Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes</p> <p>Pome Fruits: Apples, Pears, Loquats, Crabapples, and Oriental Pears.</p>	<p>Apple aphid Apple maggot Apple mealybug Apple pandemis Apple rust mite Bagworms Black cherry aphid California pearslug (pear sawfly) Cherry fruitworm Cherry maggot (Cherry fruit fly) Codling moth Cucumber beetles Eastern tent caterpillar Eyespotted bud moth European apple sawfly European earwig Fruittree leafroller Green fruitworm Gypsy moth Japanese beetle June beetle Lesser appleworm Lesser peachtree borer Lygus bugs Mealy plum aphid</p> <p>Orange tortrix Oriental fruit moth Peach twig borer Pearleaf blister mite Pear psylla Pear rust mite Periodical cicada Plum curculio Prune leafhopper Redbanded leafroller Rose chafer Rosey apple aphid Scale insects (Brown soft scale, Forbes scale, Lecanium scale, Olive scale, Oystershell scale, San Jose scale) Tarnished plant bug Tentiform leafminers Tussock moth Variegated leafroller White apple leafhopper Woolly apple aphid Yellowheaded fireworm</p>	<p>4 1/2</p>	<p>OBSERVE BEE CAUTION.</p> <p>Repeat applications as necessary up to a total of 3 times per crop but not more often than once every 7 days. An additional application at the dormant or delayed dormant timing may be made.</p> <p>For optimum scale control, apply when crawlers are present.</p> <p>For lesser peachtree borer, best results have been found by thoroughly spraying limbs and tree trunks at weekly intervals during moth flight.</p> <p>For psylla control, apply when eggs hatch or young nymphs are present.</p>

RESTRICTIONS AND PRECAUTIONS: STONE FRUIT AND POME FRUIT

- For trees taller than 10 feet, consider hiring a licensed professional.
- Do not apply within 3 days of harvest.
- Do not apply more than a total of 21 fluid ounces per 1000 SQ. FT. per crop.
- Do not apply more than a total of 13 1/2 fluid ounces per 1000 SQ. FT. during the production season.

16/25

DIRECTIONS FOR USE WITH HOSE END APPLICATOR, CONTINUED

TREE NUT CROPS

On all tree nut crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors.

TREE NUTS

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Tree Nuts: Almonds, Chestnuts, Filberts, Pecans, Walnuts	Black margined aphid Brown soft scale Calico scale Chestnut weevil Codling moth European earwig European fruit lecanium Fall webworm Filbert aphid Filbert leafroller Filbertworm Frosted scale Fruittree leafroller Hickory shuckworm Lecanium Lesser webworm Navel orangeworm Peach twig borer Pecan leaf phylloxera Pecan stem phylloxera Pecan nut casebearer Pecan spittlebug Pecan weevil San Jose scale Twig girdler Walnut caterpillar	7 1/2	<p>OBSERVE BEE CAUTION</p> <p>Repeat applications as necessary up to a total of 4 times per crop (including any applications at the dormant or delayed dormant timing) but not more often than once every 7 days.</p> <p>Use lower rates for pests attacking leaves. Use higher rates for pests attacking fruit and for higher infestations.</p> <p>For scale control, apply when crawlers are present.</p> <p>For chestnut weevil, best results have been found with 4 applications at weekly intervals beginning in late July. The last application should be made prior to shuck split.</p> <p>For peach twig borer, best results with foliar applications have been found by making applications in "popcorn" or petal fall stages when the May brood begins to hatch.</p> <p>For navel orangeworm in almonds and walnuts, best results have been found by timing early and midseason applications to correspond with moth flight peaks.</p> <p>For filbert leafroller, best results have been found by making applications when eggs are hatching, repeating application on first appearance of moths and again 3 to 4 weeks later.</p> <p>For codling moth in walnuts, best results have been found by making applications when average cross-sectional diameters of developing nuts are 0.5 to 0.75 inches and again during middle or late June as needed.</p> <p>For European earwig, thorough coverage of trunks, branches, and nuts is needed for best results.</p>

RESTRICTIONS AND PRECAUTIONS: TREE NUTS

- For trees taller than 10 feet, consider hiring a licensed professional.
- Do not apply within 14 days of harvest.
- Do not apply more than a total of 22 1/2 fluid ounces per 1000 SQ. FT. per crop, including any application at the dormant or delayed dormant timing.

17/25

DIRECTIONS FOR USE WITH A HOSE END APPLICATOR, CONTINUED

TURFGRASS

CROP	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
<p>Turfgrass: domestic lawns and recreational areas.</p>	<p>Ants Armyworm Centipedes Chiggers Cutworms Earwigs Essex skipper European chafer Fall armyworm Fiery skipper Grasshoppers Green June beetle larvae Leafhoppers</p> <p>Lucerne moth Millipedes Mosquitoes (adults) Scarab beetle adults (May beetle, June beetle, Japanese beetle, green June beetle) Sowbugs Spittlebugs Springtails Ticks Yellowstriped armyworm</p>	<p>6</p>	<p>For Armyworm, Cutworm, and Fall Armyworm Control: Do not irrigate treated areas for 24 hours following insecticide application.</p> <p>For Green June Beetle Grub (larvae) Control: Make applications when grubs are feeding near the soil surface.</p> <p>Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.</p>
	<p>Chinch bugs Sod Webworm (lawn moths) Bluegrass billbug larvae European crane fly Fleas White grubs (Japanese beetle, Chafer beetle, and <i>Phyllophaga</i> spp. larvae) Black turfgrass ataenius beetle larvae Hyperoides weevil (bluegrass weevil) larvae</p>	<p>12</p>	<p>For Chinch bug Control: Treat entire grass area rather than just damaged areas. Irrigation prior to treatment will aid in penetration of insecticide into turfgrass. Do not irrigate treated areas within 24 hours following insecticide application.</p> <p>For European Crane Fly Control: Treatments should be applied in early spring, April 1 to April 15, or as recommended by local Agricultural Extension Service agents.</p> <p>For White Grub Control: Applications should be made when grubs are feeding near the soil surface, usually during late March through May, or July through early September, or as recommended by local Agricultural Extension Service agents.</p> <p>Repeat applications as necessary up to 2 times per year but not more often than once every 7 days.</p>

19/25

DIRECTIONS FOR USE WITH A HOSE END APPLICATOR, CONTINUED

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES

CROP/SITE	PEST	FLUID OUNCES OF SEVIN® RP2 PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
<p>Ticks Which Vector Lyme Disease</p> <p>All crops/sites on this label</p>	<p><i>Ixodes</i> spp. (Deer tick, Bear tick, Black legged tick)</p> <p><i>Amblyomma</i> spp. (Lone star tick)</p>	<p>1 1/2 to 3 fl. oz. per 1000 SQ. FT.</p>	<p>Use the high rate for heavy tick infestations. Use higher spray volumes for dense ground cover or heavy leaf litter.</p> <p>Target applications for nymphal control in late spring or early summer. Control of adult tick can be obtained with late summer and fall applications.</p> <p>Do not use spot treatments. Treat entire area and perimeter areas where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary, but not more often than once every 7 days, to maintain adequate control levels.</p>
<p>Imported Fire Ants</p> <p>All crops/sites on this label</p>	<p>Imported fire ants</p>	<p>1 1/2 fl. oz. per mound.</p>	<p>MOUND DRENCH APPLICATION:</p> <p>Apply approximately 2 gallons of the hose-end spray solution over the surface of the mound. Thoroughly wet mound and surrounding areas to a 4 ft diameter (12 sq. ft.). Do not disturb mound prior to treatment. Direct flow at mound to disrupt and allow solution to flow into ant tunnels. For best results apply in cool weather (65-80°F) or in early morning or late evening hours. Repeat application if mound activity resumes after 7 days. Treat new mounds as they appear.</p>
<p>Adult Mosquito Control</p> <p>All crops/sites on this label</p>	<p>Mosquitoes (adults)</p>	<p>1 1/2 fl. oz. per 1000 SQ. FT.</p>	<p>OBSERVE BEE CAUTION.</p> <p>Treat shrubby and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early mornings or late evenings.</p> <p>Repeat applications as necessary but not more often than once every 7 days.</p>
<p>Nuisance Pest Control</p>	<p>Ants Bees Brown dog tick Centipedes Cockroaches Crickets Darkling beetle (lesser mealworm or litter beetle) Earwigs Firebrats Fleas Millipedes Scorpions Silverfish Spiders Wasps</p>	<p>12 fl. oz. per 1000 SQ. FT. as a perimeter treatment</p>	<p>PERIMETER TREATMENT DIRECTIONS:</p> <p>Observe plant response precautions.</p> <p>Apply as a coarse wet spray in a band 6 to 10 feet wide around the outside perimeter of buildings. Confine applications to areas immediately adjacent to the building. Direct application to the structure should be minimal and restricted to cracks, crevices, and areas where insects tend to congregate.</p> <p>For use in and around flower beds and ornamental plantings, apply as a coarse wet spray in and around plant beds.</p> <p>May be applied to outdoor sleeping quarters of pets. Do not treat animals. Do not allow animals to use treated area until sprays have dried. Do not treat more often than once every 7 days.</p>

RESTRICTIONS AND PRECAUTIONS:

- DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.
- NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits on painted or stained surfaces of finishes (i.e., cars, house trailers, boats, etc.) should be removed immediately by washing to prevent discoloration. Avoid application to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

CAUTION: May kill shrimp and crabs. Do not use in areas where these are important resources.

20/25

SPECIFIC USE DIRECTIONS FOR A SINGLE DOSE PACKAGE

Mix the entire contents of this packet (0.67 ounces) with one gallon of water. For safety and performance reasons, this mixed product should not be stored once opened.

VEGETABLES

For control of all the vegetable pests listed below, apply this product to effectively cover both the upper and lower surfaces of the plant to the point of runoff.

Repeat applications as necessary up to a total of 4 times for Legume vegetables, Brassica Leafy vegetables; 5 times for asparagus and leafy vegetables; 6 times for cucurbit vegetables, root and tuber vegetables; 7 times for fruiting vegetables; and 8 times for sweet corn but not more often than once every 7 days.

NOTE: This product is not registered for use on celery and sweet potatoes in California.

Do not treat asparagus more than once every 7 days as excessive residues may result.

The number in parenthesis following each vegetable grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

VEGETABLES	PESTS
Beans (including Black-eyed peas, Cowpeas, Crowder or Southern peas, Dry beans, Green beans, Lima beans, Navy beans, and Snap beans) (3 days until harvest for fresh beans and peas, 21 days until harvest for dried beans and peas)	Alfalfa caterpillar Lygus bug
Broccoli, Brussels sprouts, Cabbage, Cauliflower, Cucumber, Eggplant, Kohlrabi, Melons, Pepper, Tomato (3 days until harvest)	Apache cicada Melonworm
Carrots, Potato (7 days until harvest)	Armyworms Mexican bean beetle
Corn (2 days until harvest)	Asparagus beetle Pea leaf weevil
Asparagus (1 day until harvest)	Aster leafhopper Pea weevil
Garden beet roots, Horseradish, Parsnips, Radishes, Rutabagas, Salsify roots, Turnip roots (7 days until harvest)	Bean leaf beetle Rednecked peanutworm
Head lettuce, Chinese cabbage, Celery, Collards, Dandelions, Endive (Escarole), Garden beet tops, Hanover salads, Kale, Mustard greens, Leaf lettuce, Parsley, Salsify tops, Spinach, Swiss chard, Turnip tops (14 days until harvest)	Blister beetles Sap beetles
	Chinch bug Southwestern corn borer
	Colorado potato beetle Spittlebugs
	Corn earworm Squash bugs
	Corn rootworm adult Stink bugs
	Cowpea curculio Sweet potato weevil
	Cucumber beetle Tarnished plant bug
	Cutworms Three cornered alfalfa hopper
	European corn borer Thrips
	Fall Armyworm Tomato hornworm
	Flea beetles Tomato pinworm
	Grasshoppers Tortoise beetle
	Green cloverworm Velvetbean caterpillar
	Harlequin bug Webworms
	Imported cabbageworm Western bean cutworm
	Japanese beetle Whitefringed beetle adults
	Lace bugs Yellowstriped armyworm
	Leafhoppers
	Lima bean pod borer

SPECIFIC USE DIRECTIONS FOR A SINGLE DOSE PACKAGE (continued)

FRUITS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward upper and lower leaf surfaces and between fruit clusters, and on small limbs and trunks to the point of runoff where pests appear. Repeat applications as necessary up to a total of 8 times for tree fruits and 5 times for small fruits but not more often than once every 7 days.

NOTE: To avoid undesired apple thinning, delay use until at least 30 days after full bloom. Carbaryl insecticide may injure early dawn and sunrise strawberry varieties on the Delmarva Peninsula. The number in parenthesis following each fruit and nut grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

TREE FRUITS	SMALL FRUITS	PESTS	
Apples, Cherries, Peaches, Pears, Plums, Prunes (3 days until harvest)	Blueberries, Grapes (7 days until harvest)	Apple aphid	Lesser appleworm
Apricots, Nectarines (3 days until harvest)	Strawberries (7 days until harvest)	Apple maggot	Lesser peachtree borer
Citrus (5 days until harvest)	Blackberries, Raspberries, Dewberries (7 days until harvest)	Apple mealybug	Meadow spittlebug
		Apple pandemis	Mealy plum aphid
		Apple rust mite	Omnivorous leaftier (Strawberry fruitworm)
		Apple sucker	Omnivorous Leafroller
		Avocado leafroller	Orange tortrix
		Bagworms	Oriental fruit moth
		Black cherry aphid	Oystershell scale
		Black scale	Peach twig borer
		Blueberry maggot	Pear leaf blister mite
		Brown soft scale	Pear psyllia
		California red scale	Pear rust mite
		Cherry Fruitworm	Periodical cicada
		Cherry maggot	Plum curculio
		Citricola scale	Prune leafhopper
		Citrus cutworm	Raspberry sawfly
		Citrus root weevil	Red-banded leafroller
		Citrus snow scale	Rose chafer
		Codling moth	Rosy apple aphid
		Cucumber beetles	Saltmarsh caterpillar
		Cutworms	San Jose scale
		Eastern tent caterpillar	Snowy tree cricket
		Eightspotted forester	Strawberry weevil
		European apple sawfly	Tarnished plant bug
		European earwig	Tentiform leafminers
		European raspberry aphid	Tussock moth
		Eyespotted budmoth	Variiegated leafroller
		Forbes scale	Western grapeleaf skeletonizer
		Fruitree leafroller	Western tussock moth
		Grape berry moth	Western yellowstriped armyworm
		Grape leafhopper	West indian sugarcane borer (adults)
		Grape leaffolder	White apple leafhopper
		Green fruitworm	Woolly apple aphid
		Japanese beetles	Yellow headed fireworm
		June beetles	
		Leafhoppers	
		Leafrollers	
		Lecanium scale	

RESTRICTIONS AND PRECAUTIONS: FRUITS

- For trees taller than 10 feet, consider hiring a licensed professional.

22/25

SPECIFIC USE DIRECTIONS FOR A SINGLE DOSE PACKAGE (continued)

TREES, ORNAMENTALS, SHRUBS AND FLOWERS

This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward the upper and lower leaf surfaces and small trunks, stems and twigs to the point of runoff where pests appear. Repeat applications as necessary up to a total of 4 times for trees and 6 times for ornamentals, shrubs and flowers but not more often than once every 7 days.

TREES	PESTS		
Trees including: Shade trees, Shelter belts, Plantations, Parks and Recreational Areas	Ants	Hackberry nipplegall maker	Poinsettia hornworm
Ornamentals including: Roses, Other Herbaceous plants, Woody plants, Shrubs	Apple aphid	Holly budmoth	Psyllids
	Armyworm	Holly leafminer	Puss caterpillar
	Azalea leafminer	Jackpine budworm	Redhumped oakworm
	Bagworms	Japanese beetles	Rose aphid
	Birch leafminer	Jeffrey pine needleminer	Rose chafer
	Blister beetle	June beetles	Roseslug
	Boxelder bug	Plant bug	Saddled prominent
	Boxwood leafminer	Lace bugs	Sawflies (exposed)
	Browntail moth	Leafhoppers	Scale insects
	Cankerworms	Leafrollers	Sowbugs
	Catalpa sphinx	Locust borer	Spiny elm caterpillar
	Chiggers	Maple leafcutter	Springtails
	Cooley spruce gall aphid	Mealybugs	Spruce budworm
	Elm leaf aphid	Mimosa webworm	Spruce needleminer
	Elm leaf beetle	Nantucket pine tip moth	Subtropical pine tip moth
	Elm spanworm	Oak leafminers	Tent caterpillar
	Eriophyd mites	Oak leaf skeletonizer	Thorn bug
	European pine shoot moth	Oakworm complex	Thrips (exposed)
	Fall armyworm	Oleander caterpillar	Ticks
	Flea beetles	Olive ash borer	Walnut caterpillar
	Fulcrum rose beetles	Orange-striped oakworm	Webworms
	Gall midges	Orange tortrix	Western hemlock looper
	Gall wasps	Periodical cicada	Western spruce budworm
	Grasshoppers	Pine sawfly	Willow leaf beetles
	Green striped mapleworm	Pine spittlebug	Yellow poplar weevil
	Gypsy moth	Pitch pine tip moth	

RESTRICTIONS AND PRECAUTIONS: TREES, ORNAMENTALS, SHRUBS AND FLOWERS

- For trees taller than 10 feet, consider hiring a licensed professional

SPECIFIC USE DIRECTIONS FOR A SINGLE DOSE PACKAGE (continued)**PEST CONTROL AROUND BUILDINGS AND OTHER OUTDOOR AREAS**

This product may be used as a spot treatment around buildings such as homes, apartments, warehouses, barns and municipal and recreational areas to control the pests listed below. Thoroughly wet the outside perimeter of dwellings and other areas where pests tend to congregate. Do not apply more often than once every 7 days.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of this product to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wallpaper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

PESTS	
Ants	Earwigs
Bees	Fireboats
Wasps	Silverfish
Brown dog tick	Fleas
Carpenter ants	Millipedes
Centipedes	Scorpions
Cockroaches	Spiders
Crickets	

24/25

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.


By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

SEVIN is a registered trademark of Bayer.



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscience.us>

SEVIN RP2 Carbaryl Insecticide (MASTER) Approved 4/16/03, Notification 10/25/05

Bayer CropScience



25/05

November 14, 2005

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
1801 South Bell Street
Arlington, Virginia 22202-4501
(PM: Mr. John Hebert, Team 7)

Dear Reviewer,

Re: **SEVIN® brand RP2 Carbaryl Insecticide**
EPA Reg. No. 264-334
Notification of Minor Labeling Changes per PR Notice 98-10

In accordance with PR Notice 98-10, we are notifying the Agency of minor labeling amendments for SEVIN® brand RP2 Carbaryl Insecticide. Instructions on the use of hose end sprayers for application of the product were added on page 9 under "DIRECTIONS FOR USE WITH HOSE END APPLICATOR".

The following documents are enclosed in support of this action:

- A completed Application for Registration (EPA Form 8570-1) which includes the Certification Statement required by PR Notice 98-10.
- One (1) copy of the label with changes highlighted (page 9).

Please call me at 919/549-2718 if you need any additional information.

Sincerely,

Danielle A. Larochelle
Registration Product Manager

Attachments

Corr. # daL065-05