

ME 4™ BROMINAL® Bromoxynil
Broadleaf Herbicide

264-340

CROP USE

Controls annual broadleaf weeds in wheat, barley, oat, rye, flax, garlic, onions and newly seeded and established grasses.

INDUSTRIAL USE

Controls Russian thistle and other annual broadleaf weeds on roadsides, vacant lots, dry drainage beds, railroads and industrial sites.

For Commercial or Agricultural Use Only.

ACTIVE INGREDIENTS:

Octanoic acid, 3,5-dibromo-4-hydroxybenzotrile ester*	31.5%
Butyric acid, 3,5-dibromo-4-hydroxybenzotrile ester*	22.1%
INERT INGREDIENTS:	46.4%

*Total Bromoxynil (3,5-Dibromo-4-hydroxybenzotrile) equivalent 39.4% by weight or 4 pounds per gallon

E.P.A. Reg. No. 264-340 E.P.A. Est. 264-MO-1

U.S. PATENTS No. 4,332,613

WARNING: Keep out of reach of children.

PRECAUTIONARY STATEMENTS: See inside booklet.

IN CASE OF EMERGENCY TELEPHONE COLLECT
(24 HOURS A DAY) IN U.S.A. (304) 744-3487

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.

P. O. BOX 12014, T.W. Alexander Drive, Research Triangle Park, N.C. 27709

ME 4™ BROMINAL® and UNION CARBIDE are registered trademarks of UNION CARBIDE CORPORATION. MADE IN U.S.A.

CONTENTS 2.5 GAL. (9.46 Liters)
DO NOT DETACH BOOKLET FROM CONTAINER

TABLE OF CONTENTS

	PAGE
Accidental Ingestion or Contact	17
Application Timing	3-13
Bromigation Application through Irrigation Systems	13
Compatibility	3
Crops	
Barley	4-10
Flax	11
Garlic	12
Oat	4-10
Onions	12
Rye	4-10
Wheat	4-10
General Information	2-3
Grasses	
Established lawn and turf areas	14
Newly planted	14
Limited Warranty and Disclaimer	18
Precautionary Statements	17-18
Storage and Disposal	18
Tank Mixtures ME 4™ BROMINAL®	
2,4-D	5
Avenge (difenzoquat)	8
Banvel (dicamba)	6
Carbyne (barban)	9
Fertilizer	3
Hoelon (diclofop)	10
Karmex (diuron)	7
MCPA	5
Other pesticides and herbicides	3
Use Directions	
Crops	4-12
Industrial	17
Weeds Susceptible	15-16

GENERAL INFORMATION CROP USE

ME 4^x BROMINAL^s bromoxynil broadleaf herbicide effectively controls broadleaf weeds germinating from seed, applied when these weeds are in the seedling stage.

ME 4^x BROMINAL^s applied early kills seedling broadleaf weeds that emerge in small grains, onion, garlic and flax, thus reducing weed competition with these crops during their early growth.

ME 4^x BROMINAL^s broadleaf herbicide may be applied in the fall, winter or spring of the year depending on the area. Spraying must be done while weeds are in the seedling stage. Application of ME 4^x BROMINAL^s is not registered for use on small grains undersown with legumes.

Apply 0.5 to 1 pint of ME 4^x BROMINAL^s broadleaf herbicide per acre after small grains emerge, but before weeds are past the 3 to 4 leaf stage. Weeds that form rosettes should be treated when these weeds are less than 1-1/2 inches across.

Good spray coverage is essential for maximum weed control. Use 10 or more gallons of water per acre with ground sprayers. Nozzle pressure should be at least 30 psi. Use 2 or more gallons of water per acre with aerial sprayers according to the direction use tables. Width of swath must not exceed length of spray boom.

The most important factors for effective weed control are : 1. spraying weeds in the seedling stage; 2. Uniform spray coverage.

Do not apply when crops form a canopy over weeds. Do not apply when crops are under moisture stress. Do not apply to wheat, barley, or rye that will be exposed to cold desiccating winds in the absence of snow cover. Do not apply to small grains during or after the boot stage. If high crop mortality from winds has occurred, apply ME 4^x BROMINAL^s only after plants have sent out new roots and begin new top growth. Do not graze treated fields for 30 days after application.

ME 4^x BROMINAL^s broadleaf herbicide may be applied in liquid fertilizer or combinations of liquid fertilizer and water. Use constant agitation until the spray mixture has been completely applied.

Tank mixes permit choosing a herbicide combination most effective for the weed problems present on your farm. Use ME 4^x BROMINAL^s broadleaf herbicide alone or in combination according to the label. Follow all cautions and limitations on the labels of the products you select to mix with ME 4^x BROMINAL^s.

If tank mixtures other than those listed on this label are chosen, check compatibility by mixing the components of the tank mixtures in the same proportions and dilutions planned for the application. Do not combine ME 4^x BROMINAL^s/2,4-D or MCPA with Hoelon as the 2,4-D or MCPA will interfere with the control of grass by Hoelon.

ME 4^x BROMINAL^s contains 4 lb active bromoxynil per gallon; 0.5 pint contains 0.25 lb active ingredient; 0.75 pint contains 0.375 lb active ingredient; 1.0 pint contains 0.5 lb active ingredient.

DIRECTIONS FOR USE

DO NOT USE THIS PRODUCT FOR PURPOSES OTHER THAN THOSE RECOMMENDED ON THIS LABEL. IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

ME 4 BROMINAL and Wheat, Barley, Oats and Rye

Product/Area	Rate	Crop	Application Timing	Weeds	Spray Volume
Cereal Areas except WA, OR, ID					
ME 4 BROMINAL	1/2 - 1 (0.5-1) pint	Winter cereal: Fall through early spring applications before crop has reached the boot stage (not underseeded with legumes).		Apply to emerged susceptible weeds in the 2-4 leaf stage or when rosettes are less than 1.5 inches across	Use 10 to 20 gal. with ground or 5-10 gal. with air application. Use higher volumes for larger weeds or when growing conditions are poor.
		Spring Cereal: One leaf to boot stage.			Use 2 or more gal. with mid-west air application.
Washington, Oregon, Idaho					
BROMINAL ME 4	3/4 - 1 (0.75-1) pint			Crowweed, Tarweed, Fiddleneck, mustard, other broadleaf weeds apply before the 4 leaf stage, before rosettes are 1.5 inches across.	Use 10 to 20 gal. with ground or 5-10 gal. with air application. Use higher rates and volumes for larger weeds or where weed growth is slow due to dry or cold weather.

ME 4 BROMIRAL and MCPA - 2,4-D Tank Mixture on Wheat, Barley, Oat and Flax.

When using ME 4 BROMIRAL in combination with 2,4-D or MCPA, first mix the 2,4-D or MCPA in the water, then add the ME 4 BROMIRAL. Use the higher rate and volume for larger weeds or where weed growth is slow due to dry or cold weather. Do not include Boston in this tank mixture, as it will reduce the effectiveness of Boston.

Product/Area	Rate	Crop	Application Timing	Weeds	Spray Volume
Cereal Areas, except Washington, Oregon, Idaho					
ME 4 BROMIRAL + 2,4-D* or MCPA*	1 1/2 - 3/4 (0.5-0.75) pint + 1/4 - 1/2 (0.25- 0.5) lb. active		Application after tillering and before the forming of joints in the stem.	All weeds including wild buck- wheat, cow cockle, fiddleneck, tarweed; apply at 1 - 4 leaf stage but before 6 inches high.	Use 10-20 gal. with ground or 4-10 gal. water with air appli- cation. Use higher volume on larger weeds

Washington, Oregon, Idaho					
ME 4 BROMIRAL + 2,4-D* or MCPA*	3/4 - 1 (0.75-1) pint + 1/4 - 1/2 (0.25- 0.5) lb. active			Gromwell, tarweed, fiddle- neck, mustard, other broad- leaf weeds; apply before weeds are 6 inches high.	Use 10-20 gal. with ground or 5-10 gal. water with air application. Use higher volume on larger weeds

*2,4-D amine - WEEDAR 64 Herbicide, *2,4-D esters - WEEDONE 1V 4 Herbicide, *MCPA amine - WEEDAR MCPA concentrate, *MCPA Ester WEEDONE MCPA Ester Broadleaf Herbicide.

ME 4 BROMINAL and Banvel Tank Mixtures on Wheat

When Banvel is tank mixed with ME 4 BROMINAL, water should be added to the spray tank first, followed by ME 4 BROMINAL and Banvel.

Product/Area	Rate	Crop	Application Timing Weeds	Spray Volume
Throughout the United States				
ME 4 BROMINAL + Banvel (dicamba)	1/2-3/4 (0.5-0.75) pint + 1/16-1/8 (0.0625-0.125) pint	Spring Wheat: Apply when spring wheat is from 3-leaf up to joint stage.	Bachelor's button, knotweed, mustard, nightshade, Russian thistle, wild buckwheat, lambsquarters, lamweed, dogfennel, cow cockle; apply when weeds are small (early seedling).	Use 10 or more gal. of water per acre for ground application
Throughout the United States				
ME 4 BROMINAL + Banvel (dicamba)	1/2-3/4 (0.5-0.75) pint + 1/8 (0.125) pint	Winter Wheat: Fall through early spring application when winter wheat is from 3-leaf up to joint stage.	Foxtail, fiddleneck, miners-lettuce, knotweed, wild buckwheat, prickly lettuce, jacobsladder, alkanet, burchevil, corn cockle, corn groundwell, cow cockle, dogfennel, lamweed, falseflax, flyweed, henbit, lambsquarters, catchweed; apply when weeds are small (early seedling).	

When treating with Banvel, do not graze treated areas or harvest for dairy feed prior to crop maturity.

Banvel is a registered trademark of Velsicol Chemical Corporation.

ME 4 BROMINAL and Karnez Tank Mixture on Wheat and Barley

When Karnez is tank mixed with BROMINAL for herbicide control of weeds, use the following information on the Karnez label and especially those regarding winter stress, varieties, soil texture, organic matter, planting depth, plant vigor and replanting. Use where average annual rainfall exceeds 16 inches on winter wheat and winter barley.

Product/Area	Rate	Crop	Application Timing Weeds	Spray Volume
Washington, Oregon, Idaho				
ME 4 BROMINAL + Karnez (duron)	1/2 (0.5) pint + 1/2 (0.5) lb.	Winter wheat and winter barley.	Herbicide, quackgrass, fiddleneck, other weeds. Apply before weeds are 2 inches tall or across.	Use a minimum of 25 gal. with ground or 5-10 gal. with air application for thorough cover- age of weed foliage.
		Apply a single application in the fall after wheat has emerged, but before soil freezes or in the spring as soon as soil thaws.		

Karnez is a registered trademark of E. I. DuPont de Nemours and Company.

ME 4 BROMINAL and Carbyne (barban) Tank Mixture for Broadleaf and Wild Oats Control in Wheat and Barley

When applying a tank mix of ME 4 BROMINAL and Carbyne follow the directions for spray nozzle arrangements on the Carbyne label. Maintain vigorous agitation in the spray tank until spraying is completed. Sequential treatments have been most effective.

Do not spray this tank mixture when plants are wet with heavy dew or rain.

Do not let spray drift into adjacent fields, particularly if planted with oats, rye or buckwheat.

Do not allow livestock to graze on fields sprayed with the Carbyne 21C tank mixed with ME 4 BROMINAL until after harvest of the crop.

Product/Area	Rate	Crop	Application Timing Weeds	Spray Volume
Throughout the United States. Sequential Carbyne treatment.				
ME 4 BROMINAL + Carbyne 21C 1st application	1/2-3/4 (0.5-0.75) pint + 1-1 1/2 (1-1.5) pint	Winter type wheat and spring type wheat and barley.	Broadleaf weeds 2-4 leaves. Wild oat, canary grass in 2 leaf stage. Apply the BROMINAL tank mix only once if the split Carbyne application is used.	Use 5-10 gal. of water and sufficient pressure (minimum 45 lb.) to break spray into small droplets with ground equipment and 3-5 gal. for air application.
+ Carbyne 21C 2nd application	1-1 1/2 (1-1.5) pint	ME 4 BROMINAL can be combined with 1st or 2nd Carbyne application.	Use the high rate of Carbyne when weed population is over 50 plants per sq. ft.	
ME 4 BROMINAL + Carbyne 21C	1/2-3/4 (0.5-0.75) pint + 1-1 1/2 (1-1.5) pint	Winter type wheat and spring type wheat and barley.	Broadleaf weeds 2-4 leaves. Wild oat, canary grass in 2 leaf stage. Use the high rate of Carbyne when weed population is over 50 plants per sq. ft.	
ME 4 BROMINAL + Carbyne 21C	1/2-3/4 (0.5-0.75) pint + 2 pint	Winter type wheat and spring type wheat and barley except durum wheat.	Broadleaf weeds 2-4 leaves. Wild oat, canary grass in 2.5 - 3.5 leaf stage. do not spray when wild oats or canary grass have 4 or more leaves.	

Carbyne is a registered trademark of Velsicol Chemical Corporation.

ME 4 BROMINAL and Hoeion Tank Mixture for Broadleaf and Wild Oat and Foxtail (pigeon grass) in Wheat and Barley

When using the ME 4 BROMINAL and Hoeion tank mixture, cold (lower than 40 °F) and/or prolonged wet weather conditions increase the sensitivity of barley to Hoeion and Hoeion should not be used under these adverse weather conditions. Use the low rate of Hoeion when weather conditions are cool and moist. Do not use 3 + 3 BROMINAL (Bromoxynil and MCPA) with Hoeion as it will reduce the effectiveness of Hoeion.

Product/Area	Rate	Crop	Application Timing Weeds	Spray Volume
Throughout the United States				
ME 4 BROMINAL + Hoeion	3/4-1 (0.75-1) pint + 2 2/3 (2.66) pint	Winter type wheat and spring type wheat and barley before or after tillering.	Grasses, wild oat, foxtails, (pigeon grass) annual eye-grass, barnyardgrass, 1-3 leaves, Tiddleneck and other broadleaf weeds 2-4 leaves. Use 1 pint for 3-4 leaf broad leaf weeds.	Use at least 10 gal. with minimum 30 psi for ground or at least 5 gal with air application.
ME 4 BROMINAL + Hoeion	3/4-1 (0.75-1) pint + 2 2/3 - 3 1/3 (2.66-3.33) pint	Winter and spring type wheat before or after tillering.	Grasses, 3-4 leaves Broad-leaves, 3-4 leaves.	

Hoeion is a registered trademark of American Hoechst Corporation. Refer to the Hoeion label for list of states where applicable.

ME 4 BROMIRAL - Broadleaf Weed Control in Flax

ME 4 BROMIRAL can injure Flax if applied when the humidity is high. Temperatures at the time of treatment, or for the three days after treatment, must be below 85° F. to avoid serious Flax injury.

Product/Area	Rate	Crop	Application - Target Weeds	Spray Volume
Throughout the United States				
ME 4 BROMIRAL	1/2-1 (0.5-1) pint	Flax	Weeds 2-4 leaf stage.	Use 10-20 gal./Acre with ground or 5-10 gal./Acre with air application.

GARLIC - DIRECTIONS FOR USE

For broadleaf weed control in garlic apply 1 to 2 pints of ME 4^v BROMINAL^s per broadcast acre when the weeds are 1 to 4 inches tall. Use the lower rate on small weeds and the higher rate on 4 inch weeds. The garlic will generally have 1 to 2 true leaves and treatment should be completed before the garlic is 12 inches tall.

For ground application use 40 to 100 gallons of spray solution per broadcast acre. Good coverage of weeds is essential for good control and the higher volumes are preferred for complete coverage. For aerial application use 5 gallons of spray solution per acre.

Do not apply within 112 days of anticipated harvest.

ONIONS - DIRECTIONS FOR USE

Apply 0.5 to 0.75 pints (0.25 to 0.38 lbs active ingredient) of ME 4^v BROMINAL^s per acre on a broadcast basis. Water volume is important--use 50-70 gallons of water/acre and apply when onions have 2-5 true leaves. Concentrate spray kills onions. Good coverage is necessary for good weed control. Do not treat if thrip or sand damage have occurred and removed or reduced the waxy coating on the onion foliage. Some crop damage may occur even under ideal conditions. Do not add surfactants.

In onion-producing areas environmental conditions that reduce development of the waxy coating on the onion leaf increase the possibility of injury. Dry soil, dry onion foliage, high light intensity, low humidity, and high temperatures tend to increase the waxy coating on the onion leaf and thus reduce chance for injury. It is essential for greatest safety that the soil and onion foliage be dry. Humidity should be low and dew should be off the plants.

Applications should not be made 24 hours before or after cultivation, irrigation, or rain; longer than 24 hours if possible. It is best to spray on a bright, sunny day after dew has completely dried from onion foliage and before it has formed.

Do not use ME 4^v BROMINAL^s on onions grown under light intensity in areas such as Oregon, west of the Cascades.

APPLICATION OF ME 4* BROMINAL* THROUGH IRRIGATION SYSTEMS - BROMIGATION
- BROADLEAF WEED CONTROL SERVICE

General Information: ME 4* BROMINAL* bromoxynil broadleaf herbicide may be applied through automated (center pivot or continuous lateral move) or hand move and wheel line irrigation equipment for broadleaf weed control. Properly applied ME 4* BROMINAL* will control fiddleneck, wild buckwheat, sunflower, fanweed, Kochia, Russian thistle and other susceptible broadleaf weeds as listed on the ME 4* BROMINAL* label. For most efficient use of ME 4* BROMINAL* apply when weeds are small, before they have exerted yield reducing competition on the crop.

Directions for use: Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water. Add ME 4* BROMINAL* broadleaf herbicide or ME 4* BROMINAL* and water dilution to mix tank. For greater accuracy in calibration and distribution, inject a larger quantity of a more dilute solution per unit time.

For best performance set sprinkler to deliver approximately 0.5 inch or less of water per acre. Start sprinkler and then uniformly inject ME 4* BROMINAL* into the irrigation line so as to deliver a rate ME 4* BROMINAL* broadleaf herbicide per acre as suggested under the crop section of this label. Weeds should not be larger than 3-4 leaf growth stage. In cereal, flax, onions and garlic use the 1 pint rate on weeds with 4 leaves. In cereal, flax, and garlic use the 1 pint rate if pigweed or lambsquarters are part of the weed complex. Refer to ME 4* BROMINAL* label for detailed information on application timing. ME 4* BROMINAL* broadleaf herbicide should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to insure adequate mixing.

User precautions: Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils.

Apply ME 4* BROMINAL* only through systems containing anti-siphon, check valves and interlocking controls between the metering device and the water pump to insure simultaneous shutoff to prevent water source contamination and overflow of the mixing tank.

Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

GRASSES - DIRECTIONS FOR USE

NEWLY PLANTED GRASSES

Apply 0.75 pints of ME 4* BROMINAL* broadleaf herbicide per acre after grasses emerge and before broadleaf weeds are past the 3 to 4 leaf stage

Use 1 pints of ME 4* BROMINAL* per acre if weeds are beyond the 3 to 4 leaf stage but less than 6 inches in height and before flower formation. Weeds that form rosettes should be treated when these rosettes are less than 1 1/2 inches across.

This treatment may be used on seedling grasses such as Merion, Park, Delta or common Kentucky bluegrasses, Pennlawn, Chewings, Illahee or Alta fescues, orchard grass, Highland, Seaside or Astoria bentgrasses, and perennial ryegrasses.

Established Lawns and Turf Areas

For seeding weeds such as soliva sp., mustard, lambsquarter and pigweed in established lawns (2-3 months after seed germination) apply 1 pint per acre (3.67 fl. oz. for 10,000 sq. ft). If weeds are at the 3 - 4 leaf stage, a repeat treatment will be needed in 10-14 days for control or suppression. To control or suppress hard to kill weeds such as Black Medic, Bur Clover and Spiny Fruited Crowfoot, apply 4 pints per acre of ME 4* BROMINAL® (7.35 fl. oz. for 10,000 sq. ft).

Mix in enough water for uniform coverage of all weeds. For power sprayers equipped with booms, use at least 3 gallons of water per 2,500 square feet, or 50 gallons per acre.

Temporary turfgrass discoloration may occur particularly if excessive overlapping of spray solution occurs, or if ME 4* BROMINAL® is applied at temperatures of 85° F or above. This will usually disappear following two mowings leaving no permanent damage.

DO NOT FEED TURF GRASSES TREATED WITH ME 4* BROMINAL® TO LIVESTOCK.

WEEDS SUSCEPTIBILITY IN SEEDLING STAGE

Most Susceptible Weeds Use 1/2 (0.5) pint ME 4* BROMINAL® 0.25 lb active ingredient per acre:

Black mustard	(Brassica nigra)
Bristly star bur	(Acanthospermum hispidum)
Cocklebur	(Xanthium pennsylvanicum)
Field pennycress, farweed	(Thlaspi arvense)
Green smartweed	(Polygonum scabrum)
Kochia	(Kochia scoparia)
Lambsquarters	(Chenopodium album)
London rocket	(Sisymbrium irio)
Morningglory (annual)	(Ipomoea hederacea)
Nettleleaf goosefoot	(Chenopodium murale)
Nightshade	(Solanum nigrum, villosum)
Parthenium ragweed	(Parthenium hysterophorus)
Ragweed	(Ambrosia artemisiifolia)
Shepherdspurse	(Capsella bursa-pastoris)
Silver leaf nightshade	(Solanum elaeagnifolium)
Sowthistle (annual)	(Sonchus oleraceus)
Sunflower	(Helianthus annuus)
Treacle mustard	(Erysimum repandum)
Tumble (Jim Hill) mustard	(Sisymbrium altissimum)
Velvetleaf	(Abutilon theophrasti)
Wild buckwheat	(Polygonum convolvulus)
Wild radish	(Raphanus raphanistrum)

Most Susceptible Weeds Use 3/4 (0.75) pint of ME 4* BROMINAL:

Blue (purple) mustard	(Chorispora tenella)
Bullthistle	(Cirsium vulgare)
Corn chamomile	(Anthemis arvensis)
Corn groundsel	(Lithospermum arvense)
Cow cockle	(Saponaria vaccaria)
Dyers woad	(Isatis tinctoria)
Fiddleneck	(Amsinckia intermedia)
Fivehooked bassia	(Bassia hyssopifolia)
Jimsonweed	(Datura stramonium)
Knave) (German moss)	(Scleranthus annuus)
Mare's-tail	(Coryza canadensis)
Nodding thistle	(Carduus nutans)
Pennsylvania smartweed	(Polygonum pennsylvanicum)
Popcorn flower	(Alloccarya figurata)
Prickly (Chinese) lettuce	(Lactuca scariola)
Puncture vine	(Tribulus terrestris)
Redmaids rockpurslane	(Colandrinia caulescens)
Smallseeded false flax	(Canelina microcarpa)
Southern giant curl mustard	(Brassica japonica)
Tartary buckwheat	(Fagopyrum tataricum)
Tarweed	(hemizonia spp.)
Wild mustard	(Brassica kaber)
Winter vetch	(Vicia sativa)
Yellow rocket	(Barbarea vulgaris)
Yellow star thistle	(Centaurea solstitialis)

Less Susceptible Weeds Use 1.0 pint ME 4* BROMINAL 0.5 lb active ingredient per acre:

Bur buttercup	(Ranunculus testiculatus)
Dog fennel (Mayweed)	(Anthemis cotula)
Fumitory	(Fumaria officinalis)
Groundsel	(Senecia vulgaris)
Henbit	(Lamium amplexicaule)
Miners lettuce	(Montia perfoliata)
Pepperweed	(Lepidium campestre)
Pigweed	(Amaranthus retroflexus)
Prickly (Chinese) lettuce	(Lactuca scariola)
Tansy mustard	(Descurainia pinnata)

INDUSTRIAL USE

Russian Thistle and Annual Broadleaf weeds on roadsides, vacant lots, dry drainage beds, railroads and industrial sites.

Directions for Industrial Use

ME 4^w BROMINAL^s broadleaf herbicide effectively controls Russian thistle and certain other summer annual broadleaf weeds such as saltbrush (Atriplex), mare's-tail, puncture vine, prostrate scourge, and Bassia.

ME 4^w BROMINAL^s should be applied in early summer when Russian thistle is young and growing vigorously.

For broadcast treatment of small weeds: Apply 1 quart of ME 4^w BROMINAL^s broadleaf herbicide with 2 quarts of surfactant such as Surfel^s Spray Adjuvant in 150 gallons of water for each acre sprayed. Thorough coverage is important. Make sure spray boom is high enough to contact tops of all plants. Use adequate spray pressure to contact all leaf surfaces.

For spot treatments of scattered infestations or large weeds along fencerows or buildings: Mix 0.5 quart Me 4^w BROMINAL^s with 1 quart of surfactant such as Surfel per 100 gallons of spray solution. Cover all the weed foliage thoroughly. Use at least 200 gallons of spray solution for each acre sprayed. Best industrial results with the ME 4^w BROMINAL^s and surfactant mixture are obtained where temperatures are 85° F or higher.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful if swallowed. Avoid contact with skin, eyes, clothing. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, rinse thoroughly with water and get medical attention. In case of skin contact, wash immediately with soap and water and remove contaminated clothing.

If swallowed - drink promptly large quantities of milk, egg white, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of waste.

Avoid spray drift onto neighboring crops or pastures. Clean sprayer thoroughly before using for other purposes.

STORAGE AND DISPOSAL

Freezing does not affect the quality of this product. Do not store near fertilizers, seed, insecticides or fungicides. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL

Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon reasonable experts evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

This warranty does not extend to, and the Buyer shall be solely responsible for, any, and all loss or damage which results from the use of this product (a) in any manner which is inconsistent with the label directions, warnings or cautions.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OR PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

Weedone® LV 4, Weedone® MCPA Ester, Weedone® 64, Weedone® MCPA Concentrate, ME 4™ BROMINAL® and Sunfel® Spray Adjuvant are registered trade marks and BROMIGATION® is a service mark of Union Carbide Agricultural Product Company.