EPA Form 8570-1 (Rev. 12-90)

Previous editions are obsolete.

White - EPA File Copy (original)

Yellow - Applicant copy

LARVIN® brandThiodicarb Insecticide 75WP

WETTABLE POWDER

ACTIVE INGREDIENT

EPA Registration No. 264-341

EPA Establishment No. 3125-M0-01

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

LA USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER): If you cannot read English, do not use this product until the label has been fully explained to you.)

For PRODUCT USE Information Call 1-800-334-9745.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never give anything by mouth to an unconscious or convulsing person. Call a physician and follow General advice listed below.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Call a physician and follow General advice listed below

GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway, start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

NOTE TO PHYSICIAN

LARVIN® thiodicarb is an oxime carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number: 1-800-334-7527.

PRECAUTIONARY STATEMENTS

WARNING HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be harmful or fatal if swallowed or inhaled. May be harmful if absorbed through the skin. May be irritating to eyes and may bause mild skin sensitization. Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing spray grist

Wear long-sleeved clothing while using. Remove contaminated clothing daily; wash before reuse. Wash exposed skin thoroughly with soap and water after use and before eating or smoking. Discontinue use if allergic reaction occurs.

SIGNS AND SYMPTOMS OF OVEREXPOSURE:

Salivation, Watery eyes, Pinpoint eye pupils, Blurred vision, Muscle tremors, Difficult breathing, Excessive sweating, Abdominal cramps, Nausea, Vomiting, Diarrhea, Weakness, Headache. In severe cases, convulsions, unconsciousness and respiratory failure may occur.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. This product is toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not use LARVIN® 75WP except as directed on this label or on other Rhône-Pouleric Ag Company supplemental labeling for this product.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See **Precautionary Statement**, **Statement of Practical Treatment** and **Note to Physician** for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (WARNING), area treated with LARVIN® brand Thiodicarb Insecticide 75WP, date of application, appropriate protective clothing, appropriate reentry interval, if applicable.

STORAGE AND DISPOSAL

STORAGE:

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115°F (46°C). Do not contaminate food, water or feed by storage or disposal. NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

NEVER REUSE EMPTY CONTAINERS. Completely empty bag into application equipment. The dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If bag is burned, stay out of smoke.

ACCIDENTS: In case of a major spill of LARVIN® Thiodicarb Insecticide 75WP;

TELEPHONE (24 HOURS A DAY) IN THE USA 1-800-334-7577.

In case of minor spills or leaks follow all precautions indicated above and cleanup immediately. Soak up with sadd, earth or other suitable material and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

GENERAL INFORMATION

LARVIN® 75WP is a wettable powder concentrate formulation of Thiodicarb Insecticide that readily disperses in water for purposes of spraying by ground or air equipment. Directions on this label are based on tests and field experience relating to effectiveness, impact on environment and residues remaining in food and feed. READ THE ENTIRE LABEL AND OBSERVE ALL LABEL DIRECTIONS AND PRECAUTIONS BEFORE USE.

GENERAL DIRECTIONS

SPRAY PREPARATION

Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or slotted strainers in spray system. To prepare for spraying, fill tank approximately one-half full of water. Add LARVIN® 75WP Insecticide and mix thoroughly by mechanical or hydraulic agitation. Finish filling tank with water to desired volume and thoroughly mix. Do not store spray mixture for prolonged periods. If tank-mixes are to be used, LARVIN® 75WP must be fully dispersed in water first, followed by the addition of the intended tank-mix material.

APPLICATION

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, Professional Consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area. Apply a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground for sweet corn. Repeat application as needed to maintain control. Use higher desage rates for heavier infestations or larger larvae. Use lower rates for light to moderate infestations.

To clean the sprayer after use, drain and flush with water.

DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including center pivot and solid set. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS

First prepare a suspension of LARVIN® brand Thiodicarb Insecticide 75WP in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of LARVIN® brand 75WP, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of LARVIN® brand 75WP per 1 to 4 gallons of water are recommended). Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of LARVIN® brand 75WP into the irrigation water line so as to deliver the desired rate per acre. The suspension of LARVIN® brand 75WP should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with LARVIN® brand Thiodicarb Insecticide 75WP has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve, and low pressure crain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed, expors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.**

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

COMPATIBILITY

LARVIN® 75WP, when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using any tank mixture of LARVIN® 75WP Insecticide, the compatibility of the mixture be established. Add a small amount of LARVIN® 75WP Insecticide to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. THIS PRODUCT IS NOT MISCIBLE WITH VEGETABLE OIL DILUENTS. TANK MIXTURES SHOULD BE USED WITHIN 6 HOUES AFTER MIXING.

NOTE: LARVIN Thiodicarb Insecticide is hydrolytically sensitive to degradation of active ingredient by strong acids, strong bases, and certain heavy metal oxides and salts of certain fungicides such as Maneb, Mancozeb, Copper Count N-8, and Bordeaux mixture. DO NOT USE LARVIN 75®WP INSECTICIDE IN SPRAY MIXTURES WITH THE ABOVE METAL PRODUCTS. DO NOT ADD LARVIN® 75WP TO WATER WITH pH VALUES BELOW 3.0 OR ABOVE 8.5. If necessary, water should be buffered within this range before adding LARVIN® Thiodicarb Insecticide

RECOMMENDED USES

		DOSAGE PER ACRE		
CROP	INSECT	POUNDS ACTIVE	FLUID OUNCES	SPECIFIC DIRECTIONS
Sweet Corn (Fresh Market Only)	Corn earworm European corn borer Armyworms Western bean cutworm	0.5 to 0.75 (Florida Only)	0.66 to 1.00	Insects attacking silks and ears: Apply at 1-7 day intervals starting when silks first appear and continuing until silks begin to dry or infestation potential ceases. Follow local recommendations for proper timing. Whorl or foliage feeders: For best
				results, use sprays of 15 gallons or more per acre applied by ground equipment using solid or hollow cone nozzles directed into the plant whorl. Also refer to general directions for use.

PREHARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

To avoid illegal residues in or on: SWEET CORN

- After silk initiation, do not exceed 7.5 pounds of active ingredient (10 pounds of LARVIN® 75WP) per acre per season.
- Do not allow livestock to graze treated fields.
- · Do not feed treated corn silage (green plant) or fodder to livestock.
- · Ears may be harvested on day of last application.

FIELD USE CONVERSION TABLE

DOSAGE P		
POUNDS ACTIVE	FLUID OUNCES	ACRES TREATED BY 10 LBS. OF LARVIN® 75WP
0.25	0.33	30.3
0.33	0.44	22.7
0.40	0.53	18.9 • • •
0.45	0. 60	16.6
0.50	0. 66	15.2•
0.55	0.73	13.7
0.60	0.80	12.5
0.65	0.87	11.5
0.70	0. 93	10.7°
0.75	1.00	10.0 • • • •

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experiences. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO THE BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company P.O. Box 12014, 2 T. W. Alexander Drive Research Triangle Park, North Carolina 27709

U.S. Patent No.: 4,382,957

LARVIN® is a registered trademark of Rhône-Poulenc LARVIN® 75WP (M ASTER) Notification 6/25/93.

