

# LARVIN<sup>®</sup> brand DF Thiodicarb Insecticide/Ovicide

## 80% Dry Flowable

### ACTIVE INGREDIENT:

(THIODICARB) Dimethyl N, N' [thiobis [(methylimino) carbonyloxy]] bis [ethanimidothioate] .....80% by wt.

INERT INGREDIENTS.....20% by wt.

EPA Reg. No. 264-378

EPA Est. No. 264-MO-02

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-800-334-9745

### STATEMENT OF PRACTICAL TREATMENT

Thiodicarb is an N-methyl carbamate.

**IF SWALLOWED:** If patient is conscious and alert, give 2 to 3 glasses of water or milk to drink. If available, give one tablespoon of Syrup of Ipecac to induce vomiting. If vomiting does not occur within 20 minutes of the first dose, follow with a second and last dose of 1 teaspoon. Alternatively, induce vomiting by touching back of throat with finger. Do not make an unconscious person vomit. GET MEDICAL ATTENTION.

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. GET MEDICAL ATTENTION.

**IF ON SKIN:** Immediately flush skin with plenty of water.

**IF INHALED:** Remove to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. GET MEDICAL ATTENTION.

#### GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway, start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

#### NOTE TO PHYSICIAN

All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains an N-methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.


Gastric lavage may be used if this product has been swallowed. LARVIN<sup>®</sup> poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent over atropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number : 1-800-334-7577.



2913

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

#### WARNING

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Illness may result rapidly following overexposure to this product. May be irritating to eyes and may cause mild skin sensitization. Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Keep away from domestic animals. Discontinue use if allergic reaction occurs. Do not eat or use tobacco while using this product.

#### SIGNS AND SYMPTOMS OF OVEREXPOSURE

Salivation, Muscle tremors, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache.

**IN SEVERE CASES, CONVULSIONS, UNCONSCIOUSNESS, AND RESPIRATORY FAILURE MAY OCCUR.**

#### ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, invertebrates and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from area treated. This product is toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the treatment area.

#### GENERAL INFORMATION

LARVIN<sup>®</sup> brand DF Thiodicarb Insecticide/Ovicide is a dry flowable concentrate (water dispersible granule) formulation that readily disperses in water for spraying by ground or air equipment. Directions on this label are based on tests and field experience relating to effectiveness, impact on environment and residues remaining in food and feed. **READ THE ENTIRE LABEL AND OBSERVE ALL LABEL DIRECTIONS AND PRECAUTIONS BEFORE USE.**

### DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Do not use this product except as directed on this label or on other Rhône-Poulenc Ag Company supplemental labeling for this product. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crops, meat and milk.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

See PRECAUTIONARY STATEMENT, STATEMENT OF PRACTICAL TREATMENT, and NOTE TO PHYSICIAN for information on accidental exposures.

## STORAGE AND DISPOSAL

### STORAGE

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115°F (46°C). NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

If container is damaged, before clean-up, put on long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and chemical-resistant headgear for overhead exposure.

### PESTICIDE DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Contamination with this product will render water, food or feed unfit for human or animal consumption. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL

NEVER REUSE EMPTY CONTAINERS. Completely empty container into application equipment. Then dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

**ACCIDENTS:** In case of a major spill of LARVIN® brand DF Thiodicarb Insecticide/Ovicide TELEPHONE (24 HOURS A DAY) IN THE U.S.A. 1-800-334-7577.

In case of minor spills, follow all precautions indicated above and cleanup immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

## GENERAL DIRECTIONS

### SPRAY PREPARATION

Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent slotted strainers in spray system. To prepare for spraying, fill tank to approximately 1/2 the needed volume of water. Add the required amount of this insecticide and mix thoroughly by mechanical or hydraulic agitation. Finish filling tank with water to desired volume and thoroughly mix. Do not store spray mixture for prolonged periods. If tank mixes are to be used, LARVIN® brand DF Thiodicarb Insecticide/Ovicide must be fully dispersed in water first, followed by the addition of the intended tank-mix material.

### COMPATIBILITY

Thiodicarb Insecticide/Ovicide when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using LARVIN® brand DF Thiodicarb Insecticide/Ovicide in any tank mixture the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. THIS PRODUCT IS NOT COMPATIBLE WITH NON-EMULSIFIED VEGETABLE OR CROP OIL DILUENTS.** FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

**NOTE:** Thiodicarb Insecticide/Ovicide is hydrolytically sensitive to degradation of active ingredient by strong acids, strong bases, and certain heavy metal oxides and salts of certain fungicides. Degradation can cause reduced effectiveness, plant phytotoxicity and increased handling hazards. **DO NOT USE THIS PRODUCT IN SPRAY MIXTURES WITH THE ABOVE TYPE OF PRODUCTS INCLUDING MIXTURES WITH Maneb, Mancozeb (except Manzate 200 and Dithane M-45 wettable powders where mixtures are used immediately after mixing), Copper Count N-8, and Bordeaux mixture. DO NOT ADD LARVIN® BRAND DF THIODICARB INSECTICIDE/OVICIDE TO WATER WITH pH VALUES BELOW 3.0 OR ABOVE 8.5.** If necessary, water should be buffered within this range before adding LARVIN® brand DF Thiodicarb Insecticide/Ovicide.

## DIRECTIONS FOR AERIAL OR GROUND SPRAY APPLICATION

### APPLICATION TIMING

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, Professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

### APPLICATION INSTRUCTIONS

Apply a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground unless otherwise directed under crop specific directions. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground. For best results, it is important to obtain thorough and uniform spray coverage of the plant. Repeat application as needed to maintain control. Use higher dosage rates for heavy infestations, large larvae, or dense foliage. The specific length of residual control depends on environmental factors, plant growth, dosage rate, and degree of insect infestation. Although LARVIN® has some contact activity, larvae not feeding on treated foliage will not be controlled. When banding, determine the amount of chemical to use per acre by dividing the band width by the row width and multiplying by the appropriate broadcast rate.

To clean the sprayer after use, drain and flush with water. Use rinsings on crop according to label instructions or dispose of in an approved manner (See STORAGE AND DISPOSAL).

4 9 13

**RECOMMENDED USES FOR BROCCOLI, CABBAGE AND CAULIFLOWER**

**SPRAY VOLUME FOR BROCCOLI, CABBAGE AND CAULIFLOWER:** Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COLE CROPS (BROCCOLI, CABBAGE, CAULIFLOWER)	Alfalfa Looper Beet Armyworm Cabbage Flea Beetle Imported Cabbageworm	0.4 to 0.8	0.5 to 1.0	Spray as needed. As insect populations increase and/or new plant growth is rapid, the spray interval should be shorter. Use higher dosage rates for heavier infestations or large larvae. Use lower rates for light to moderate infestations.  <b>NOTE:</b> Regional differences have been noted in the susceptibility of certain strains of diamondback moth to thiodicarb. If local experience indicates inadequate control, use an alternate pesticide.	Do not exceed 6.0 pounds of active ingredient (7.5 pounds of this product) per acre per season. Do not apply less than 7 days before harvest.
	Cabbage looper Diamondback moth	0.6 to 1.0	0.75 to 1.25		

**RECOMMENDED USES FOR COTTON**

**INTEGRATED PEST MANAGEMENT (IPM) USE OF THIS PRODUCT**

LARVIN® Thiodicarb Insecticide/Ovicide has both contact and residual ovicidal activity which can be effectively utilized in IPM programs (See ovicide section). LARVIN® Thiodicarb Insecticide/Ovicide has been shown to leave substantial populations of beneficial insects and spiders after use. The lower rates allow for maximum beneficial survival and faster rebound of beneficial populations. Larvicidal and ovicidal control coupled with retention of beneficial insects and spiders can offer significant benefits to those producers utilizing integrated pest control programs.

**SPRAY VOLUME FOR COTTON**

LARVIN® brand DF Thiodicarb Insecticide/Ovicide should be applied in a minimum finished spray volume of 2 gallons per acre by aircraft and 5 gallons per acre by ground equipment. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON	Ovicide: Cotton bollworm Tobacco budworm	0.125 to 0.25	0.16 to 0.31	Apply on a 3 to 5 day schedule when moth flights begin or sufficient eggs are present. Thorough spray coverage is necessary as eggs must be contacted by LARVIN® for activity to occur. Eggs laid on untreated foliage or not contacted by LARVIN® sprays will not be controlled. If significant number of larvae survive, use higher recommended LARVIN® rates or 0.125 to 0.25 lbs ai/A of LARVIN® in tank mixture combinations with organophosphates, pyrethroids or <i>Bacillus thuringiensis</i> such as DIPEL® ES registered for use on cotton at their label recommended rates.  Observe all restrictions and precautions which appear on the labels of all products to be tank mixed with LARVIN®	Do not apply less than 28 days before harvest.  Do not exceed 0.9 pound active ingredient per acre per application or 5.4 pounds active ingredient per acre per season. Do not apply more than 0.9 pound active ingredient per acre of this product in any 7 day period.  In Arizona and California, apply before bolls begin to open and do not exceed 1.8 pounds active ingredient per acre per season.
	Larvicide: Cotton bollworm Tobacco budworm	0.6 to 0.9	0.75 to 1.13	In this rate range, LARVIN® is a direct contact and residual contact ovicide, as well as an effective larvicide. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Use the higher rate when insect larvae populations are heavier or when infestations are predominantly large larvae (>1/4 inch in size).	

RECOMMENDED USES FOR COTTON (CONTINUED)

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON (cont)	Larvicide (Cont.): Armyworms (such as: Beet, Fall, etc.) Cabbage looper Cotton leaf perforator Cotton leafworm Soybean looper Fleahopper (Suppression) Pink bollworm (Suppression) Plant bugs (Suppression)	0.6 to 0.9	0.75 to 1.12	As pest pressure increases, reduce intervals between applications and/or increase dosage rate as indicated in specific directions below. <b>A rate of 0.75 pounds (0.60 lbs. active) should be used when:</b> For low to moderate insect populations and when maximum survival of beneficial insects and spiders is desired. <b>Rates of 0.94 to 1.12 pounds (0.75 to 0.90 lbs. active) should be used when:</b> Insect populations are heavy or infestation is predominantly large larvae. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Applications for suppression of Pink bollworm should be timed for the presence of adult moths.	Do not apply less than 28 days before harvest.  Do not exceed 0.9 pound active ingredient per acre per application or 5.4 pounds active ingredient per acre per season. Do not apply more than 0.9 pound active ingredient per acre of this product in any 7 day period.  In Arizona and California, apply before bolls begin to open and do not exceed 1.8 pounds active ingredient per acre per season.
	Boll weevil (Suppression)	0.6 to 0.9	0.75 to 1.12	LARVIN® will reduce boll weevil populations comparable to most pyrethroids.  <b>Light Boll Weevil Populations:</b> When spraying LARVIN® on a 5 to 7 day worm control schedule, control of boll weevil will usually be adequate to maintain boll weevil populations below established economic thresholds.  <b>Moderate to Heavy Boll Weevil Populations:</b> When spraying LARVIN® on a worm control schedule, application of a registered boll weevil control product and/or tank mix of LARVIN® plus a registered boll weevil product should be used. A 3 to 5 day interval between sprays may be required to maintain boll weevil control under heavy infestations.	
	Cutworms	0.6	24.0	See "SPECIAL DIRECTIONS ON 'CUTWORMS'"	

**RECOMMENDED USES FOR LEAFY VEGETABLES**

**SPRAY VOLUME FOR LEAFY VEGETABLES:** Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
<b>LEAFY VEGETABLES</b> (Amaranth, arrugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible leaved, garland), cress (garden, upland), dandelion, dock, endive, Florence fennel, lettuce (head, leaf), orach, parsley, purslane (garden, winter), rhubarb, spinach(leaf, fine, new zealand), Swiss chard	Alfalfa looper Armyworms (Such as: Beet, Fall, Southern) <i>Heliothis zea</i> (Bollworm, Corn earworm, Tomato fruitworm)	0.4 to 0.75	0.50 to 0.94	Spray as needed. Use the higher dosage rates for heavier infestations of larger larvae. Use the lower rates for light to moderate infestations.	Do not exceed 1.5 pounds of active ingredient (1.9 pounds of LARVIN® brand DF) per acre per season. Do not apply less than 14 days before harvest.
	Cabbage looper	0.6 to 0.75	0.75 to 0.94		

**RECOMMENDED USES FOR SOYBEANS  
(STANDARD AND LOW SPRAY VOLUMES)**

**STANDARD SPRAY VOLUME FOR SOYBEANS**

Apply in a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum of 10 gallons per acre by ground.

**LOW SPRAY VOLUME FOR SOYBEANS**

Apply in a minimum finished spray volume of 1 gallon per acre by aircraft. Only use the low volume aerial application (i. e. minimum of 1 GPA) where that spray volume will provide adequate coverage of the crop.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
SOYBEAN (Standard Spray Volume Applications)	Podworms (Bollworm, Corn earworm, Tobacco bud- worm) Armyworms (Such as: Beet, Fall, Southern, Yellow- striped, etc.) Green cloverworm Velvetbean caterpillar Woollybear caterpillar	0.25 to 0.75	0.31 to 0.94	Use the lower rates for low to moderate populations and maximum protection of beneficials.  For loopers, use low rate (0.45) for newly hatched worms, moderate rate (0.60) for young worms, and use high rates (0.75) for large worms. Use at least 5 gallons of finished spray material per acre if applied by air.	Do not feed forage, hay or straw to livestock. Do not apply less than 28 days before harvest.  Do not exceed 0.75 pound active ingredient per acre per application or 3.0 pounds active ingredient per acre per season. Do not apply more than 0.75 pound active ingredient per acre of this product in any 7 day period.
	Soybean looper Cabbage looper Mexican bean beetle Bean leaf beetle Stinkbugs (Suppression) Three-Cornered alfalfa hopper (Suppression)	0.45 to 0.75	0.56 to 0.94		
	Cutworms	See "SPECIAL DIRECTIONS ON CUTWORMS"			
SOYBEAN (Low Spray Volume Applications)	Podworms (Bollworm, Corn earworm Tobacco bud- worm) Green Cloverworm Velvetbean caterpillar Armyworms (such as Fall, Beet, Southern, etc.) Woollybear caterpillar	0.25 to 0.75	0.31 to 0.94	For best results, use the lower rates for low to moderate populations and maximum protection of beneficials.  When applying at 1 gallon of spray volume per acre by air, use 1 quart of emulsified crop or vegetable oil (e.g., emulsified corn, soybean or cottonseed oil). Water alone may be used as a diluent for spray volumes of 2 gallons or higher per acre.  <b>TANK MIX DIRECTIONS FOR EMULSIFIED CROP OR VEGETABLE OILS:</b> Fill spray tank with approximately 3/4 of the water required for the spray mixture. Add the required amount of LARVIN® DF to the spray tank with the agitator running. Then add the desired amount of emulsified crop or vegetable oil. Continue agitation while adding the remainder of water.	FOR USE ONLY IN MARYLAND, VIRGINIA, NORTH CAROLINA, DELAWARE, SOUTH CAROLINA, GEORGIA, ALABAMA, MISSISSIPPI, LOUISIANA, ARKANSAS AND MISSOURI.  Do not feed forage, hay, or straw to livestock. Do not apply less than 28 days before harvest.  Do not exceed 0.75 pound active ingredient per acre per application or 3.0 pounds active ingredient per acre per season. Do not apply more than 0.75 pound active ingredient per acre of this product in any 7 day period.



**RECOMMENDED USES FOR SWEET CORN**

**SPRAY VOLUME FOR SWEET CORN:** Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground. For aerial applications, only water may be used as a diluent and total spray volume must be at least 2 gallons per acre.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
SWEET CORN (Florida Fresh Market Only)	Corn earworm	0.5	0.63	<p><b>Whorl Feeders:</b> Spray as needed. As insect populations increase and/or when new plant growth is rapid, the spray interval should be shorter. For best results, use spray volumes of 15 gallons or more per acre applied by ground equipment using solid or hollow cone nozzles directed into the plant whorl.</p> <p><b>Insects Attacking Silks And Ears:</b> Apply at 1 to 7 day intervals starting when silks first appear and continuing until harvest, if necessary. As insect populations increase and/or silk growth is rapid, the spray interval should be shorter. Use the higher dosage rate and apply at 1 to 2 day intervals following periods of heavy moth flights. Follow local recommendations for proper timing.</p> <p><b>Foliage Feeders:</b> Use at least 2 gallons finished spray volume per acre by air or 10 gallons per acre by ground.</p>	<p>After silk initiation, do not exceed 7.5 pounds of active ingredient (9.38 pounds of this product) per acre per season.</p> <p>Do not allow livestock to graze treated fields.</p> <p>Do not feed treated corn silage (green plant) or fodder to livestock.</p> <p>Ears may be harvested on day of last application.</p>
	European corn borer	to 0.75	to 0.94		
	Amyworms (Such as Fall, Beet, etc).				
	Western bean cutworm				
	Cutworms	See "SPECIAL DIRECTIONS ON "CUTWORMS"			

**RECOMMENDED USES FOR CUTWORMS**

**SPECIAL DIRECTIONS ON CUTWORMS**

**SPRAY VOLUME FOR CUTWORMS:** Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON, SOYBEANS, SWEET CORN (Florida Fresh Market Only)	Cutworms	0.5 to 0.75	0.63 to 0.94	<p><b>Postemergence Rescue Treatments:</b> For best results, apply as a broadcast spray when cutworms are actively feeding. If cutworms are feeding below the ground, control will be more difficult. Light cultivation with a rotary hoe before spraying may enhance control. If banding, use a minimum width of 10 inches over the row. When banding, determine the amount of chemical to use per acre, by dividing the band width by the row width and multiplying by the appropriate broadcast rate.</p>	Refer to each specific crop for use restrictions.

**RECOMMENDED USES FOR ORNAMENTALS AND TREES**

For dilute-spray ground applications to ornamentals (such as: Trees, Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants and Non-bearing fruits): Apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage. For aerial application to ornamental plants, shrubs and trees (except forests), apply the specified dosage per acre in a minimum total spray volume of 5 gallons to provide thorough coverage. BEFORE MAKING TREATMENTS REFER TO "PLANT TOLERANCE INFORMATION" LISTED ON THIS LABEL.

SITE	PEST	POUNDS OF LARVIN® BRAND DF PER 100 GALS SPRAY/ OR PER ACRE	SPECIFIC DIRECTIONS	USE RESTRICTIONS
<p><b>ORNAMENTALS</b> such as: Trees (except forests), Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants, Non-bearing fruit trees <b>(Citrus:</b> Grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats and hybrids;  <b>Nut Crops:</b> almonds, filberts, pecans, walnuts;  <b>Pome Fruits:</b> apples, pears, loquats, crabapples, oriental pears, quince;  <b>Stone Fruits:</b> peaches, apricots, nectarines, plums, prunes and cherries)</p>	<p>Armyworms  Bagworm Cankerworms (such as: spring, fall)  Gypsy moth Leafrollers (such as: omnivorous, fruittree)  Spruce budworms Webworms (such as: fall, mimosa)  Tent caterpillars</p>	<p>0.63 to 0.94</p>	<p>See "PLANT TOLERANCE INFORMATION" and "APPLICATION" sections of this label.  Apply when pests appear and thereafter as needed not to exceed 6 applications per site per season.</p>	<p>Do not use on fruit trees or vines that will bear edible fruit within one year.</p>

**RECOMMENDED USES FOR NON-CROP AREAS**

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
<b>NON-CROP AREAS</b> (Wastelands, Roadsides, Rights-of-Way, Grassways, Shelter belts, and Fencerows)	Armyworms	0.40 to 0.76	0.50	Treat non-crop areas where pests congregate to reduce infestation levels or to prevent movement into croplands. Apply when pests appear. Make additional applications as needed, not to exceed 4.56 lbs. of active ingredient per site per season.	Do not allow livestock to graze treated areas. Do not use treated plants as food or feed.
	Bagworm		to		
	Budworms	0.76	0.95		
	Cankerworms (such as: spring, fall)				
	Cutworms				
	Gypsy moth				
	Heliothis spp. (such as: corn earworm)				
	Leafrollers (such as: omnivorous, fruittree)				
	Loopers				
	Oakworms				
	Sawflies				
	Skippers				
	Tent caterpillars				
	Webworms (such as: fall, mimosa)				

**ORNAMENTAL PLANT TOLERANCE INFORMATION**

LARVIN<sup>®</sup> brand Thiodicarb Insecticide/Ovicide has been tested on many species of ornamental plants. Results indicate plants listed below are tolerant to the higher rates of this product. Horticultural varieties of many plants differ significantly in sensitivity to pesticides. Before making applications to plants not listed, it is recommended that a small portion of the crop be treated to determine its tolerance to this product. Plants stressed from heat, moisture or other factors are usually more sensitive to sprays. Phytotoxic reactions can be minimized by spraying when stress is minimal such as early mornings or late evenings.

**ORNAMENTAL PLANTS TOLERANT TO THIODICARB INSECTICIDE/OVICIDE**

TREES	SHRUBS	HERBACEOUS GREENHOUSE PLANTS		HERBACEOUS BEDDING PLANTS
Apple, (Crab)	Arborvitae	<i>Acalypha hispida</i>	Ixora	Calendula
Birch, (Weeping, White)	Aucuba	<i>Acalypha wilkesiana</i>	Jade Plant	Canna
Honeylocust, (Sunburst)	Azalea	<i>Aechmea lueddemanniana</i>	<i>Kalanchoe daigremontiana</i>	Chrysanthemum
Maple, (Sugar)	Boxwood	<i>Alternanthera versicolor</i>	<i>Maranta leuconeura erythronera</i>	Coleus
Pine, (Scotch)	Camellia	Aluminum Plant	<i>Neprolepis exaltata</i>	Geranium
Spruce, (Norway)	Elaeagnus	<i>Begonia semperflorens</i>	<i>Peperomia magnoliifolia</i>	Hosta
	Euonymus	Cactus	<i>Peperomia obtusifolia</i>	Impatiens
	Fatsia, (Japanese)	<i>Ceropegia woodii</i>	<i>Philodendron oxycardium</i>	Lantana
	Gardenia	<i>Chamaedorea elegans</i>	Piggyback Plant	Periwinkle
	Hawthorn	<i>Chlorophytum comosum</i>	<i>Pilea grandis</i>	Petunia
	Holly, (Japanese)	Dieffenbachia	Poinsettia	Salvia
	Holly, (Yaupon)	<i>Dracaena sanderana</i>	Pseuderanthemum	Snapdragon
	Juniper	Echeveria	<i>Rhoeo spathacea</i>	Verbena
	Lantana	Fern	<i>Sansevieria trifascia</i>	Violet
	Ligustrum	<i>Ficus diversifolia</i>	Senecio	
	Lilac	Gloxinia	<i>Spathiphyllum wallisii</i>	
	Mahonia	<i>Gynura samentosa</i>	<i>Syngonium podophyllum</i>	
	Osmanthus	<i>Hechtia tellandsioides</i>	Torenia	
	Pittosporum	Hoya	<i>Zebrina pendula</i>	
	Pyracantha	<i>Iboza riparia</i>	<i>Zygocactus truncata</i>	
	Rhododendron	Ivy, (Swedish)		
	Rose, (Hybrid Tea)			
	Yew			

**FIELD USE CONVERSION TABLE**

Pounds Active	Pounds LARVIN <sup>®</sup> DF	Acres Treated by 12.5 lbs. of LARVIN <sup>®</sup> brand DF
0.125	0.16	78.0
0.25	0.31	40.0
0.33	0.41	30.3
0.40	0.50	25.0
0.45	0.56	22.2
0.50	0.63	20.2
0.55	0.69	18.4
0.60	0.75	16.7
0.65	0.81	15.4
0.70	0.87	14.4
0.75	0.94	13.3
0.80	1.00	12.5
0.90	1.13	11.2

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experiences. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

### NOTICE TO THE BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

LARVIN is a registered trademark of Rhône-Poulenc.  
U.S. Patent No.: 4,382,957

LARVIN brand DF Thiodicarb Insecticide/Ovicide 80% Dry Flowable (PENDING) Submitted 2/21/96.

# LARVIN<sup>®</sup> brand DF Thiodicarb Insecticide/Ovicide

## 80% Dry Flowable

### ACTIVE INGREDIENT:

(THIODICARB) Dimethyl N, N' [thiobis [(methylimino) carbonyloxy]] bis [ethanimidothioate] .....80% by wt.

INERT INGREDIENTS.....20% by wt.

EPA Reg. No. 264-378

EPA Est. No. 264-MO-02

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-334-9745

### STATEMENT OF PRACTICAL TREATMENT

Thiodicarb is an N-methyl carbamate.

**IF SWALLOWED:** If patient is conscious and alert, give 2 to 3 glasses of water or milk to drink. If available, give one tablespoon of Syrup of Ipecac to induce vomiting. If vomiting does not occur within 20 minutes of the first dose, follow with a second and last dose of 1 teaspoon. Alternatively, induce vomiting by touching back of throat with finger. Do not make an unconscious person vomit. GET MEDICAL ATTENTION.

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. GET MEDICAL ATTENTION.

**IF ON SKIN:** Immediately flush skin with plenty of water.

**IF INHALED:** Remove to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. GET MEDICAL ATTENTION.

#### GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway, start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.** If poisoning is suspected in animals, contact a veterinarian.

#### NOTE TO PHYSICIAN

All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains an N-methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. LARVIN<sup>®</sup> poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent over atropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number: 1-800-334-7577.

**ACCEPTED**  
**MAY 13 1996**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 264-378

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

#### WARNING

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Illness may result rapidly following overexposure to this product. May be irritating to eyes and may cause mild skin sensitization. Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Keep away from domestic animals. Discontinue use if allergic reaction occurs. Do not eat or use tobacco while using this product.

#### SIGNS AND SYMPTOMS OF OVEREXPOSURE

Salivation, Muscle tremors, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache.

**IN SEVERE CASES, CONVULSIONS, UNCONSCIOUSNESS, AND RESPIRATORY FAILURE MAY OCCUR.**

#### ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, invertebrates and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from area treated. This product is toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the treatment area.

#### GENERAL INFORMATION

LARVIN® brand DF Thiodicarb Insecticide/Ovicide is a dry flowable concentrate (water dispersible granule) formulation that readily disperses in water for spraying by ground or air equipment. Directions on this label are based on tests and field experience relating to effectiveness, impact on environment and residues remaining in food and feed. READ THE ENTIRE LABEL AND OBSERVE ALL LABEL DIRECTIONS AND PRECAUTIONS BEFORE USE.

### DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Do not use this product except as directed on this label or on other Rhône-Poulenc Ag Company supplemental labeling for this product. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crops, meat and milk.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

See PRECAUTIONARY STATEMENT, STATEMENT OF PRACTICAL TREATMENT, and NOTE TO PHYSICIAN for information on accidental exposures.

## STORAGE AND DISPOSAL

### STORAGE

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115°F (46°C). NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

If container is damaged, before clean-up, put on long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and chemical-resistant headgear for overhead exposure.

### PESTICIDE DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Contamination with this product will render water, food or feed unfit for human or animal consumption. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL

NEVER REUSE EMPTY CONTAINERS. Completely empty container into application equipment. Then dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

**ACCIDENTS:** In case of a major spill of LARVIN® brand DF Thiodicarb Insecticide/Ovicide TELEPHONE (24 HOURS A DAY) IN THE U.S.A. 1-800-334-7577.

In case of minor spills, follow all precautions indicated above and cleanup immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

## GENERAL DIRECTIONS

### SPRAY PREPARATION

Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent slotted strainers in spray system. To prepare for spraying, fill tank to approximately 1/2 the needed volume of water. Add the required amount of this insecticide and mix thoroughly by mechanical or hydraulic agitation. Finish filling tank with water to desired volume and thoroughly mix. Do not store spray mixture for prolonged periods. If tank mixes are to be used, LARVIN® brand DF Thiodicarb Insecticide/Ovicide must be fully dispersed in water first, followed by the addition of the intended tank-mix material.

### COMPATIBILITY

Thiodicarb Insecticide/Ovicide when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using LARVIN® brand DF Thiodicarb Insecticide/Ovicide in any tank mixture the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. THIS PRODUCT IS NOT COMPATIBLE WITH NON-EMULSIFIED VEGETABLE OR CROP OIL DILUENTS.** FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

**NOTE:** Thiodicarb Insecticide/Ovicide is hydrolytically sensitive to degradation of active ingredient by strong acids, strong bases, and certain heavy metal oxides and salts of certain fungicides. Degradation can cause reduced effectiveness, plant phytotoxicity and increased handling hazards. **DO NOT USE THIS PRODUCT IN SPRAY MIXTURES WITH THE ABOVE TYPE OF PRODUCTS INCLUDING MIXTURES WITH Maneb, Mancozeb (except Manzate 200 and Dithane M-45 wettable powders where mixtures are used immediately after mixing), Copper Count N-8, and Bordeaux mixture. DO NOT ADD LARVIN® BRAND DF THIODICARB INSECTICIDE/OVICIDE TO WATER WITH pH VALUES BELOW 3.0 OR ABOVE 8.5.** If necessary, water should be buffered within this range before adding LARVIN® brand DF Thiodicarb Insecticide/Ovicide.

## DIRECTIONS FOR AERIAL OR GROUND SPRAY APPLICATION

### APPLICATION TIMING

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, Professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

### APPLICATION INSTRUCTIONS

Apply a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground unless otherwise directed under crop specific directions. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground. For best results, it is important to obtain thorough and uniform spray coverage of the plant. Repeat application as needed to maintain control. Use higher dosage rates for heavy infestations, large larvae, or dense foliage. The specific length of residual control depends on environmental factors, plant growth, dosage rate, and degree of insect infestation. Although LARVIN® has some contact activity, larvae not feeding on treated foliage will not be controlled. When banding, determine the amount of chemical to use per acre by dividing the band width by the row width and multiplying by the appropriate broadcast rate.

To clean the sprayer after use, drain and flush with water. Use rinsings on crop according to label instructions or dispose of in an approved manner (See STORAGE AND DISPOSAL).



4 9 13

**RECOMMENDED USES FOR BROCCOLI, CABBAGE AND CAULIFLOWER**

**SPRAY VOLUME FOR BROCCOLI, CABBAGE AND CAULIFLOWER:** Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COLE CROPS (BROCCOLI, CABBAGE, CAULIFLOWER)	Alfalfa Looper Beet Armyworm Cabbage Flea Beetle Imported Cabbageworm	0.4 to 0.8	0.5 to 1.0	Spray as needed. As insect populations increase and/or new plant growth is rapid, the spray interval should be shorter. Use higher dosage rates for heavier infestations or large larvae. Use lower rates for light to moderate infestations.  <b>NOTE:</b> Regional differences have been noted in the susceptibility of certain strains of diamondback moth to thiodicarb. If local experience indicates inadequate control, use an alternate pesticide.	Do not exceed 6.0 pounds of active ingredient (7.5 pounds of this product) per acre per season. Do not apply less than 7 days before harvest.
	Cabbage looper Diamondback moth	0.6 to 1.0	0.75 to 1.25		

## RECOMMENDED USES FOR COTTON

### INTEGRATED PEST MANAGEMENT (IPM) USE OF THIS PRODUCT

LARVIN® Thiodicarb Insecticide/Ovicide has both contact and residual ovicidal activity which can be effectively utilized in IPM programs (See ovicide section). LARVIN® Thiodicarb Insecticide/Ovicide has been shown to leave substantial populations of beneficial insects and spiders after use. The lower rates allow for maximum beneficial survival and faster rebound of beneficial populations. Larvicidal and ovicidal control coupled with retention of beneficial insects and spiders can offer significant benefits to those producers utilizing integrated pest control programs.

### SPRAY VOLUME FOR COTTON

LARVIN® brand DF Thiodicarb Insecticide/Ovicide should be applied in a minimum finished spray volume of 2 gallons per acre by aircraft and 5 gallons per acre by ground equipment. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON	Ovicide: Cotton bollworm Tobacco budworm	0.125 to 0.25	0.16 to 0.31	Apply on a 3 to 5 day schedule when moth flights begin or sufficient eggs are present. Thorough spray coverage is necessary as eggs must be contacted by LARVIN® for activity to occur. Eggs laid on untreated foliage or not contacted by LARVIN® sprays will not be controlled. If significant number of larvae survive, use higher recommended LARVIN® rates or 0.125 to 0.25 lbs ai/A of LARVIN® in tank mixture combinations with organophosphates, pyrethroids or <i>Bacillus thuringiensis</i> such as DIPEL® ES registered for use on cotton at their label recommended rates.  Observe all restrictions and precautions which appear on the labels of all products to be tank mixed with LARVIN®	Do not apply less than 28 days before harvest. Do not exceed 0.9 pound active ingredient per acre per application or 5.4 pounds active ingredient per acre per season. Do not apply more than 0.9 pound active ingredient per acre of this product in any 7 day period.  In Arizona and California, apply before bolls begin to open and do not exceed 1.8 pounds active ingredient per acre per season.
	Larvicide: Cotton bollworm Tobacco budworm	0.6 to 0.9	0.75 to 1.13	In this rate range, LARVIN® is a direct contact and residual contact ovicide, as well as an effective larvicide. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Use the higher rate when insect larvae populations are heavier or when infestations are predominantly large larvae (>1/4 inch in size).	

RECOMMENDED USES FOR COTTON (CONTINUED)

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON (cont)	Larvicide (Cont.): Armyworms (such as: Beet, Fall, etc.) Cabbage looper Cotton leaf perforator Cotton leafworm Soybean looper Fleahopper (Suppression) Pink bollworm (Suppression) Plant bugs (Suppression)	0.6 to 0.9	0.75 to 1.12	As pest pressure increases, reduce intervals between applications and/or increase dosage rate as indicated in specific directions below. <b>A rate of 0.75 pounds (0.60 lbs. active) should be used when:</b> For low to moderate insect populations and when maximum survival of beneficial insects and spiders is desired. <b>Rates of 0.94 to 1.12 pounds (0.75 to 0.90 lbs. active) should be used when:</b> Insect populations are heavy or infestation is predominantly large larvae. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Applications for suppression of Pink bollworm should be timed for the presence of adult moths.	Do not apply less than 28 days before harvest.  Do not exceed 0.9 pound active ingredient per acre per application or 5.4 pounds active ingredient per acre per season. Do not apply more than 0.9 pound active ingredient per acre of this product in any 7 day period.  In Arizona and California, apply before bolls begin to open and do not exceed 1.8 pounds active ingredient per acre per season.
	Boll weevil (Suppression)	0.6 to 0.9	0.75 to 1.12	LARVIN® will reduce boll weevil populations comparable to most pyrethroids.  <b>Light Boll Weevil Populations:</b> When spraying LARVIN® on a 5 to 7 day worm control schedule, control of boll weevil will usually be adequate to maintain boll weevil populations below established economic thresholds.  <b>Moderate to Heavy Boll Weevil Populations:</b> When spraying LARVIN® on a worm control schedule, application of a registered boll weevil control product and/or tank mix of LARVIN® plus a registered boll weevil product should be used. A 3 to 5 day interval between sprays may be required to maintain boll weevil control under heavy infestations.	
	Cutworms	0.6	24.0	See "SPECIAL DIRECTIONS ON CUTWORMS"	

**RECOMMENDED USES FOR LEAFY VEGETABLES**

**SPRAY VOLUME FOR LEAFY VEGETABLES:** Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
<b>LEAFY VEGETABLES</b> (Amaranth, arrugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible leaved, garland), cress (garden, upland), dandelion, dock, endive, Florence fennel, lettuce (head, leaf), orach, parsley, purslane (garden, winter), rhubarb, spinach(leaf, fine, new zealand), Swiss chard)	Alfalfa looper	0.4	0.50	Spray as needed. Use the higher dosage rates for heavier infestations of larger larvae. Use the lower rates for light to moderate infestations.	Do not exceed 1.5 pounds of active ingredient (1.9 pounds of LARVIN® brand DF) per acre per season. Do not apply less than 14 days before harvest.
	Armyworms (Such as: Beet, Fall, Southern) <i>Heliothis zea</i> (Bollworm, Corn earworm, Tomato fruitworm)	to 0.75	to 0.94		
	Cabbage looper	0.6 to 0.75	0.75 to 0.94		

## RECOMMENDED USES FOR SOYBEANS (STANDARD AND LOW SPRAY VOLUMES)

### STANDARD SPRAY VOLUME FOR SOYBEANS

Apply in a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum of 10 gallons per acre by ground.

### LOW SPRAY VOLUME FOR SOYBEANS

Apply in a minimum finished spray volume of 1 gallon per acre by aircraft. **Only** use the low volume aerial application (i. e. minimum of 1 GPA) where that spray volume will provide adequate coverage of the crop.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
SOYBEAN (Standard Spray Volume Applications)	Podworms (Bollworm, Corn earworm, Tobacco bud- worm) Armyworms (Such as: Beet, Fall, Southern, Yellow- striped, etc.) Green cloverworm Velvetbean caterpillar Woollybear caterpillar	0.25 to 0.75	0.31 to 0.94	Use the lower rates for low to moderate populations and maximum protection of beneficials.  For loopers, use low rate (0.45) for newly hatched worms, moderate rate (0.60) for young worms, and use high rates (0.75) for large worms. Use at least 5 gallons of finished spray material per acre if applied by air.	Do not feed forage, hay or straw to livestock. Do not apply less than 28 days before harvest.  Do not exceed 0.75 pound active ingredient per acre per application or 3.0 pounds active ingredient per acre per season. Do not apply more than 0.75 pound active ingredient per acre of this product in any 7 day period.
	Soybean looper Cabbage looper Mexican bean beetle Bean leaf beetle Stinkbugs (Suppression) Three-Cornered alfalfa hopper (Suppression)	0.45 to 0.75	0.56 to 0.94		
	Cutworms	See "SPECIAL DIRECTIONS ON CUTWORMS"			
SOYBEAN (Low Spray Volume Applications)	Podworms (Bollworm, Corn earworm Tobacco bud- worm) Green Cloverworm Velvetbean caterpillar Armyworms (such as Fall, Beet, Southern, etc.) Woollybear caterpillar	0.25 to 0.75	0.31 to 0.94	For best results, use the lower rates for low to moderate populations and maximum protection of beneficials.  When applying at 1 gallon of spray volume per acre by air, use 1 quart of emulsified crop or vegetable oil (e.g., emulsified corn, soybean or cottonseed oil). Water alone may be used as a diluent for spray volumes of 2 gallons or higher per acre.  <b>TANK MIX DIRECTIONS FOR EMULSIFIED CROP OR VEGETABLE OILS:</b> Fill spray tank with approximately 3/4 of the water required for the spray mixture. Add the required amount of LARVIN® DF to the spray tank with the agitator running. Then add the desired amount of emulsified crop or vegetable oil. Continue agitation while adding the remainder of water.	FOR USE ONLY IN MARYLAND, VIRGINIA, NORTH CAROLINA, DELAWARE, SOUTH CAROLINA, GEORGIA, ALABAMA, MISSISSIPPI, LOUISIANA, ARKANSAS AND MISSOURI.  Do not feed forage, hay, or straw to livestock. Do not apply less than 28 days before harvest.  Do not exceed 0.75 pound active ingredient per acre per application or 3.0 pounds active ingredient per acre per season. Do not apply more than 0.75 pound active ingredient per acre of this product in any 7 day period.

### RECOMMENDED USES FOR SWEET CORN

**SPRAY VOLUME FOR SWEET CORN:** Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground. For aerial applications, only water may be used as a diluent and total spray volume must be at least 2 gallons per acre.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
SWEET CORN (Florida Fresh Market Only)	Corn earworm European corn borer Armyworms (Such as Fall, Beet, etc.) Western bean cutworm	0.5 to 0.75	0.63 to 0.94	<p><b>Whorl Feeders:</b> Spray as needed. As insect populations increase and/or when new plant growth is rapid, the spray interval should be shorter. For best results, use spray volumes of 15 gallons or more per acre applied by ground equipment using solid or hollow cone nozzles directed into the plant whorl.</p> <p><b>Insects Attacking Silks And Ears:</b> Apply at 1 to 7 day intervals starting when silks first appear and continuing until harvest, if necessary. As insect populations increase and/or silk growth is rapid, the spray interval should be shorter. Use the higher dosage rate and apply at 1 to 2 day intervals following periods of heavy moth flights. Follow local recommendations for proper timing.</p> <p><b>Foliage Feeders:</b> Use at least 2 gallons finished spray volume per acre by air or 10 gallons per acre by ground.</p>	<p>After silk initiation, do not exceed 7.5 pounds of active ingredient (9.38 pounds of this product) per acre per season.</p> <p>Do not allow livestock to graze treated fields.</p> <p>Do not feed treated corn silage (green plant) or fodder to livestock.</p> <p>Ears may be harvested on day of last application.</p>
	Cutworms	See "SPECIAL DIRECTIONS ON "CUTWORMS"			

### RECOMMENDED USES FOR CUTWORMS

#### SPECIAL DIRECTIONS ON CUTWORMS

**SPRAY VOLUME FOR CUTWORMS:** Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
COTTON, SOYBEANS, SWEET CORN (Florida Fresh Market Only)	Cutworms	0.5 to 0.75	0.63 to 0.94	<p><b>Postemergence Rescue Treatments:</b> For best results, apply as a broadcast spray when cutworms are actively feeding. If cutworms are feeding below the ground, control will be more difficult. Light cultivation with a rotary hoe before spraying may enhance control. If banding, use a minimum width of 10 inches over the row. When banding, determine the amount of chemical to use per acre, by dividing the band width by the row width and multiplying by the appropriate broadcast rate.</p>	Refer to each specific crop for use restrictions.

**RECOMMENDED USES FOR ORNAMENTALS AND TREES**

For dilute-spray ground applications to ornamentals (such as: Trees, Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants and Non-bearing fruits): Apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage. For aerial application to ornamental plants, shrubs and trees (except forests), apply the specified dosage per acre in a minimum total spray volume of 5 gallons to provide thorough coverage. BEFORE MAKING TREATMENTS REFER TO "PLANT TOLERANCE INFORMATION" LISTED ON THIS LABEL.

SITE	PEST	POUNDS OF LARVIN® BRAND DF PER 100 GALS SPRAY/ OR PER ACRE	SPECIFIC DIRECTIONS	USE RESTRICTIONS
<p><b>ORNAMENTALS</b> such as: Trees (except forests), Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants, Non-bearing fruit trees <b>(Citrus:</b> Grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats and hybrids;  <b>Nut Crops:</b> almonds, filberts, pecans, walnuts;  <b>Pome Fruits:</b> apples, pears, loquats, crabapples, oriental pears, quince;  <b>Stone Fruits:</b> peaches, apricots, nectarines, plums, prunes and cherries)</p>	<p>Armyworms Bagworm Cankerworms (such as: spring, fall) Gypsy moth Leafrollers (such as: omnivorous, fruittree) Spruce budworms Webworms (such as: fall, mimosa) Tent caterpillars</p>	<p>0.63 to 0.94</p>	<p>See "PLANT TOLERANCE INFORMATION" and "APPLICATION" sections of this label.  Apply when pests appear and thereafter as needed not to exceed 6 applications per site per season.</p>	<p>Do not use on fruit trees or vines that will bear edible fruit within one year.</p>

## RECOMMENDED USES FOR NON-CROP AREAS

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® DF		
<b>NON-CROP AREAS</b> (Wastelands, Roadsides, Rights-of-Way, Grassways, Shelter belts, and Fencerows)	Armyworms	0.40 to 0.76	0.50 to 0.95	Treat non-crop areas where pests congregate to reduce infestation levels or to prevent movement into croplands. Apply when pests appear. Make additional applications as needed, not to exceed 4.56 lbs. of active ingredient per site per season.	Do not allow livestock to graze treated areas. Do not use treated plants as food or feed.
	Bagworm				
	Budworms				
	Cankerworms (such as: spring, fall)				
	Cutworms				
	Gypsy moth				
	Heliothis spp. (such as: corn earworm)				
	Leafrollers (such as: omnivorous, fruittree)				
	Loopers				
	Oakworms				
	Sawflies				
	Skippers				
	Tent caterpillars				
Webworms (such as: fall, mimosa)					



## ORNAMENTAL PLANT TOLERANCE INFORMATION

LARVIN® brand Thiodicarb Insecticide/Ovicide has been tested on many species of ornamental plants. Results indicate plants listed below are tolerant to the higher rates of this product. Horticultural varieties of many plants differ significantly in sensitivity to pesticides. Before making applications to plants not listed, it is recommended that a small portion of the crop be treated to determine its tolerance to this product. Plants stressed from heat, moisture or other factors are usually more sensitive to sprays. Phytotoxic reactions can be minimized by spraying when stress is minimal such as early mornings or late evenings.

### ORNAMENTAL PLANTS TOLERANT TO THIODICARB INSECTICIDE/OVICIDE

TREES	SHRUBS	HERBACEOUS GREENHOUSE PLANTS		HERBACEOUS BEDDING PLANTS
Apple, (Crab)	Arborvitae	<i>Acalypha hispida</i>	Ixora	Calendula
Birch, (Weeping, White)	Aucuba	<i>Acalypha wilkesiana</i>	Jade Plant	Canna
Honeylocust, (Sunburst)	Azalea	<i>Aechmea lueddemanniana</i>	<i>Kalanchoe daigremontiana</i>	Chrysanthemum
Maple, (Sugar)	Boxwood	<i>Alternanthera versicolor</i>	<i>Maranta leuconeura erythroneura</i>	Coleus
Pine, (Scotch)	Camellia	Aluminum Plant	<i>Nephrolepis exaltata</i>	Geranium
Spruce, (Norway)	Elaeagnus	<i>Begonia semperflorens</i>	<i>Peperomia magnoliifolia</i>	Hosta
	Euonymus	Cactus	<i>Peperomia obtusifolia</i>	Impatiens
	Fatsia, (Japanese)	<i>Ceropegia woodii</i>	<i>Philodendron oxycardium</i>	Lantana
	Gardenia	<i>Chamaedorea elegans</i>	Piggyback Plant	Periwinkle
	Hawthorn	<i>Chlorophytum comosum</i>	<i>Pilea grandis</i>	Petunia
	Holly, (Japanese)	Dieffenbachia	Poinsettia	Salvia
	Holly, (Yaupon)	<i>Dracaena sanderana</i>	Pseuderanthemum	Snapdragon
	Juniper	Echeveria	<i>Rhoeo spathacea</i>	Verbena
	Lantana	Fern	<i>Sansevieria trifascia</i>	Violet
	Ligustrum	<i>Ficus diversifolia</i>	Senecio	
	Lilac	Gloxinia	<i>Spathyiphyllum wallisii</i>	
	Mahonia	<i>Gynura sarmentosa</i>	<i>Syngonium podophyllum</i>	
	Osmanthus	<i>Hechtia tellandsioides</i>	Torenia	
	Pittosporum	Hoya	<i>Zebrina pendula</i>	
	Pyracantha	<i>Iboza riparia</i>	<i>Zygocactus truncata</i>	
	Rhododendron	Ivy, (Swedish)		
	Rose, (Hybrid Tea)			
	Yew			

### FIELD USE CONVERSION TABLE

Pounds Active	Pounds LARVIN® DF	Acres Treated by 12.5 lbs. of LARVIN® brand DF
0.125	0.16	78.0
0.25	0.31	40.0
0.33	0.41	30.3
0.40	0.50	25.0
0.45	0.56	22.2
0.50	0.63	20.2
0.55	0.69	18.4
0.60	0.75	16.7
0.65	0.81	15.4
0.70	0.87	14.4
0.75	0.94	13.3
0.80	1.00	12.5
0.90	1.13	11.2

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experiences. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

### NOTICE TO THE BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

LARVIN is a registered trademark of Rhône-Poulenc.  
U.S. Patent No.: 4,382,957

LARVIN brand DF Thiodicarb Insecticide/Ovicide 80% Dry Flowable (PENDING) Submitted 2/21/96.