

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

# LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide Aqueous Flowable

**ACTIVE INGREDIENT**

THIODICARB: Dimethyl N, N' [(thiois [(methylimino) carbonyloxy]] bis [ethanimidothioate]).....	32.5% by wt.
INERT INGREDIENTS .....	67.5% by wt.

CONTAINS 3.2 pounds active thiodicarb per U.S. gallon

EPA Registration No. 264-379

EPA Establishment No. 264-MO-02

## KEEP OUT OF REACH OF CHILDREN

### WARNING

### AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

MAY BE FATAL IF SWALLOWED. MAY BE HARMFUL IF INHALED OR ABSORBED THROUGH THE SKIN. ANTIDOTE IS ATROPINE SULFATE. SEE STATEMENT OF PRACTICAL TREATMENT AND OTHER PRECAUTIONARY STATEMENTS ON LEFT PANEL. READ THE ENTIRE LABEL BEFORE USING THIS THIS PESTICIDE.

For Emergency Information ONLY Call 24 Hours A Day 1-800-334-7577.


For Product Handling, Use and General Information CALL 1-800-334-9745.

RHONE-POULENC AG COMPANY  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

Net contents 2.5 gallons  
(9.46 liters)

U.S. Patent No.: 4,382,957

LARVIN® is the registered trademark of RHONE-POULENC for thiodicarb insecticides



**ACCEPTED**

APR 12 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 264-379

**BEST AVAILABLE COPY**

2  
14

## PRECAUTIONARY STATEMENTS

### WARNING

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Illness may result rapidly following overexposure to this product. May be fatal if swallowed. May be harmful if inhaled or absorbed through the skin. May be irritating to eyes and may cause mild skin sensitization. Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Keep away from domestic animals.

Wear long-sleeved clothing, full-length trousers, head covering and protective gloves while using this product. Remove contaminated clothing daily; wash before reuse. Wash exposed skin thoroughly with soap and water after use and before eating or using tobacco. Discontinue use if allergic reaction occurs. Do not eat or use tobacco while using this product.

#### SIGNS AND SYMPTOMS OF OVEREXPOSURE:

Salivation, Watery eyes, Pinpoint eye pupils, Blurred vision, Muscle tremors, Difficult breathing, Excessive sweating, Abdominal cramps, Nausea, Vomiting, Diarrhea, Weakness, Headache. In severe cases, convulsions, unconsciousness and respiratory failure may occur.

#### ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

#### STATEMENT OF PRACTICAL TREATMENT

##### GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway, start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: If patient is conscious and alert, give 2 to 3 glasses of water or milk to drink. If available, give one tablespoon of Syrup of Ipecac to induce vomiting. If vomiting does not occur within 20 minutes of the first dose, follow with a second and last dose of 1 teaspoon. Alternatively, induce vomiting by touching back of throat with finger. Do not make an unconscious person vomit. GET MEDICAL ATTENTION.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. GET MEDICAL ATTENTION.

IF ON SKIN: Immediately flush skin with plenty of water while removing contaminated clothing and shoes. Wash clothes separately before reuse.

IF INHALED: Remove to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. GET MEDICAL ATTENTION.

**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/82**

**NOTE TO PHYSICIAN**

All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains a methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product has been swallowed. LARVIN® poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number : 1-800-334-7577.

**ENVIRONMENTAL HAZARDS**


This product is toxic to fish, birds, invertebrates and other wildlife. Do not apply directly to lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from area treated. This product is toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the treatment area.

**GENERAL INFORMATION**

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide is an aqueous flowable formulation that readily disperses in water for spraying by ground or air equipment. Directions on this label are based on tests and field experience relating to effectiveness, impact on environment and residues remaining in food and feed. **READ THE ENTIRE LABEL AND OBSERVE ALL LABEL DIRECTIONS AND PRECAUTIONS BEFORE USE.**

**DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.



**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88**

Do not use this product except as directed on this label or on other Rhône-Poulenc Ag Company supplemental labeling for this product. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crops, meat and milk.

**GENERAL WORKER PROTECTION STATEMENTS**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See Precautionary Statement, Statement of Practical Treatment and Note to Physician for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (WARNING), area treated with LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide, date of application, appropriate protective clothing, appropriate reentry interval, if applicable.

**STORAGE AND DISPOSAL**

**STORAGE:** Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115°F (46°C). NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

If container is damaged, before clean-up, put on long-sleeved shirt, full-length trousers, head covering and protective gloves.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Contamination with this product will render water, food or feed unfit for human or animal consumption. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** NEVER REUSE EMPTY CONTAINERS. Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

**ACCIDENTS:** In case of a major spill of LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide: TELEPHONE (24 HOURS A DAY) IN THE U.S.A.: 1-800-334-7577.

**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88**

In case of minor spills or leaks follow all precautions indicated above and cleanup immediately. Soak up with sand, earth or other suitable material and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

**GENERAL DIRECTIONS**

**SPRAY PREPARATION:** TO ASSURE A UNIFORM PRODUCT, AGITATE OR SHAKE ALL CONTAINERS OF THIS PRODUCT PRIOR TO USE. Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent slotted strainers in spray system. To prepare for spraying, fill tank to approximately 1/2 the needed volume of water. Add the required amount of this insecticide and mix thoroughly by mechanical or hydraulic agitation. Finish filling tank with water to desired volume and thoroughly mix. Do not store spray mixture for prolonged periods. If tank mixes are to be used, LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide must be fully dispersed in water first, followed by the addition of the intended tank-mix material.

**COMPATABILITY:** Thiodicarb Insecticide when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide in any tank mixture the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. THIS PRODUCT IS NOT COMPATIBLE WITH NON-EMULSIFIED VEGETABLE OIL DILUENTS. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

**NOTE:** Thiodicarb Insecticide is hydrolytically sensitive to degradation of active ingredient by strong acids, strong bases, and certain heavy metal oxides and salts of certain fungicides. Degradation can cause reduced effectiveness, plant phytotoxicity and increased handling hazards. DO NOT USE THIS PRODUCT IN SPRAY MIXTURES WITH THE ABOVE TYPE OF PRODUCTS INCLUDING MIXTURES WITH Maneb, Mancozeb (except Manzate 200 and Dithane M-45 wettable powders where mixtures are used immediately after mixing), Copper Count N-8, and Bordeaux mixture. Do not use a four way tank mixture of LARVIN® brand 3.2 + Pix® + chlordimeform + Guthion® or Parathion as this mixture is incompatible. DO NOT ADD LARVIN® brand 3.2 THIODICARB INSECTICIDE/OVICIDE TO WATER WITH pH VALUES BELOW 3.0 OR ABOVE 8.5. If necessary, water should be buffered within this range before adding LARVIN® brand Thiodicarb Insecticide/Ovicide.

**DIRECTIONS FOR AERIAL OR GROUND SPRAY APPLICATION**

**APPLICATION TIMING:** Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, Professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

**APPLICATION INSTRUCTIONS:** Apply a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground unless otherwise directed under crop specific directions. For best results, it is important to obtain thorough and uniform spray coverage of the plant. Repeat application as needed to maintain control. Use higher dosage rates for heavy

6  
11

**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88**

infestations, large larvae, or dense foliage. The specific length of residual control depends on environmental factors, plant growth, dosage rate, and degree of insect infestation. When banding, determine the amount of chemical to use per acre by dividing the band width by the row width and multiplying by the appropriate broadcast rate.

To clean the sprayer after use, drain and flush with water. Use rinsings on crop according to label instructions or dispose of in an approved manner (See STORAGE AND DISPOSAL).

**DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS**

Apply this product only through sprinkler irrigation systems including center pivot and solid set. Do not apply this product through any other type of irrigation system.

**SPRAY PREPARATION:** Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

**APPLICATION INSTRUCTIONS:** First prepare a suspension of LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of LARVIN® 3.2, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of LARVIN® 3.2 per 1 to 4 gallons of water are recommended). Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of LARVIN® 3.2 into the irrigation water line so as to deliver the desired rate per acre. The suspension of LARVIN® 3.2 should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

**NOTE:** When treatment with LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

**GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS**

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually

**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88**

shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

RECOMMENDED USES

COTTON

INTEGRATED PEST MANAGEMENT (IPM) USE OF THIS PRODUCT: LARVIN® Thiodicarb Insecticide/Ovicide has both contact and residual ovicidal activity which can be effectively utilized in IPM programs (See ovicide section). LARVIN® Thiodicarb Insecticide/Ovicide has been shown to leave substantial populations of beneficial insects and spiders after use. The lower rates allow for maximum beneficial survival and faster rebound of beneficial populations. Larvicidal and ovicidal control coupled with retention of beneficial insects and spiders can offer significant benefits to those producers utilizing integrated pest control programs.

SPRAY VOLUME FOR COTTON: LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide should be applied in a minimum finished spray volume of 2 gallons per acre by aircraft and 5 gallons per acre by ground equipment.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Fluid Ounces		
COTTON (NOT FOR USE IN CALIFORNIA AND ARIZONA)	Ovicide:				
	Cotton bollworm	0.125	5.0	Apply in tank mix combinations with pyrethroid or organophosphate insecticides at their standard rate. Thorough spray coverage is necessary as eggs must be contacted by LARVIN® in order for activity to occur. Eggs laid on untreated foliage or not contacted by LARVIN® will not be controlled. Initiate sprays when sufficient eggs and newly hatched larvae are present.	Do not allow livestock to graze fields. Do not apply less than 28 days before harvest.
Tobacco budworm	to 0.25	to 10.0			
	Larvicide:				
	Cotton bollworm	0.6	24.0	In this rate range, LARVIN® is a direct contact and residual contact ovicide, as well as an effective larvicide. Use the higher rate when insect larvae populations are heavier or when infestations are predominantly large larvae (> 1/4 inch in size). Larvae present in bolls will not be controlled.	
	Tobacco budworm	to 0.9	to 36.0		

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Fluid Ounces		
COTTON (Cont.)	Larvicide (Cont.):	0.6	24.0	<p>As pest pressure increases, reduce intervals between applications and/or increase dosage rate as indicated in specific directions below.</p> <p><u>A rate of 24 fluid ounces (0.60 lbs. active) should be used when:</u></p> <p>Maximum survival of beneficial insects and spiders is desired.</p> <p><u>Rates at 30 to 36 fluid ounces (0.75 to 0.90 lbs. active) should be used when:</u></p> <p>Insect populations are heavy or infestation is predominantly large larvae. Larvae present in bolls will not be controlled.</p> <p>Applications for control of Pink bollworm should be timed for the presence of adult moths.</p>	<p>Do not allow livestock to graze fields. Do not apply less than 28 days before harvest.</p>
	Amyworms	to	to		
	(such as:	0.9	36.0		
	Beet, Fall, etc.)				
	Cabbage looper				
	Cotton leafperforator				
	Cotton leafworm				
	Fleahopper (suppression)				
	Pink bollworm				
	Plant bugs				
(Suppression)					
Soybean looper					
	Boll weevil	0.6	24.0	<p>LARVIN® will reduce boll weevil populations comparable to most pyrethroids.</p> <p><u>LIGHT BOLL WEEVIL POPULATIONS</u></p> <p>When spraying LARVIN® on a 5 to 7 day worm control schedule, control of boll weevil will usually be adequate to maintain boll weevil populations below established economic thresholds.</p> <p><u>MODERATE TO HEAVY BOLL WEEVIL POPULATIONS:</u></p> <p>When spraying LARVIN® on a worm control schedule, intersprays of a registered boll weevil control product and/or tank mix of LARVIN® plus a registered boll weevil product should be used. A 3 to 5 day interval between sprays may be required to maintain boll weevil control under heavy infestations.</p>	
(Suppression)	to	to			
		0.9	36.0		
	Cutworms	0.6	24.0	See "SPECIAL DIRECTIONS ON CUTWORMS"	

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Fluid Ounces		
SOYBEAN	Podworms	0.25	10.0	Use the lower rates for low to moderate populations and maximum protection of beneficials.  Apply in a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground.	Do not feed forage, hay, or straw to livestock. Do not apply less than 28 days before harvest.
	(Bollworm,	to	to		
	Corn earworm,	0.4	16.0		
	Tobacco budworm)				
	Armyworms (Such as: Beet, Fall, Southern, Yellow-striped, etc.)				
	Green cloverworm				
	Velvetbean caterpillar				
	Woolly bear caterpillar				
	Soybean looper	0.45	18.0		
	Cabbage looper	to	to		
	Mexican bean beetle	0.75	30.0		
	Bean leaf beetle				
	Stinkbugs (Suppression)				
	Three-Cornered alfalfa hopper (Suppression)				
Cutworms	See Below		See "SPECIAL DIRECTIONS ON CUTWORMS"		

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Fluid Ounces		
SWEET CORN (Florida Fresh Market Only)	Corn earworm European corn borer Armyworms (Such as Fall, Beet, etc) Western bean cutworm	0.5 to 0.75	20 to 30	<p><u>Whorl feeders:</u> Spray as needed. As insect populations increase and/or when new plant growth is rapid, the spray interval should be shorter. For best results use sprays of 15 gallons or more per acre applied by ground equipment using solid or hollow cone nozzles directed into the plant whorl.</p> <p><u>Insects attacking silks and ears:</u> Apply at 1-7 day intervals starting when silks first appear and continuing until harvest, if necessary. As insect populations increase and/or silk growth is rapid, the spray interval should be shorter. Use the higher dosage rate and apply at 1 to 2 day interval following periods of heavy moth flights. Follow local recommendations for proper timing.</p> <p>Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground.</p> <p><u>Foliage Feeders:</u> Use at least 2 gallons finished spray volume per acre by air or 10 gallons per acre by ground.</p>	<p>After silk initiation, do not exceed 7.5 pounds of active ingredient (300 fluid ounces of this product) per acre per season.</p> <p>Do not allow livestock to graze treated field.</p> <p>Do not feed treated corn silage (green plant) or fodder to livestock.</p> <p>Ears may be harvested on day of last application.</p>
Cutworms	See Below			See "SPECIAL DIRECTIONS ON CUTWORMS."	

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

SPECIAL DIRECTIONS ON CUTWORMS

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Fluid Ounces		
CROPS INDICATED ABOVE	Cutworms	0.5 to 0.75	20.0 to 30.0	<u>Postemergence rescue treatments:</u> For best results, apply as a broadcast spray when cutworms are actively feeding. If cutworms are feeding below the ground, control will be more difficult. Light cultivation with a rotary hoe before spraying may enhance control. If banding, use a minimum width of 10 inches over the row. When banding, determine the amount of chemical to use per acre, by dividing the band width by the row width and multiplying by the appropriate broadcast rate. Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground.	Refer to each specific crop for use restrictions.

LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88

FIELD USE CONVERSION TABLE

DOSAGE PER ACRE		ACRES TREATED PER VOLUME OF LARVIN® brand 3.2		
Pounds Active	Fluid Ounces	1 gallon	2.5 gallons	5 gallons
0.125	5.0	25.6	64.0	128.0
0.25	10.0	12.8	32.0	64.0
0.33	13.2	9.7	24.2	48.5
0.40	16.0	8.0	20.0	40.0
0.45	18.0	7.1	17.7	35.5
0.50	20.0	6.4	16.0	32.0
0.55	22.0	5.9	14.5	29.0
0.60	24.0	5.3	13.2	26.5
0.65	26.0	4.9	12.3	24.6
0.70	28.0	4.6	11.4	22.8
0.75	30.0	4.2	10.5	21.4
0.80	32.0	4.0	10.0	20.0
0.90	36.0	3.6	8.9	17.8
1.00	40.0	3.2	8.0	13.0

1 U.S. Gallon Equals 128 Fluid Ounces

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experiences. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES

**LARVIN® brand 3.2 Thiodicarb Insecticide/Ovicide-PENDING-2/4/88**

ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**NOTICE TO THE BUYER**

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.