(PROPOSED LABEL)

with comment

WEEDONE® CB Woody Plant Herbicide

364-39

FOR CONTROLLING WOODY PLANTS IN FENCEROWS, ON ROADSIDES, RAILROADS, UTILITY RIGHTS-OF-WAY AND OTHER NON-CROP AREAS.

ACTIVE INGREDIENTS:

<pre>2-(2,4-Dichlorophenoxy)propionic acid, butoxyethanol ester* 2,4-Dichlorophenoxyacetic acid, butoxyethanol ester**</pre>			
INERT INGREDIENTS		•	76.4%

- *2-(2,4-Dichlorophenoxy)propionic acid equivalent 8.2% by weight or 0.66 pounds per gallon.***
- ** 2,4-Dichlorophenoxyacetic acid equivalent 8.2% by weight or 0.66 pounds per gallon.***

***Isomer specific by ADAC method No. 6-D01-5.

EPA Reg. No. 264-393

EPA Est No. 264-MO-1

Keep Out Of Reach of Children WARNING See Side Panel For Additional Precautionary Statements

IN CASE OF EMERGENCY
PHONE COLLECT (24 HOURS A DAY)
IN U.S.A. (304) 744-3487

CUNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
P. O. Box 12014, T.W. Alexander Drive, Research Triangle Park, N.C. 27709

GENERAL INFORMATION

WEEDONE® CB Woody Plant Herbicide is a ready-to-apply formulation for controlling unwanted woody plants by basal, stump or stem application. WEEDONE CB Woody Plant Herbicide is applied directly to the woody plant stem; it is not necessary to cut or frill the bark. WEEDONE CB Woody Plant Herbicide will not freeze under normal storage or application conditions, so it can be applied during the growth period or in the dormant season.

For most efficient treatment with WEEDONE CB Woody Plant Herbicide use the nozzles and spray equipment suggested in the Directions For Use.

During the growing season, vegetation near treated stems may be discolored and even killed. WEEDONE CB Woody Plant Herbicide normally allows ground cover plants to become reestablished in the treated area.

Species controlled by WEEDONE CB Herbicide include: chestnut oak ironwood (Ostrya) red oak ironwood (Carpinus) (blue beech) white oak American elm bur oak red elm water oak hackberry pin oak sour orange black oak redcedar bear (scrub) oak persimmon red maple* spruce ash* Sassafras sweetgum tallowtree blackgum sumac hickory aspen birch black locust White pine

balsam fir
Virginia pine*
southern pines*
red pine
sycamore

black cherry*
mazzard cherry*
choke cherry
poison ivy
Ailanthus
osageorange

pin cherry

^{*}Treat at least lower 36 inches of stem as specified in Directions For Use.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly. Apply this product only as specified on this label.

Do not dilute WEEDONE CB Woody Plant Herbicide with water. Do not apply WEEDONE CB Woody Plant Herbicide when stems are wet. Rainfall which follows application of WEEDONE CB to dry stems does not normally reduce brush control.

Do not graze meat or dairy animals on treated area.

FOR CONTROL OF WOODY PLANTS IN FENCEROWS, ON ROADSIDES, RAILROADS, UTILITY RIGHTS-OF-WAY AND OTHER NON-CROP AREAS

Average application volume of WEEDONE CB Woody Plant Herbicide is 5 gallons per acre, although actual amount used will depend on the amount and size of brush in the treated area. Total spray volume should be approximately 10% of that required for a conventional basal treatment. Do not exceed a total application volume of 20 gallons WEEDONE CB Woody Plant Herbicide per acre.

BASAL BARK TREATMENT:

To control general mixed brush up to 4 inches DBH (trunk diameter at breast height) including oak, hickory, sweetgum, blackgum, and many others, apply WEEDONE CB Woody Plant Herbicide undiluted to the lower 18-24 inches of the plant stem. Treat all around the stem including the root collar area and any exposed roots. It is not necessary for WEEDONE CB Woody Plant Herbicide to run down or puddle in the root collar area.

To control conifers larger than 2 inches DBH, or cherry, ash and maple, increase treatment height to cover the lower 36 inches of stem bark.

WEEDONE CB Woody Plant Herbicide basal bark treatment will control top growth of root-suckering species such as aspen, sumac, Sassafras, persimmon and black locust, but regrowth from lateral roots may occur the following growing season.

STUMP TREATMENT:

Control stems larger than 4 inches DBH, by cutting the tree, leaving an 18-24 inch tall stump and treating the sides and root collar area of the remaining stump.

If other oil-miscible woody plant herbicides are mixed with WEEDONE CB Woody Plant Herbicide to increase the spectrum of species controlled, do not exceed the ratio of 1 quart per 3 quarts WEEDONE CB Woody Plant Herbicide. Mix herbicide with WEEDONE CB Woody Plant Herbicide thoroughly before application.

APPLICATION EQUIPMENT

WEEDONE CB Woody Plant Herbicide can be applied from a hand pump back pack sprayer or a hand-carried pump-up pressurized sprayer such as Solo 425, or Chapin Model 152.

For most efficient use of the spray mixture, apply with a gun such as the Spraying Systems Model 23L-7676 equipped with a Spraying Systems 5500 Y series adjustable ConeJet nozzle. Equip all nozzles with a 200 mesh screen.

An alternate system which will reduce but not eliminate dripping between trigger releases is a 200 mesh TeeJet strainer and check valve number 4193A-SS having a spring opening that operates at 10 psi with the Spraying Systems 5500 Y series adjustable ConeJet nozzle mounted on a conventional basal spray wand.

Since WEEDONE CB Woody Plant Herbicide is applied as a concentrate, the most efficient nozzle size will vary with the diameter of the brush stems to be treated:

-For average brush of 1-3 inches DBH, or for heavy stands of brush with most stems less than 1 inch DBH use the Spraying Systems 5500 adjustable ConeJet nozzle with a Y2 orifice.

- For scattered stands or large trees or brush use the Spraying Systems 5500 adjustable ConeJet nozzle with a Y3 orifice. Spray pressure of 15 psi is optimum for most use. Do not use spray pressure higher than 20 psi.

Precautionary Statements

Hazard to Humans and Domestic Animals WARNING
Harmful if swallowed. Causes skin irritation. Causes Eye Irritation. Do not get on skin or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly before eating or smoking.

Statement of Practical Treatment

In case of skin contact, wash with plenty of soap and water. Get medical attention if irritation develops.

In case of eye contact, flush with plenty of water.

If swallowed, drink promptly large quantities of milk, egg white, gelatin solution or if these are not available, drink large quantities of water.

Avoid alcohol. Get medical attention.

Do not induce vomiting.

Do not give anything by mouth to an unconscious person.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not graze meat or dairy animals on treated area.

Do not apply when weather conditions favor drift. Do not use the same spray equipment for other purposes unless thoroughly cleaned. Do not contaminate water used for domestic or irrigation purposes. Do not use in or near a greenhouse. Do not apply WEEDONE CB Woody Plant Herbicide directly to or permit spray mist to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers, or other desirable crop or ornamental plants which are susceptible to 2,4-D or 2,4-DP herbicides. Do not apply near susceptible, plants since very small quantities of the 2,4-D or 2,4-DP will cause severe injury during the growing or dormant periods. Crops contacted by WEEDONE CB Woody Plant Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Drift may be reduced by: (1) applying as near to the target as possible; (2) by decreasing the pounds of pressure at the nozzle tips; (3) by using nozzles which produce a coarse spray pattern; (4) by not applying when wind is blowing toward susceptible crops or valuable plants.

Storage and Disposal Statements

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides, or fungicides. Pesticides, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

Container <u>Disposal</u>
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

(Limited warranty and Disclaimer to be placed here)

WEEDONE is a registered trademark of Union Carbide Agricultural Products Company, Inc.

ConeJet and TeeJet are registered trademarks of Spraying Systems Company.

...:

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warranta (a) that this product conforms to the chemical description of the label. (b) that this product is reasonably it for the purposes set forth in the directions for use when it is used in accordance with such direction, and (c) that the directions, warnings and other statements on this tabel are based upon responsible experts evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues, and upon reports of field experience. Tests have not been made on all verieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED. WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to, and the Buyer shall be solely responsible for, any and all loss or damage which results from the use of this product in any manner which is inconsistent with the tabel directions, warnings or cautions.

INCONSISION WITH THE TABLET CHOISE, WAITINGS OF CAUTIONS
BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR
SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS,
LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE
OR HANDLING OF THIS PRODUCT, WHETHER OR NOT SUCH
LIABILITY IS BASED IN CONTRACT, NEGLIGENCE, STRICT
LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE
MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE
REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY
OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE
CLAIMED IN NO EVENT SHALL MANUFACTURER OR SELLER
BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL
DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS
PRODUCT.

II PHUBANIE COLOR	EPA REGISTRATION NO.	DATE OF ISSUANCE
U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS		FES 18 1889
REGISTRATION DIVISION (NH-567) MASHINGTON, L.C. 27467	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: THE STRATION	NAME OF PESTICIDE PRODUC	Ť
(Under the Lederal Insection by Lancing Court Rock interference of the Lancing Court Rock interference of the Lancing Court Rock interference of the Lancing Court Rock in		
AME AND ADDRESS OF HE DISTRANT IN ILIE ZIP		
г	7	
L	J	
NOTE: Changes in Tabeling fermula differing in substance submitted to and accepted by the Registration Division pri- moduct always refer to the above U.S. EPA registration nu-	or to use of the label in commerc	with this registration must be e. In any correspondence on the
On the basis of information furnished by the registrant, the the Federal Insecticide, Fungicide, and Rodenticide Act.	above named posticide is hereb	y Registered/Reregistered unit
A copy of the labeling accepted in connection with this Re	egistration/Reregistration is retu	uned herewith.
Registration is in no way to be construed as an indursement health and the environment, the Administrator, on his moti- icide in accordance with the Act. The acceptance of any n Act is not to be construed as giving the registrant a right by others.	on, may at any time suspend or come in connection with the regis	ancel the registration of a pesi tration of a product under this
ATTACHMENT IS APPLICABLE		
IGNATURE OF APPROVING OFFICIAL		DATE
		<u>i & 1983</u>

- 3. Similarive (5) contestor your final point of lambling refere you release the product for supposent. He set to the 1-73 Enclosure for a lictuar losality of that printed lambling.
- 4. Submit the data lister below, contractor in accordance with the proposed test guidelines referred to below, no liver that six months from the date of registration.
 - o Perferal magister, Voi. 43, 35, 163, pp. 3733 = 37401, Anguse 22, 1978. Austr invalueion 250 = 153.81=3.

If these conditions are not complian with, the regardention will be subject to cancellation in accordance with FICRA sec. 3(a). Your oftense for aniphent or the product constitutes acceptance of these conditions.

A stamped copy of the tabel is inclosed for year respects.

Richard F. Mountmort (M)
Product hundger (23)
Fungioids-Merals, to Meanon
Registration division (NS-757)

and tosure

.