



RHÔNE-POULENC AG COMPANY

Certification of Compliance Statement

I, being an authorized representative of Rhône-Poulenc Ag Company, certify that all containers of WEEDONE® CB Ready-To-Use Woody Plant Herbicide, EPA Reg. No. 264-393, produced by October 23, 1993, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by April 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Company Representative: Karen S. Shearer

Title: Registration Manager

Signature: Karen S. Shearer

Date: February 26, 1993

298

WEEDONE® CB brand READY-TO-USE

Woody Plant Herbicide

For Controlling Woody Plants In Forest Management Programs And On Fencerows, Roadsides, Drainage Ditchbanks, Railroads, Utility Rights-Of-Way and Other Non-Crop Areas.

ACTIVE INGREDIENTS:

2-(2,4-Dichlorophenoxy)propionic acid, butoxyethyl ester* 11.7%
2,4-Dichlorophenoxyacetic acid, butoxyethyl ester** 11.9%

INERT INGREDIENTS: 76.4%

*2-(2,4-Dichlorophenoxy)propionic acid equivalent 8.2% by weight or 2/3 pounds per gallon.

**2,4-Dichlorophenoxyacetic acid equivalent 8.2% by weight or 2/3 pounds per gallon.

*Isomer specific by AOAC method No. 6.DO1-5.

Contains Petroleum Distillates

EPA Reg. No. 264-393

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada amplamente.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: If patient is conscious and alert, give 2 to 3 glasses of water to drink. Do not induce vomiting. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

PRECAUTIONARY STATEMENTS

WARNING

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled, or absorbed through the skin. Causes skin and eye irritation. Do not get in eyes, on skin or clothing. Avoid breathing spray mist. When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. For aerial applicators in enclosed cockpits the applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove all soiled clothing as soon as possible and shower.

NOTE: (1) For containers over 1 gallon, but less than 5 gallons, persons engaged in pouring of this product must also wear coverage of chemical-resistant apron. (2) For containers of 5 gallons or more, a mechanical transfer system (probe and pump) must be used for transferring the contents of the container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect fish, aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift. Do not contaminate water used for domestic or irrigation purposes.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D or 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D or 2,4-DP pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

CLEANING OF EQUIPMENT: Do not use the same spray equipment for other purposes unless thoroughly cleaned. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply WEEDONE® CB brand Ready-To-Use Woody Plant Herbicide directly to or permit spray mist to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers, or other desirable crop or ornamental plants which are susceptible to 2,4-D or 2,4-DP herbicides.

Do not allow spray to contact susceptible plants since very small quantities of the 2,4-D or 2,4-DP will cause severe injury during the growing or dormant periods. Crops contacted by WEEDONE® CB brand Ready-To-Use Woody Plant Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction. Vapors from this product may injure highly sensitive plants already growing in the immediate vicinity or planted there two to three weeks after spraying.

Do not use in or near a greenhouse.

Drift may be reduced by: (1) applying as near to the target as possible; (2) by decreasing the pounds of pressure at the nozzle tips; (3) by using nozzles which produce a coarse spray pattern; (4) by not applying when wind is blowing toward susceptible crops or valuable plants.

Do not graze meat or dairy animals on treated area.

Do not dilute WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE with water.

Do not apply WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE when stems are wet. Rainfall which follows application of WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE to dry stems does not normally reduce brush control.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

AGRICULTURAL USE REQUIREMENTS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance.

It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted-entry interval, and notification to workers. For any requirements specific to your State, consult the agency in your state responsible for pesticide regulation.

Do not enter or allow agricultural workers on farms and forests to enter into treated areas during the restricted entry interval (REI) of 24 hrs. For early reentry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, wear eye protection, chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. During aerial application, human flaggers either must wear chemical-resistant headgear or an approved jacket in enclosed vehicles.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Wastes representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local procedures.

SPECIES CONTROLLED BY WEEDONE® CB BRAND READY-TO-USE WOODY PLANT HERBICIDE

Alanthus (tree-of-heaven)	<i>Ailanthus altissima</i>	Juniper, common	<i>Juniperus communis</i>
Alder	<i>Ainus sp.</i>	Laurel, sheep	<i>Kalmia angustifolia</i>
Apple seedlings	<i>Malus sp.</i>	Locust	<i>Robinia sp.</i>
Aralia (devils-walkingstick)	<i>Aralia spinosa</i>	Locust, black	<i>Robinia pseudoacacia</i>
Ash	<i>Fraxinus sp.</i>	Magnolia (Sweet bay)	<i>Magnolia virginiana</i>
Ash, green	<i>Fraxinus pennsylvanica</i>	*Maple	<i>Acer sp.</i>
*Ash, white	<i>Fraxinus americana</i>	*Maple, bigleaf	<i>Acer macrophyllum</i>
Aspen	<i>Populus sp.</i>	*Maple, red	<i>Acer rubrum</i>
Aspen, big-tooth	<i>Populus grandidentata</i>	*Maple, sugar	<i>Acer saccharum</i>
Aspen, quaking	<i>Populus tremuloides</i>	Mulberry	<i>Morus sp.</i>
Balm-of-Gilead	<i>Populus gileadensis</i>	Oak	<i>Quercus sp.</i>
Bayberry, northern	<i>Myrica pennsylvanica</i>	Oak, bur	<i>Quercus macrocarpa</i>
Beautybush	<i>Kolkwitzia amabilis</i>	Oak, chestnut	<i>Quercus prinus</i>
Beech	<i>Fagus sp.</i>	Oak, northern red	<i>Quercus rubra</i>
Birch	<i>Betula sp.</i>	Oak, pin	<i>Quercus palustris</i>
*Birch, gray	<i>Betula populifolia</i>	Oak, southern red	<i>Quercus falcata</i>
Birch, paper	<i>Betula papyrifera</i>	Oak, water	<i>Quercus nigra</i>
*Birch, river	<i>Betula nigra</i>	Oak, white	<i>Quercus alba</i>
Blackberry	<i>Rubus sp.</i>	Orange, sour	<i>Citrus aurantium</i>
Boxelder	<i>Acer negundo</i>	Osageorange	<i>Maclura pomifera</i>
Buckeye	<i>Aesculus sp.</i>	Persimmon, common	<i>Diospyros virginiana</i>
Buckeye, Ohio	<i>Aesculus glabra</i>	*Pine, shortleaf **	<i>Pinus echinata</i>
Butternut	<i>Juglans cinerea</i>	*Pine, Virginia **	<i>Pinus virginiana</i>
Briars	<i>Rosa sp.</i>	Pine, jack	<i>Pinus banksiana</i>
Camphor tree	<i>Cinnamomum camphor</i>	Pine, red	<i>Pinus resinosa</i>
Catalpa, western	<i>Catalpa speciosa</i>	Pine, white	<i>Pinus strobus</i>
Cherry	<i>Prunus virginiana</i>	Poison-ivy	<i>Rhus radicans</i>
*Cherry, black	<i>Prunus serotina</i>	Poison-oak	<i>Rhus toxicodendron</i>
Cherry, pin	<i>Prunus pennsylvanica</i>	Poplar	<i>Populus sp.</i>
Chokecherry, common	<i>Prunus virginiana</i>	Poplar, balsam	<i>Populus balsamifera</i>
Cottonwood	<i>Populus sp.</i>	Prickly-ash	<i>Zanthoxylum americanum</i>
Dogwood	<i>Cornus sp.</i>	Prickly pear	<i>Opuntia sp.</i>
Elder	<i>Sambucus sp.</i>	Raspberry	<i>Rubus sp.</i>
Elder, Pacific red	<i>Sambucus callicarpa</i>	Redcedar, eastern	<i>Juniperus virginiana</i>
Elm, American	<i>Ulmus americana</i>	Rhododendron, Canadian	<i>Rhododendron canadense</i>
Elm, Chinese	<i>Ulmus parvifolia</i>	Rose	<i>Rosa sp.</i>
Elm, red	<i>Ulmus fulva</i>	Rose, multiflora	<i>Rosa multiflora</i>
Elm, winged	<i>Ulmus alata</i>	Sassafras	<i>Sassafras albidum</i>
Fir	<i>Abies sp.</i>	Serviceberry (shadbush)	<i>Amelanchier sp.</i>
Fir, Balsam	<i>Abies balsamea</i>	Sourwood	<i>Oxydendrum arboreum</i>
Fir, Douglas	<i>Pseudotsuga menziesii</i>	Spruce, white	<i>Picea glauca</i>
Grape	<i>Vitis sp.</i>	Sumac	<i>Rhus sp.</i>
Grape, Fox	<i>Vitis labrusca</i>	Sumac, staghorn	<i>Rhus typhina</i>
Gum, black	<i>Nyssa sylvatica</i>	Sumac, poison	<i>Rhus vernix</i>
Gum, sweet	<i>Liquidambar styraciflua</i>	Swamp cyrilla	<i>Cyrilla racemiflora</i>
Hackberry	<i>Celtis occidentalis</i>	Sweetfern	<i>Comptonia peregrina</i>
Hardhack	<i>Spiraea tomentosa</i>	Sycamore, American	<i>Platanus occidentalis</i>
Hawthorn	<i>Crataegus sp.</i>	Tallowtree, Chinese	<i>Sapium sebiferum</i>
Hawthorn, English	<i>Crataegus oxyacantha</i>	Tulip, poplar (yellow)	<i>Liriodendron tulipifera</i>
Hazel, beaked	<i>Corylus cornuta</i>	Walnut	<i>Juglans sp.</i>
Hickory	<i>Carya sp.</i>	Walnut, black	<i>Juglans nigra</i>
Hickory, pignut	<i>Carya glabra</i>	Wax myrtle	<i>Myrica cerifera</i>
Hickory, shagbark	<i>Carya ovata</i>	(southern bayberry)	
Honeysuckle	<i>Lonicera japonica</i>	Willow	<i>Salix sp.</i>
Hornbeam	<i>Carpinus sp.</i>	Willow, Bebb	<i>Salix bebbiana</i>
Huckleberry	<i>Vaccinium sp.</i>	Witchhazel	<i>Hamamelis sp.</i>
Ironwood	<i>Ostrya virginiana</i>	White-rod	<i>Virburnum cassinoides</i>
Ivy, English	<i>Hedera helix</i>		

*Treat at least the lower 36 inches of stem as specified in Directions For Use.

**Not effective on southern pines larger than 2 1/2" in stem diameter (DBH)

FLORIDA DRAINAGE DITCHBANK SPECIES CONTROLLED BY WEEDONE® CB BRAND READY-TO-USE WOODY PLANT HERBICIDE INCLUDE THE FOLLOWING:

Australian-pine	(<i>Casuarina equisetifolia</i>)	Lantana, largeleaf	(<i>Lantana camara</i>)
Brazilian peppertree	(<i>Schinus terebinthifolius</i>)	Sea Myrtle	(<i>Baccharis sp.</i>)
Cabbage palmetto	(<i>Sabal palmetto</i>)		

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE can be applied from a hand pump back pack sprayer or a hand-carried pump-up pressurized sprayer such as Chapin Model¹ 152, Sears 1443, or Birchmeier Iris² 10, Cooper Pegler Model CP 15³ or Solo⁴ models 425/475 (equipped with Atlas gaskets).

For most efficient use of the spray mixture, apply with a gun such as the B & G Extenda-wand equipped with a Spraying System 5500 Y series adjustable ConeJet⁵ nozzle. Equip all nozzles with a 200 mesh screen.

An alternate system which will reduce but not eliminate dripping between trigger releases is a 200 mesh TeeJet⁵ strainer and check valve number 4193A-SS having a spring opening that operates at 10 psi with the Spraying Systems 5500 Y series adjustable ConeJet nozzle mounted on a conventional basal spray wand.

Since **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE** is applied as a concentrate, the most efficient nozzle size will vary with the diameter of the brush stems to be treated:

For average brush of 1-3 inches DBH, or for heavy stands of brush with most stems less than 1 inch DBH use the Spraying Systems 5500 adjustable ConeJet nozzle with a Y2 orifice.

For scattered stands or large trees or brush or to treat large stumps, use the Spraying Systems 5500 adjustable ConeJet nozzle with a Y3 orifice.

For small stems less than 1 inch diameter or to limit amount of **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE** applied, use the Spraying Systems 5500 adjustable ConeJet nozzle with a Y1 orifice.

Spray pressure of 15 psi is optimum for most use. Do not use spray pressure higher than 20 psi.

EQUIPMENT MAINTENANCE

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE may affect O rings and other seals in certain sprayer units. The most effective method of minimizing this problem is to drain the sprayer completely every evening. **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE** is a concentrate and can be returned to the original container.

If daily applicator draining cannot be performed, prolonged exposure of specific wetted rubber parts, starting after two weeks, will result in reduced applicator reliability. Viton⁶, Nitrile Rubber, Buna N., Neoprene⁶ and EPDM are particularly susceptible to swelling after prolonged exposure to **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE**. Tests have shown Atlas-Poly (tetrafluoroethylene-co-propylene) to have excellent long-term resistance to **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE**. Solo Inc., Newport News Virginia, telephone (804) 245-4228, currently provides Atlas replacement gaskets (part # 4900337) for the Solo Model 425 Applicator. Other applicator manufacturers are currently evaluating other alternative rubbers having long-term resistance to **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE**.

To prevent time loss in the field, have available replacement parts whenever you start out to spray.

Sprayers can be cleaned of **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE** at the end of the season, or prior to shifting to other herbicides, by first rinsing with kerosene, then following with a triple rinse of hot water and detergent. At the end of the spray season replace all O rings and/or seals showing signs of deterioration. Such a maintenance program will prolong the life of your equipment.

GENERAL INFORMATION

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE is a ready-to-apply formulation that can be used in basal, stump and stem applications.

WEEDONE® CB is a ready-to-use product, applied undiluted, using the special application equipment described on the label. It controls over 100 major woody plant species, including oak, hickory, multiflora rose, sweetgum and blackgum when applied to the lower 18 to 24 inches of the plant stem. During the growing season, vegetation near treated stems may be discolored and even killed. **WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE** normally allows ground cover plants to become re-established in the treated area.

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE can be applied in both dormant and growing seasons. This year-round capability lengthens treatment programs and improves crew utilization.

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE is ready-to-apply and requires no mixing with oil, thus reducing application time.

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE will not freeze under normal storage or use conditions.

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE gives effective control of numerous unwanted woody plant species.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly.

FOR CONTROLLING WOODY PLANTS IN FOREST MANAGEMENT PROGRAMS AND ON FENCE-ROWS, ROADSIDES, DRAINAGE DITCHBANKS, RAILROADS, UTILITY RIGHTS-OF-WAY AND OTHER NON-CROP AREAS.

Average application volume of WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE is 3 gallons per acre, although actual amount used will vary from 1 to 5 gallons depending on the amount and size of brush in the treated area. Total spray volume should be approximately 10% of that required for a conventional basal treatment.

BASAL BARK TREATMENT

To control general mixed brush up to 4 inches DBH (trunk diameter at breast height) including oak, hickory, sweetgum, blackgum and many others, apply WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE undiluted to the lower 18 to 24 inches of the plant stem. Treat all around the stem including the root collar area and any exposed roots. It is not necessary for WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE to run down or puddle in the root collar area.

To control Southern pines less than 2 1/2 inches DBH, increase treatment height to cover the lower 36 inches of stem bark.

To control ash, cherry or maple increase treatment height to cover the lower 36 inches of stem bark.

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE basal bark treatment will control top growth of root-suckering species such as aspen, sumac, Sassafras, persimmon and black locust, but regrowth from lateral roots may occur the following growing season. Fall and winter treated stems may briefly leaf out in the following spring before final kill. WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE will not usually control woody stems larger than 4 inches DBH. For large stems, cut the tree and treat the stump as directed in the section on stump treatment.

STUMP TREATMENT

Control stems larger than 4 inches DBH, by cutting the tree and thoroughly treating the bark and root collar area of the remaining stump. The stump can be cut at any height - low or high as long as the remaining wood, bark and root collar area is thoroughly treated.

Cutting the stump as close to the ground as possible will reduce the amount of WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE needed for control.

Stumps do not require immediate treatment permitting separate cut and spray operations for greater efficiency.

FENCEROW TREATMENT

To control brush in fencerows apply WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE to brush or stumps as directed above. Do not allow WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE spray or drift to contact crops, forage plants or ornamentals.

DRAINAGE DITCHBANK TREATMENT

WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE effectively removes unwanted brush along ditchbanks that interferes with the growth of bank-holding perennial grasses. WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE is an excellent management tool, allowing you to selectively remove unwanted brush while leaving desired species. Do not allow spray or spray drift to contact water.

To control brush on drainage ditchbanks apply WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE to brush or stumps as directed above. Do not allow WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE spray or spray drift to contact crops, forage plants or ornamentals.

TANK MIXTURES WITH OTHER HERBICIDES

When mixing WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE with other herbicides, follow all directions, precautions and limitations on both product labels.

WEEDONE® CB/Garlon⁷ 4

To improve control on (resistant) difficult-to-control species such as red maple, white ash and black cherry, apply a mixture of 7 parts WEEDONE® CB brand WOODY PLANT HERBICIDE and 1 part Garlon 4 Herbicide (7 pints + 1 pint). Cover the lower 18" to 24" of brush stem thoroughly as with other WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE applications.

Add the required amount of Garlon 4 to the WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE in the sprayer tank. Shake or stir the sprayer to thoroughly mix the two ingredients. Once mixed, the Garlon 4 and WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE tank mixture should not need additional agitation.

WEEDONE® CB/Tordon⁷ K

To improve control of pine and root-suckering species such as sassafras, black locust or sumac, apply a mixture of 7 parts WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE and 1 part Tordon® K Herbicide. Cover the lower 18" to 24" of brush stem thoroughly as with other WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE applications.

Add the required amount of Tordon® Herbicide to the WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE in the sprayer tank. Shake or stir the sprayer to thoroughly mix the two ingredients and continue to agitate the sprayer every few minutes while spraying. WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE tank mixtures with Tordon® K Herbicide will separate on standing. Use entire spray mixture, do not allow this tank mixture to remain in the sprayer overnight. Do not apply this tank mixture to drainage ditch banks.

798

WEEDONE® CB/Banvel® OS

To improve control of pines, ash, persimmon, osageorange and certain other difficult-to-control species, apply a mixture of 7 parts WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE and 1 part Banvel OS Herbicide. Cover the lower 18" to 24" of brush stem thoroughly as with other WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE applications.

Add the required amount of Banvel OS to the WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE in the sprayer tank. Shake or stir the sprayer to thoroughly mix the two ingredients. Once mixed, the Banvel OS and WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE tank mixture should not need additional agitation.

Banvel Herbicide can be substituted for an equal amount of Banvel OS, but a suitable emulsifier or compatibility agent must be added to enable the Banvel Herbicide to disperse in the WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE. Test suitable mixtures before using. Banvel Herbicide/WEEDONE® CB brand READY-TO-USE WOODY PLANT HERBICIDE tank mixtures will separate on standing, do not allow this tank mixture to remain in the sprayer overnight or for any prolonged period without agitation. Do not apply this tank mixture to drainage ditchbanks.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warning and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

¹ Chapin is a registered trademark of R.E. Chapin Mfg. Works, Inc.

² Birchmeier Iris is a registered trademark of Birchmeier & Company.

³ CP-15 is a registered trademark of Cooper Pegler and Company, Ltd.

⁴ Solo 425 is a registered trademark of Solo Incorporated.

⁵ ConeJet and TeeJet are registered trademarks of Spraying Systems Company.

⁶ Viton and Neoprene are registered trademarks of E.I. duPont de Nemours and Company and Copolymer Rubber and Chemical.

⁷ Garlon and Tordon are registered trademarks of DowElanco.

⁸ Banvel is a registered trademark of Sandoz Corporation.

Rhône-Poulenc Ag Company

P.O. Box 12014, 2 T.W. Alexander Drive

Research Triangle Park, North Carolina 27709

WEEDONE is a registered trademark of Rhône-Poulenc

WEEDONE® CB Ready-to-Use Woody Plant Herbicide (PENDING) Submitted 2/26/93.