


PREP™
PLANT REGULATOR FOR COTTON



FOR COMMERCIAL USE OR AGRICULTURAL USE ONLY.

NOT FOR RESIDENTIAL USE.

ACTIVE INGREDIENT:

Ethephon (2-Chloroethyl) phosphonic acid* 55.4%

INERT INGREDIENTS: 44.6%

*This product contains 6 pounds ethephon per gallon.

EPA Reg. No. 264-418

EPA Est. No. 264-PA-01

KEEP OUT OF REACH OF CHILDREN

DANGER

PRACTICAL TREATMENT

IF IN EYES: Flush with water for at least 15 minutes and call a physician.
IF ON SKIN OR CLOTHING: Remove clothing and wash skin thoroughly with soap and water. Get medical attention.


FOR ADDITIONAL PRECAUTIONARY STATEMENTS: SEE SIDE PANEL

For Emergency Information ONLY Call 24 Hours A Day 1-800-521-8235

For Product Handling, Use and General Information Call 1-800-334-9745

The use of this product for a variety of plant growth regulation uses is protected by various United States and foreign patents including U.S. Patents 3,879,188 and 4,240,819. No license is granted to use this product in countries other than the United States or for any use not contemplated by this label. Liability for patent infringement may result from use or sale of this product outside the United States.

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
P. O. Box 12014, T.W. Alexander Drive
Research Triangle Park, North Carolina 27709



MADE IN U.S.A.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CORROSIVE. Causes eye and skin damage. May be fatal if absorbed through skin. Harmful if swallowed or inhaled. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear full length trousers, long sleeved shirt, protective gloves, and goggles or face shield when handling. Change to clean work clothes daily. Wash thoroughly with soap and water after handling and before eating, or smoking. Remove and wash contaminated clothing before reuse. Keep away from domestic animals.

STATEMENT OF PRACTICAL TREATMENT

Remove the patient from immediate source of exposure and assure that the individual is breathing. Use artificial respiration if necessary, to assure continued breathing. Call a physician in cases of suspected poisoning. If poisoning is suspected in animals, call a veterinarian.

IF IN EYES: Hold eyelids open and flush with a gentle stream of water for at least 15 minutes. Get medical attention, preferably an ophthalmologist.

IF ON SKIN: Immediately wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

NOTE TO PHYSICIAN

There is no specific antidote.


All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Victims of severe overexposure, by inhalation, should be kept under medical observation for up to 72 hours for delayed onset of pulmonary edema. In a victim of overexposure by ingestion, careful gastric lavage is required due to the possibility of stomach or esophageal perforation. This material is an acid, but use of alkaline substances to neutralize it is contraindicated.

ENVIRONMENTAL HAZARDS

Do not apply PREP™ Plant Regulator for cotton through any type of irrigation system.

Do not apply directly to water. Do not contaminate water used for irrigation or domestic purposes.



USE PRECAUTIONS

Avoid spray drift to nearby crops as this product will cause modifications in plant growth. Plant injury or reduced yields will result.

Do not plant another crop within 30 days after treatment.

Mix only the amount of spray you expect to use each day. Do not allow mixed solution to stand over night.

GENERAL INFORMATION

A foliar spray of PREP™ Plant Regulator will accelerate opening of mature unopened cotton bolls and enhance defoliation which can result in earlier harvest with an increased recoverable yield. PREP™ Plant Regulator treatment allows increased efficiency from a once-over harvest.

Some premature drop of small, immature bolls may be associated with treatment. Do not apply before sufficient, mature, unopened bolls have developed to produce the desired yield of cotton.

A boll is mature when it is too hard to be dented when squeezed between thumb and fingers, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.


For best results, thorough and uniform spray coverage is required. It is essential that the bolls are contacted by the spray to obtain optimum boll opening. Observe treated fields closely. Harvest at optimum boll opening and defoliation, generally 14 to 21 days after treatment. Sustained cool weather with average day and night temperatures of 65°F or lower after treatment with PREP™ Plant Regulator will delay the boll opening response.

DO NOT APPLY PREP™ PLANT REGULATOR IF RAIN IS EXPECTED WITHIN 6 TO 8 HOURS. Rainfall within 6 to 8 hours of application may reduce product performance.

Certified cotton seed growers should observe boll and seed maturity before treatment with PREP™ Plant Regulator. Bolls not completely mature might open with treatment of PREP™ Plant Regulator giving lower quality seed.

Spray deposits of PREP™ Plant Regulator, when left on aircraft acrylic plastic type windshields for longer than 1 hour, may react with this material, forming an opaque film or spots on the surface which cannot be removed by washing with water or any other cleaner. Rinse windshields with water immediately after each tankload of PREP™ Plant Regulator has been applied. Thoroughly rinse the aircraft with clean water after each day PREP™ Plant Regulator is applied.

PREP™ Plant Regulator may also affect certain paints, and if accidental deposits on painted objects such as cars, trucks, agricultural equipment, etc. occurs, these objects should be rinsed with water as soon as possible (within 1 hour) to prevent permanent staining of the painted surface.



SPRAY PREPARATION

Add 1/2 to 3/4 of the required amount of water to the spray tank. Start agitation. Add the required amount of PREP™ Plant Regulator, and the remaining amount of water. Prepare only as much spray solution as can be used on the day of mixing. Do not allow spray solution to stand overnight. Do not spill the concentrated product on spray equipment, or any airplane parts. ANY SPILLS SHOULD BE RINSED IMMEDIATELY WITH PLENTY OF WATER.

Use of a nurse tank is highly recommended for avoiding possible spills of concentrated formulation on spray equipment or any airplane parts.

EQUIPMENT CLEANING

Because of the acidic nature of this product, prolonged exposure to spray deposit will damage acrylic plastics, certain paints, and metals.

Rinse thoroughly with a detergent and water all exposed acrylic plastic-type materials (e.g., aircraft windshields), and painted surfaces within an hour after exposure to spray deposits.

At the end of each day, rinse thoroughly with a detergent and water all the metal parts of the aircraft and the associated spray equipment exposed to the spray deposits.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply PREP™ Plant Regulator through any type of irrigation system.

BOLL OPENING

Rates For Boll Opening

CONDITIONS EXPECTED 5-7 DAYS AFTER TREATMENT	PREP™ PINTS/ACRE	ONE GALLON OF PREP™ TREATS	SPRAY VOLUME GAL./ACRE	
			GROUND	AIR
Hot and dry 80°F or higher	1 1/3 (1.0 lb a.i.)	6 Acres	15 to 50	2 to 5*
Dry with temperature 75°F to 80°F	2 (1.5 lb a.i.)	4 Acres	15 to 50	2 to 5*
Cool but higher than 65°F	2 2/3 (2.0 lb a.i.)	3 Acres	15 to 50	2 to 5*
Rank cotton	2 2/3 (2.0 lb a.i.)	3 Acres	25 to 50	5

*NOTE: For California and Arizona use a volume of no less than 5 gallons per acre for aerial applications

Time of Application (Boll Opening)

Apply when there are sufficient mature unopened bolls to produce the desired yield, but not later than when 60% of the bolls are opened (see General Information section on how to test for boll maturity). Do not tank mix PREP™ Plant Regulator with a desiccant if the cotton is to be spindle harvested.

PRECONDITIONING FOR DEFOLIATION

Rates For Preconditioning For Defoliation

CONDITIONS EXPECTED 5-7 DAYS AFTER TREATMENT	PREP™ PINTS/ACRE	ONE GALLON OF PREP™ TREATS	SPRAY VOLUME GAL./ACRE	
			GROUND	AIR
Hot and dry 80°F or higher	2/3 (0.5 lb a.i.)	12 Acres	15 to 50	2 to 5*
Cool but higher than 65° and also for rank cotton	1 1/3 (1.0 lb a.i.)	6 Acres	15 to 50	2 to 5*

*NOTE: For California and Arizona use a volume of no less than 5 G.P.A. for aerial applications.


Time of Application (Preconditioning)

Apply PREP™ Plant Regulator 4 to 7 days prior to application of defoliant or tank mixed with the defoliant (see section on Compatibility). Do not tank mix PREP™ with desiccants unless plant desiccation is required.

Pre-treatment With Defoliants Prior to PREP™ Plant Regulator

If the cotton is overly rank or laying down in the middles and good spray coverage of the bolls with PREP™ is difficult, a pretreatment with defoliants will be useful to improve boll coverage with PREP™. Use dosage rates of PREP™ Plant Regulator recommended for boll opening. Read and observe all appropriate label use directions and precautions for the defoliant used.

Do not use a defoliant before there are sufficient mature unopened bolls to produce the desired yield (see General Information section on How to Test for Boll Maturity).



COMPATIBILITY

PREP™ Plant Regulator is compatible with DEF, FOLEX, DROPP, HARVADE, METHYL PARATHION, GUTHION AND MALATHION, and may be applied in sequence or as a tank mixture (DO NOT TANK MIX WITH DESICCANTS IF COTTON IS TO BE SPINDLE HARVESTED). In some cases slight reduction in boll opening response has been observed when tank mixes with defoliants were used.

Good agitation in the spray tank is essential and a tank mixture should not be allowed to stand without agitation for more than 5 to 10 minutes. Read and observe all appropriate label use directions and precautions for the defoliants and insecticides used.

PRECAUTION

DO NOT TANK MIX PREP™ PLANT REGULATOR WITH DEFOLIANTS CONTAINING SODIUM CHLORATE BECAUSE THIS RESULTS IN THE FORMATION OF HYPOCHLOROUS ACIDS WHICH UPON HEATING EMIT TOXIC CHLORINE FUMES.

WHEN TO HARVEST

Do not harvest cotton sooner than 7 days after a treatment with PREP™ Plant Regulator. Observe the treated crop and harvest when optimum boll opening has been reached. Too early harvest might reduce the full advantage of the treatment and too late a harvest may result in reduced quality and loss of lint which will drop from the plant.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food, or feed by storage or disposal.


If container is broken or contents have spilled, follow all precautions indicated above and clean up immediately. Before cleaning up, put on full length trousers, long sleeved shirt, protective gloves, and goggles or face shield. Soak up spill with absorbent media such as sand, earth or other suitable material and dispose of waste at an approved waste disposal facility.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent) the empty containers. Then offer for recycling, or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.




LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warning and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.



ACCEPTED
SEP - 3 1987
Under Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 264-418

FOR COMMERCIAL USE OR AGRICULTURAL USE ONLY.

NOT FOR RESIDENTIAL USE.

ACTIVE INGREDIENT:

Ethephon (2-Chloroethyl) phosphonic acid* 55.4%

INERT INGREDIENTS: 44.6%

*This product contains 6 pounds ethephon per gallon.

EPA Reg. No. 264-418

EPA Est. No. 264-PA-01

KEEP OUT OF REACH OF CHILDREN

DANGER

PRACTICAL TREATMENT

IF IN EYES: Flush with water for at least 15 minutes and call a physician.
IF ON SKIN OR CLOTHING: Remove clothing and wash skin thoroughly with soap and water. Get medical attention.

FOR ADDITIONAL PRECAUTIONARY STATEMENTS: SEE SIDE PANEL

For Emergency Information ONLY Call 24 Hours A Day 1-800-521-8235

For Product Handling, Use and General Information Call 1-800-334-9745

The use of this product for a variety of plant growth regulation uses is protected by various United States and foreign patents including U.S. Patents 3,879,188 and 4,240,819. No license is granted to use this product in countries other than the United States or for any use not contemplated by this label. Liability for patent infringement may result from use or sale of this product outside the United States.

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
P. O. Box 12014, T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

MADE IN U.S.A.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CORROSIVE. Causes eye and skin damage. May be fatal if absorbed through skin. Harmful if swallowed or inhaled. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear full length trousers, long sleeved shirt, protective gloves, and goggles or face shield when handling. Change to clean work clothes daily. Wash thoroughly with soap and water after handling and before eating, or smoking. Remove and wash contaminated clothing before reuse. Keep away from domestic animals.

STATEMENT OF PRACTICAL TREATMENT

Remove the patient from immediate source of exposure and assure that the individual is breathing. Use artificial respiration if necessary, to assure continued breathing. Call a physician in cases of suspected poisoning. If poisoning is suspected in animals, call a veterinarian.

IF IN EYES: Hold eyelids open and flush with a gentle stream of water for at least 15 minutes. Get medical attention, preferably an ophthalmologist.

IF ON SKIN: Immediately wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

NOTE TO PHYSICIAN

There is no specific antidote.

All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Victims of severe overexposure, by inhalation, should be kept under medical observation for up to 72 hours for delayed onset of pulmonary edema. In a victim of overexposure by ingestion, careful gastric lavage is required due to the possibility of stomach or esophageal perforation. This material is an acid, but use of alkaline substances to neutralize it is contraindicated.

ENVIRONMENTAL HAZARDS

Do not apply PREP™ Plant Regulator for cotton through any type of irrigation system.

Do not apply directly to water. Do not contaminate water used for irrigation or domestic purposes.

USE PRECAUTIONS

Avoid spray drift to nearby crops as this product will cause modifications in plant growth. Plant injury or reduced yields will result.

Do not plant another crop within 30 days after treatment.

Mix only the amount of spray you expect to use each day. Do not allow mixed solution to stand overnight.

GENERAL INFORMATION

A foliar spray of PREP™ Plant Regulator will accelerate opening of mature unopened cotton bolls and enhance defoliation which can result in earlier harvest with an increased recoverable yield. PREP™ Plant Regulator treatment allows increased efficiency from a once-over harvest. Some premature drop of small, immature bolls may be associated with treatment. Do not apply before sufficient, mature, unopened bolls have developed to produce the desired yield of cotton.

A boll is mature when it is too hard to be dented when squeezed between thumb and fingers, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.

For best results, thorough and uniform spray coverage is required. It is essential that the bolls are contacted by the spray to obtain optimum boll opening. Observe treated fields closely. Harvest at optimum boll opening and defoliation, generally 14 to 21 days after treatment. Sustained cool weather with average day and night temperatures of 65°F or lower after treatment with PREP™ Plant Regulator will delay the boll opening response.

DO NOT APPLY PREP™ PLANT REGULATOR IF RAIN IS EXPECTED WITHIN 6 TO 8 HOURS. Rainfall within 6 to 8 hours of application may reduce product performance.

Certified cotton seed growers should observe boll and seed maturity before treatment with PREP™ Plant Regulator. Bolls not completely mature might open with treatment of PREP™ Plant Regulator giving lower quality seed.

Spray deposits of PREP™ Plant Regulator, when left on aircraft acrylic plastic type windshields for longer than 1 hour, may react with this material, forming an opaque film or spots on the surface which cannot be removed by washing with water or any other cleaner. Rinse windshields with water immediately after each tankload of PREP™ Plant Regulator has been applied. Thoroughly rinse the aircraft with clean water after each day PREP™ Plant Regulator is applied

PREP™ Plant Regulator may also affect certain paints, and if accidental deposits on painted objects such as cars, trucks, agricultural equipment, etc. occurs, these objects should be rinsed with water as soon as possible (within 1 hour) to prevent permanent staining of the painted surface.

4 a 7

SPRAY PREPARATION

Add 1/2 to 3/4 of the required amount of water to the spray tank. Start agitation. Add the required amount of PREP™ Plant Regulator, and the remaining amount of water. Prepare only as much spray solution as can be used on the day of mixing. Do not allow spray solution to stand overnight. Do not spill the concentrated product on spray equipment, or any airplane parts. ANY SPILLS SHOULD BE RINSED IMMEDIATELY WITH PLENTY OF WATER.

Use of a nurse tank is highly recommended for avoiding possible spills of concentrated formulation on spray equipment or any airplane parts.

EQUIPMENT CLEANING

Because of the acidic nature of this product, prolonged exposure to spray deposit will damage acrylic plastics, certain paints, and metals.

Rinse thoroughly with a detergent and water all exposed acrylic plastic-type materials (e.g., aircraft windshields), and painted surfaces within an hour after exposure to spray deposits.

At the end of each day, rinse thoroughly with a detergent and water all the metal parts of the aircraft and the associated spray equipment exposed to the spray deposits.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply PREP™ Plant Regulator through any type of irrigation system.

BOLL OPENING

Rates For Boll Opening

CONDITIONS EXPECTED 5-7 DAYS AFTER TREATMENT	PREP™ PINTS/ACRE	ONE GALLON OF PREP™ TREATS	SPRAY VOLUME GAL./ACRE	
			GROUND	AIR
Hot and dry 80°F or higher	1 1/3 (1.0 lb a.i.)	6 Acres	15 to 50	2 to 5*
Dry with temperature 75°F to 80°F	2 (1.5 lb a.i.)	4 Acres	15 to 50	2 to 5*
Cool but higher than 65°F	2 2/3 (2.0 lb a.i.)	3 Acres	15 to 50	2 to 5*
Rank cotton	2 2/3 (2.0 lb a.i.)	3 Acres	25 to 50	5

*NOTE: For California and Arizona use a volume of no less than 5 gallons per acre for aerial applications

Time of Application (Boll Opening)

Apply when there are sufficient mature unopened bolls to produce the desired yield, but not later than when 60% of the bolls are opened (see General Information section on how to test for boll maturity). Do not tank mix PREP™ Plant Regulator with a desiccant if the cotton is to be spindle harvested.

PRECONDITIONING FOR DEFOLIATION:

Rates For Preconditioning For Defoliation

CONDITIONS EXPECTED 5-7 DAYS AFTER TREATMENT	PREP™ PINTS/ACRE	ONE GALLON OF PREP™ TREATS Acres	SPRAY VOLUME GAL./ACRE	
			GROUND	AIR
Hot and dry 80°F or higher	2/3 (0.5 lb a.i.)	12 Acres	15 to 50	2 to 5*
Cool but higher than 65° and also for rank cotton	1 1/3 (1.0 lb a.i.)	6 Acres	15 to 50	2 to 5*

*NOTE: For California and Arizona use a volume of no less than 5 G.P.A. for aerial applications.

Time of Application (Preconditioning)

Apply PREP™ Plant Regulator 4 to 7 days prior to application of defoliant or tank mixed with the defoliant (see section on Compatibility). Do not tank mix PREP™ with desiccants unless plant desiccation is required.

Pre-treatment With Defoliants Prior to PREP™ Plant Regulator

If the cotton is overly rank or laying down in the middles and good spray coverage of the bolls with PREP™ is difficult, a pretreatment with defoliants will be useful to improve boll coverage with PREP™. Use dosage rates of PREP™ Plant Regulator recommended for boll opening. Read and observe all appropriate label use directions and precautions for the defoliant used.

Do not use a defoliant before there are sufficient mature unopened bolls to produce the desired yield (see General Information section on How to Test for Boll Maturity).

COMPATIBILITY

PREP™ Plant Regulator is compatible with DEF, FOLEX, DROPP, HARVADE, METHYL PARATHION, GUTHION AND MALATHION, and may be applied in sequence or as a tank mixture (DO NOT TANK MIX WITH DESICCANTS IF COTTON IS TO BE SPINDLE HARVESTED). In some cases slight reduction in boll opening response has been observed when tank mixes with defoliant were used.

Good agitation in the spray tank is essential and a tank mixture should not be allowed to stand without agitation for more than 5 to 10 minutes. Read and observe all appropriate label use directions and precautions for the defoliant and insecticides used.

PRECAUTION

DO NOT TANK MIX PREP™ PLANT REGULATOR WITH DEFOLIANTS CONTAINING SODIUM CHLORATE BECAUSE THIS RESULTS IN THE FORMATION OF HYPOCHLOROUS ACIDS WHICH UPON HEATING EMIT TOXIC CHLORINE FUMES.

WHEN TO HARVEST

Do not harvest cotton sooner than 7 days after a treatment with PREP™ Plant Regulator. Observe the treated crop and harvest when optimum boll opening has been reached. Too early harvest might reduce the full advantage of the treatment and too late a harvest may result in reduced quality and loss of lint which will drop from the plant.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food, or feed by storage or disposal.

If container is broken or contents have spilled, follow all precautions indicated above and clean up immediately. Before cleaning up, put on full length trousers, long sleeved shirt, protective gloves, and goggles or face shield. Soak up spill with absorbent media such as sand, earth or other suitable material and dispose of waste at an approved waste disposal facility.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent) the empty containers. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

7 97

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warning and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.