

411 19 264-420 1076

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A) 	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460 Application for Pesticide:	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number <div style="font-size: 24pt; text-align: center;">196635</div>
----------------	---	---	---

Section I

1. Company/Product Number <div style="text-align: center; font-size: 18pt;">264-420</div>	2. EPA Product Manager Dennis Edwards	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SEVIN brand Carbaryl Insecticide Spray Ready-To-Use	PM#	
5. Name and Address of Applicant (Include ZIP Code) Rhone-Poulenc Ag Company 2 T. W. Alexander Drive RTP, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION: As required by PR Notice 93-3, the Environmental Hazards section of this label has been amended to prohibit the application.

NOTIFICATION
 LABEL NOT REVIEWED
 PER PR NOTICE 93-3
 DATE 6/29/93

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted.		If "Yes," Unit Package wgt. No. per container	If "Yes," Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label Is Affixed To Product				<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other (_____)	

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name <div style="text-align: center; font-size: 18pt;">Larry R. Hodges</div>	Title <div style="text-align: center; font-size: 18pt;">Registration Manager</div>	Telephone No. (Include Area Code) <div style="text-align: center; font-size: 18pt;">919-549-2870</div>
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped) <div style="text-align: center; font-size: 24pt;"> </div>
2. Signature 	3. Title <div style="text-align: center; font-size: 18pt;">Registration Manager</div>	<div style="text-align: center; font-size: 24pt;"> </div>
4. Typed Name <div style="text-align: center; font-size: 18pt;">Larry R. Hodges</div>	5. Date <div style="text-align: center; font-size: 18pt;">June 24, 1993</div>	

2874

SEVIN[®] brand Carbaryl Insecticide Spray Ready-To-Use

ACTIVE INGREDIENT: Carbaryl (1-naphthyl N-methylcarbamate)0.126%

INERT INGREDIENTS:99.874%

This product contains the toxic inert ingredient formaldehyde.

E.P.A. Reg. No. 264 - 420

E.P.A. Est. No. 264-

KEEP OUT OF REACH OF CHILDREN CAUTION CUIDADO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USER** Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.** If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash skin thoroughly with soap and water.

NOTE TO PHYSICIAN

Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

MAY BE HARMFUL IF SWALLOWED. Avoid breathing spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or disposal of wastes.

BEE CAUTION

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.


PLANT RESPONSE PRECAUTIONS

Do not use on Boston ivy, Virginia creeper and Maidenhair fern as injury will result. During early season, it may also injure Virginia and Sand pines.

DO NOT PLANT ROTATIONAL FOOD AND FEED CROPS NOT LISTED ON THIS OR OTHER CARBARYL LABELS IN CARBARYL TREATED SOIL.

NOTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 864

DATE: 6/29/93



216

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING

Shake container before using. Hold sprayer 8 to 12 inches from the area to be treated. Spray until surface is wet.

STORAGE AND DISPOSAL

STORAGE

Store unused product in an area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F.

PESTICIDE AND CONTAINER DISPOSAL

Do not reuse empty container. Rinse thoroughly before discarding in trash.

Securely wrap original container in several layers of newspaper and discard in trash.

VEGETABLE CROPS


For control of all the vegetable pests listed below, apply this product in adequate volume to effectively cover both the upper and lower surfaces of the plant.

NOTE: THIS PRODUCT IS NOT REGISTERED FOR USE ON CELERY AND SWEET POTATOES IN CALIFORNIA.

DO NOT TREAT ASPARAGUS MORE THAN ONCE EVERY 3 DAYS AS EXCESSIVE RESIDUES MAY RESULT

THE NUMBER IN PARENTHESIS FOLLOWING EACH VEGETABLE CROP GROUPING REFERS TO THE MINIMUM INTERVAL THAT MUST BE OBSERVED BETWEEN THE DATE OF THE LAST APPLICATION AND THE DATE OF HARVEST.

CROPS	PESTS
<p>Beans (including Black-eyed peas, Crowder or Southern peas, Dry beans, Green beans, Lima beans, Navy beans, and Snap beans), Carrots, Corn, Cucumber, Eggplant, Melons, Okra, Peanuts, Pepper, Potato, Pumpkin, Squash, Sweet Potato, Tomato (0 days until harvest)</p> <p>Asparagus (1 day until harvest)</p> <p>Broccoli, Brussels sprouts, Cabbage, Cauliflower, Cowpeas, Garden beet roots, Head Lettuce, Horseradish, Kohlrabi, Parsnips, Peas, Radishes, Rutabagas, Salsify roots, Turnip roots (3 days until harvest)</p> <p>Chinese cabbage, Celery, Collards, Dandelion, Endive (Escarole) Garden beet tops, Hanover salads, Kale, Mustard greens, Leaf lettuce, Parsley, Salsify tops, Spinach, Swiss chard, Turnip tops (14 days until harvest)</p>	<p>Alfalfa caterpillar</p> <p>Apache cicada</p> <p>Armyworms</p> <p>Asparagus beetle</p> <p>Aster leafhopper</p> <p>Bean leaf beetle</p> <p>Blister beetles</p> <p>Chinch bug</p> <p>Colorado potato beetle</p> <p>Corn earworm</p> <p>Corn rootworm adult</p> <p>Cowpea curculio</p> <p>Cucumber beetle</p> <p>Cutworms</p> <p>European corn borer</p> <p>Fall armyworm</p> <p>Flea beetles</p> <p>Grasshoppers</p> <p>Green cloverworm</p> <p>Harlequin bug</p> <p>Imported cabbageworm</p> <p>Japanese beetle</p> <p>Lace bugs</p> <p>Leafhoppers</p> <p>Lima bean pod borer</p> <p>Lygus bugs</p> <p>Melonworm</p> <p>Mexican bean beetle</p> <p>Pea leaf weevil</p> <p>Pea weevil</p> <p>Rednecked peanutworm</p> <p>Sap beetles</p> <p>Southwestern corn borer</p> <p>Spittlebugs</p> <p>Squash bugs</p> <p>Stink bugs</p> <p>Sweet potato weevil</p> <p>Tarnished plant bug</p> <p>Three cornered alfalfa hopper</p> <p>Thrips</p> <p>Tomato hornworm</p> <p>Tomato pinworm</p> <p>Tortoise beetle</p> <p>Velvetbean caterpillar</p> <p>Webworms</p> <p>Western bean cutworm</p> <p>Whitefringed beetle adults</p> <p>Yellowstriped armyworm</p>



FRUIT CROPS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward upper and lower leaf surfaces and between fruit clusters, and on small limbs and trunks where pests appear.

NOTE: TO AVOID UNDESIREED APPLE THINNING, DELAY USE UNTIL AT LEAST 30 DAYS AFTER FULL BLOOM.

CARBARYL INSECTICIDE MAY INJURE EARLY DAWN AND SUNRISE STRAWBERRY VARIETIES ON THE DELMARVA PENINSULA.

THE NUMBER IN PARENTHESIS FOLLOWING EACH FRUIT AND NUT CROP GROUPING REFERS TO THE MINIMUM INTERVAL THAT MUST BE OBSERVED BETWEEN THE DATE OF THE LAST APPLICATION AND THE DATE OF HARVEST.

TREE FRUITS	SMALL FRUITS	PESTS	
Apples, Cherries, Peaches, Pears, Plums, Prunes (1 day until harvest) Apricots, nectarines (3 days until harvest) Citrus (5 days until harvest)	Blueberries, Grapes (0 days until harvest) Strawberries (1 day until harvest) Blackberries, Raspberries, Dewberries (7 days until harvest)	Apple Aphid Apple maggot Apple mealybug Apple pandemis Apple rust mite Apple sucker Avocado leafroller Bagworms Black cherry aphid Black scale Blueberry maggot Brown soft scale California orangedog California pearslug (pear sawfly) California red scale Cherry fruitworm Cherry maggot Citricola scale Citrus cutworm Citrus root weevil Citrus snow scale Codling moth Cucumber beetles Cutworms Eastern tent caterpillar Eightspotted forester European apple sawfly European earwig European raspberry aphid Eyespotted budmoth Forbes scale Fruittree leafroller Grape berry moth Grape leafhopper Grape leafroller Green fruitworm Japanese beetle June beetles Leafhoppers Leafrollers Lecanium scales	Lesser appleworm Lesser peachtree borer Meadow spittlebug Mealy plum aphid Omnivorous leaftier (Strawberry fruitworm) Omnivorous leafroller Orange tortrix Oriental fruit moth Oystershell scale Peach twig borer Pear leaf blister mite Pear psylla Pear rust mite Periodical cicada Plum curculio Prune leafhopper Raspberry sawfly Red-banded leafroller Rose chafer Rosy apply aphid Saltmarsh caterpillar San Jose scale Snowy tree cricket Strawberry weevil Tarnished plant bug Tentiform leafminers Tussock moth Variegated leafroller Western grapeleaf skeletonizer Western tussock moth Western yellowstriped armyworm West Indian sugarcane borer (adults) White apple leafhopper Woolly apple aphid Yellow headed fireworm

TREES, ORNAMENTALS, SHRUBS AND FLOWERS


This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward the upper and lower leaf surfaces and small trunks, stems and twigs where pests appear.

NOTE: THIS PRODUCT IS NOT REGISTERED FOR USE ON CELERY AND SWEET POTATOES IN CALIFORNIA.

DO NOT TREAT ASPARAGUS MORE THAN ONCE EVERY 3 DAYS AS EXCESSIVE RESIDUES MAY RESULT

THE NUMBER IN PARENTHESIS FOLLOWING EACH VEGETABLE CROP GROUPING REFERS TO THE MINIMUM INTERVAL THAT MUST BE OBSERVED BETWEEN THE DATE OF THE LAST APPLICATION AND THE DATE OF HARVEST.

TREES	PESTS		
Trees (including) Shade trees, Shelter belts, Plantations, Parks and Recreational areas Ornamentals (including) Roses, Other Herbaceous plants, Woody plants, Shrubs	Ants	Hackberry nipplegall maker	Rise aphid
	Apple aphid	Holly budmoth	Pose chafer
	Armyworm:	Holly leafminer	Roseslug
	Azalea leafminer	Jackpine budworm	Saddled prominent
	Bagworms	Japanese beetle	Sawflies (exposed)
	Birch leafminer	Jeffrey pine needleminer	Scale insects
	Blister beetle	June beetles	Sowbugs
	Boxelder bug	Plant bugs	Spiny elm caterpillar
	Boxwood leafminer	Lace bugs	Springtails
	Browntail moth	Leafhoppers	Spruce budworm
	Cankerworms	Leafrollers	Spruce needleminer
	Catalpa sphinx	Locust borer	Subtropical pine tip moth
	Chiggers	Maple leafcutter	Tent caterpillar
	Cooley spruce gall aphid	Mealybugs	Thorn bug
	Cutworms	Mimosa webworm	Thrips (exposed)
	Cypress tip moth	Nantucket pine tip moth	Ticks
	Douglas-fir tussock moth	Oak leafminers	Walnut caterpillar
	Eastern spruce gall aphid	Oak leaf skeletonizer	Webworms
	Elm leaf aphid	Oakworm complex	Western hemlock looper
	Elm leaf beetle	Oleander caterpillar	Western spruce budworm
	Elm spanworm	Olive ash borer	Willow leaf beetles
	Eriophyid mites	Orange-striped oakworm	Yellow poplar weevil
	European pine shoot moth	Orange tortrix	
	Fall armyworm	Periodical cicada	
	Flea beetles	Pine sawfly	
	Fuller rose beetles	Pine spittlebug	
	Gall midges	Pitch pine tip moth	
	Gall wasps	Poinsetta hornworm	
	Grasshoppers	Psyllids	
	Green striped mapleworm	Puss caterpillar	
	Gypsy moth	Redhumped oakworm	



PEST CONTROL AROUND BUILDINGS AND OTHER OUTDOOR AREAS

This product may be used as a spot treatment around buildings such as homes, apartments, warehouses, barns and municipal and recreational areas to control the pests listed below. Thoroughly wet the outside perimeter of dwellings and other areas where pests tend to congregate.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of SEVIN brand Carbaryl Insecticide Spray Ready-To-Use to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wallpaper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

PESTS	
Ants	Earwigs
Bees	Firebrats
Wasps	Silverfish
Brown dog ticks	Fleas
Carpenter ants	Millipedes
Centipedes	Scorpions
Cockroaches	Spiders
Crickets	

DISCLAIMER OF LIABILITY

BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

SEVIN® is a registered trademark of Rhône-Poulenc.

SEVIN® brand Carbaryl Insecticide Spray Ready-To-Use (MASTER) Notification 6/23/93.