

JUN 4 1967

Mr. J.S. Lovell
Union Carbide Agricultural Products Company, Inc.
T.W. Alexander Drive
P.O. Box 12014
Research Triangle Park, NC 27709

Dear Mr. Lovell:

Subject: Change in Forage Crops/Wasteland Directions for Use Added
Sevin Brand ULV Carbaryl Insecticide
EPA Registration No. 264-422
Your Application Date: September 3, 1966

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Dennis Edwards
Acting Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

91741:Edwards:186393:E-4:KENCO:3/10/87:3/18/87:kim:vo:tar

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

DIRECTIONS FOR USE (con't)

WASTELANDS, RIGHTS-OF-WAY, HEDGEROWS, DITCHBANKS, ROADSIDES INSECT CONTROL

For grasshoppers, black grass bugs, mormon cricket, range caterpillars, range crane fly: Apply 1/2 to 1 quart of this product per acre. Use the lower rate for nymphs and immature insects on small plants or sparse vegetation. Apply the higher rate for adults or when vegetation is thick.

Apply a maximum of 2 applications per year. Allow at least 14 days between applications. Do not allow foraging or cut for hay within 14 days of last application by ground. Aerially treated areas may be grazed or cut for hay on day of treatment.

Carefully mark swaths to avoid over-application.

IMPORTANT: Before using SEVIN® brand ULV Carbaryl Insecticide, read and carefully observe the Precautionary Statements and all other information appearing on the product label. This bulletin contains revised directions for use of this product which may not appear on the package label. Follow directions carefully.

OBSERVE ALL DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE FEDERAL LABEL FOR SEVIN® BRAND ULV CARBARYL INSECTICIDE.

August 1986

6672y/8-20-86

264-422-3/3

PRODUCT BULLETIN



UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
T. W. ALEXANDER DRIVE
P.O. BOX 12014
RESEARCH TRIANGLE PARK, NC 27709

SEVIN® brand ULV Carbaryl
Insecticide. EPA Reg. No. 264-422
Revised Directions for Use on:
Forage Crops: Alfalfa, Pastures,
Rangeland, Wasteland/Rights-of-
Way/Hedgerows/Ditchbanks/Roadsides

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

CAREFULLY CALIBRATE ALL APPLICATION EQUIPMENT.

ALFALFA INSECT CONTROL

For grasshoppers: Apply 1/2 to 1 1/2 quarts of this products per acre. To avoid possible injury to tender alfalfa foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.

Apply only once per cutting

Do not apply to alfalfa within 7 days of harvest.

PASTURE INSECT CONTROL

For grasshoppers, true armyworm, fall armyworm, chinch bugs, essex skipper, striped grass looper, thrips: Apply 1/2 to 1 quart of this product per acre. Use the lower rate for nymphs and immature insects on small plants or sparse vegetation. Apply the higher rate for adults or when vegetation is thick.

Apply a maximum of 2 applications per year. Allow at least 14 days between applications. Do not allow foraging or cut for hay within 14 days of last application by ground. Aerially treated pastures may be grazed or cut for hay on day of treatment.

Carefully mark swaths to avoid over-application.

RANGELAND INSECT CONTROL (FOR AERIAL APPLICATION ONLY)

For grasshoppers, black grass bugs, mormon cricket, range caterpillars, range crane fly: Apply 1/2 to 1 quart of this product per acre. Use the lower for nymphs and immature insects on small plants or sparse vegetation. Apply the higher rate for adults or when vegetation is thick.

Apply a maximum of 2 applications per year. Allow at least 14 days between applications.

Carefully mark swaths to avoid over-application

RECEIVED
with COMMENTS
In EPA Letter Dated

JUN 04 1987

6672y/8-20-86

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 264-422.