

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A)		United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number <h2 style="margin: 0;">188590</h2>	
		Application for Pesticide:			

Section I

1. Company/Product Number <h3 style="margin: 0;">264-426</h3>	2. EPA Product Manager <h3 style="margin: 0;">D. Edwards</h3>	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <h3 style="margin: 0;">TEMIK 15G brand Aldicarb Pesticide for California only</h3>		
5. Name and Address of Applicant (Include ZIP Code) Rhone-Poulenc Ag Company 2 T. W. Alexander Drive Research Triangle Park, NC 27707 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION: PR Notice 93-1 - Restricted Use
PR Notice 93-3 - Wetlands

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted.		If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Other (_____) <input type="checkbox"/> Stenciled					

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Warren A. Davis	Title Senior Registration Manager	Telephone No. (Include Area Code) 919-549-2379
--------------------------------	---	--

Certification		6. Date Application Received (Stamped) <div style="border: 1px solid black; height: 100px; width: 100%;"></div>
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		
2. Signature 	3. Title Senior Registration Manager	
4. Typed Name Warren A. Davis	5. Date May 26, 1993	

2 of 11

RESTRICTED USE PESTICIDE

ACUTE TOXICITY AND GROUND WATER CONTAMINATION

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

TEMIK[®] brand 15G ALDICARB PESTICIDE

For Sale and Use in California Only

For Control of Certain Insects, Mites, and Nematodes

ACTIVE INGREDIENT: Aldicarb [2-methyl-2-(methylthio) propionaldehyde O- (methylcarbamoyl)oxime].....15%

INERT INGREDIENTS:85%

EPA Reg. No. 264-426

EPA Est. No. 264-GA-01



KEEP OUT OF REACH OF CHILDREN DANGER POISON PELIGRO



PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

TRANSLATION TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything orally to an unconscious or convulsing person. Call a physician and follow General advice.

IF IN EYES: Flush eyes with plenty of water and get medical attention.

IF ON SKIN: Wash thoroughly with soap and water. Remove contaminated clothing and wash before reuse.

IF INHALED: Call a physician and follow General advice.

PLANT PROTECTIVE CORP
JUN 8 1993
DATE _____

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Illness may be produced rapidly following over-exposure to TEMIK[®] Aldicarb. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN**. If poisoning is suspected in animals, contact a veterinarian.

NOTE TO PHYSICIAN

TEMIK[®] brand aldicarb is a methyl carbamate insecticide which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. TEMIK[®] brand aldicarb poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of the administration of parenteral atropine sulfate. Caution should be exercised to prevent over-atropinization. Mild cases may be given 1 to 2 mg intra-muscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until the patient is fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs such as (pyridine-2-aldoxime methiodide) are **NOT** recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples must be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number: 1 (800) 334-7577.

BEST AVAILABLE COPY

3 7 11

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

FATAL IF SWALLOWED. Causes cholinesterase inhibition. May be fatal or harmful by skin or eye contact or by breathing dust. Rapidly absorbed through skin or eyes. Do not get on skin or in eyes. Do not breathe dust. Keep away from domestic animals. Wear long-sleeved clothing, full-length trousers, head coverings, and protective gloves when handling. Always stand up-wind from hopper when loading. Wash hands and face before eating, drinking, or using tobacco. Bathe at the end of work day, washing entire body and hair with soap and water. Change clothing daily. Wash contaminated clothing in strong washing soda solution and rinse thoroughly before reusing.

SIGNS AND SYMPTOMS OF OVEREXPOSURE

Salivation, Muscle tremor, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache.

In severe cases, convulsions, unconsciousness, and respiratory failure may occur.

ANTIDOTE STATEMENT: ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

ENVIRONMENTAL HAZARDS

DECOMPOSITION AND MOVEMENT IN SOIL: This product is readily decomposed into harmless residues under most use conditions. However, a combination of permeable and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F (10°C) at application time tend to reduce degradation and promote movement of residues to groundwater. If the above describes your local use conditions and groundwater in your area is used for drinking, do not use this product without first contacting Rhône-Poulenc at 1 (800) 334-9745.

TOXIC TO FISH, BIRDS, AND WILDLIFE: This product is toxic to fish, birds, aquatic invertebrates and other wildlife. Birds feeding on exposed granules in treated areas may be killed. Cover or incorporate granules in spill areas. Run-off from treated areas may be hazardous to fish in neighboring areas. For terrestrial uses, do apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated area without protective footwear. Persons coming in direct contact with wet treated soil after the initial irrigation or rainfall following treatment shall wear coveralls, rubber or neoprene boots, and gloves. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application. Regular long-sleeved shirt, full-length trousers, and head covering should be worn when working in treated fields. See PRECAUTIONARY STATEMENT, STATEMENT OF PRACTICAL TREATMENT, and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (DANGER-POISON), area treated with TEMIK® brand 15G Aldicarb Pesticide, date of application, appropriate protective clothing, and re-entry requirements.

STORAGE AND DISPOSAL

STORAGE

Store unused TEMIK® brand Aldicarb Pesticide in original container only, in well ventilated clean dry area out of reach of children and animals. Do not store in areas where temperature averages 115° F (46° C) or greater. Do not store in or around the home or home garden.

If container is broken, before cleaning up, put on long-sleeved shirt, full-length trousers, head covering and rubber gloves. Do not get dust or granules on skin or in eyes. Do not breathe dust. Always stand upwind from spill when cleaning up. Sweep up and bury any small spills or excess TEMIK® Aldicarb at least 18 inches deep in soil, isolated from water supplies and food crops.

PESTICIDE DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or The Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or as allowed by state and local authorities, by open burning. If container is burned, stay away from and do not breathe or contact smoke.

IN CASE OF TRANSPORTATION OR WAREHOUSE EMERGENCY INVOLVING A SPILL, FIRE, OR EXPOSURE, CALL 1 (800) 334-7577 TWENTY-FOUR HOURS A DAY IN THE USA.

BEST AVAILABLE COPY

GENERAL PRECAUTIONS

Use TEMIK® brand 15G Aldicarb Pesticide only in accordance with label directions, warnings, and precautions. DO NOT USE ON ANY CROP NOT LISTED ON THIS OR SUPPLEMENTAL LABELING AS ANY RESIDUES REMAINING MAY BE ILLEGAL OR HARMFUL. DO NOT STORE OR USE IN OR AROUND THE HOME OR HOME GARDEN. Use higher rates on heavy organic or clay soils. Do not exceed the maximum label rate. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crop, meat, and milk. Treatments in excess of 7 pounds per acre made directly in the seed furrow of cotton, sugar beets, soybeans, or drybeans may delay plant emergence and reduce plant stand. Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, TEMIK® brand 15G Aldicarb Pesticide and water must be on the same side of the plant row.

Calibrate and adjust application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. For any left over material, see instructions for STORAGE AND DISPOSAL in this booklet.

Deep disc any spills at row ends immediately to prevent birds from feeding on exposed granules.

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of groundwater contamination.

Do not apply in Del Norte or Humboldt counties in California.

PREHARVEST, GRAZING, MAXIMUM USE RATE, AND GROUNDWATER LIMITATIONS

TO AVOID EXCESSIVE RESIDUES IN OR ON:

COTTONSEED

- Apply between March 1 and September 1 only.
- Do not make more than one 'at-planting' application and one postemergence application per crop. Do not apply within 90 days of harvest.
- Do not feed cotton forage to livestock or allow livestock to graze in treated area.
- Do not exceed a total of 21 pounds per acre.

PEANUTS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not hog-off treated fields.
- Do not feed hay or vines to livestock.
- Do not exceed a total of 20 pounds per acre.

SUGAR BEETS

- Apply only between March 1 and September 1.
- Do not make more than one at-planting application and 2 postemergence applications per crop.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.
- Do not exceed a total of 28 pounds per acre.

SOYBEANS

- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay, or straw to livestock.
- Do not exceed a total of 20 pounds per acre.

DRY BEANS


- Apply only between March 1 and September 1.
- Do not make more than one application per crop.
- Do not harvest within 90 days of application.
- Do not feed green forage, hay or straw to livestock.
- Do not use green pods as food for humans.
- Do not exceed a total of 7 pounds per acre.

SWEET POTATOES

- Do not make more than one application per crop.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.
- Do not exceed a total of 20 pounds per acre.

SORGHUM

- Do not make more than one application per year.
- Do not harvest within 90 days of application.
- Do not feed green forage to livestock.
- Do not exceed a total of 7 pounds per acre.



ROTATIONAL CROPS RESTRICTION

- Do not plant any crop not listed on this label in soil treated with TEMIK® brand 15G Aldicarb Pesticide within 10 months after the last application with the following exceptions:
 - Do not plant wheat or barley within 6 months after last application.
 - Do not plant corn within 8 months of last application.
 - Do not plant melons within 8 months after last application.
 - Do not plant other cucurbits (such as cucumbers and squash) within 8 months after last application.
 - Do not plant bulb crops (such as onions or garlic) within 6 months after the last application.
 - Do not plant fruiting vegetables (such as tomatoes or eggplant) within 8 months after the last application.
 - Do not plant brassica crops (such as broccoli or cabbage) within 6 months after the last application.

MEAT AND MILK

- Do not allow livestock to graze in treated areas before harvest.

GROUNDWATER


- Observe Environmental Hazard precautions regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- Do not apply within 50 feet of any drinking water well.
- Do not wash, load, or empty application equipment near any well, as this practice is a potential source of groundwater contamination.
- In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.
- Do not apply in Del Norte or Humboldt counties in California.

COMPATIBILITY

Pesticidal activity of TEMIK® brand 15G Aldicarb Pesticide is not affected by normal applications of fertilizers or other pesticides. Its effectiveness may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix TEMIK® aldicarb with other materials before application.

GENERAL INFORMATION


TEMIK® brand 15G Aldicarb Pesticide controls certain insects, mites, and nematodes. When applied into moist soil at planting and/or postemergence the active ingredient aldicarb is rapidly absorbed by roots and translocated to all parts of the plant. Rainfall or irrigation soon after application will ensure prompt uptake of aldicarb. Control often lasts more than six weeks varying with growing conditions, rate of use, and pests. Crop yields are usually increased with treatments of TEMIK® brand 15G Aldicarb Pesticide.



DIRECTIONS FOR USE AND RECOMMENDED APPLICATIONS

To provide maximum performance and to minimize hazard to birds, work granules into or cover with soil to a depth of 2 inches or more. Deep disc any spills at row ends immediately to prevent birds from feeding on granules. When a range or rates are recommended, use the higher rate if pest infestations are expected to be severe.

Crop & Time Of Application	Pests Controlled	Pounds Per Acre	Ounces/1000 Feet of Row	Recommended Application
COTTON At Planting	Aphids, Thrips	3.5 to 5	<u>40" row spacing</u> 4.5 to 6	Drill granules just below seed line or place in seed furrow and cover with soil. If seed and TEMIK® are hill-dropped, TEMIK® rates may be reduced by one-half.
	Fleahoppers Leafminers, mites Overwintering boll weevil (adults feeding on foliage), Plant bugs, including Lygus	4 to 7	5 to 8.5	
At first squaring	Leafhoppers Fleahoppers	7 to 14	8.5 to 17	Side-dress granules 8 to 16 inches to one side of the plant row and 2 to 6 inches deep (usually at or below bottom of water furrow)
	Mites, Plant bugs, including Lygus	14	17	
From squaring through early bloom Do not exceed a total of 21 lbs/A.	Cotton leaf perforator Whiteflies	14	17	
GRAIN SORGHUM At Planting	Nematodes Greenbug Chinch Bug	7	<u>36" row spacing</u> 7.5	Apply granules in seed furrow and cover with soil.
PEANUTS At planting	Thrips	7 to 14	<u>36" row spacing</u> 7.5 to 15	Apply granules in seed furrow and cover with soil. In Southwest use high rate only.
	Nematodes	14 to 20	15 to 22	Apply granules in a 6 to 12 inch band and work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed into treated zone.




Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
SUGAR BEETS At planting or within one week before planting	Nematodes	14	22" row spacing 9.5	Apply granules in a 4 to 6 inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed in or above treated zone, OR Where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
At planting	Aphids	7 to 14	4.5 to 9.5	Drill granules 1 to 3 inches below seed line. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre. Repeat applications will be required for continued protection against virus vectors (aphids and leafhoppers).
	Leafminers Leafhoppers	14	9.5	
	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules in a 2 to 4 inch band over seed row and immediately work into the soil or cover the soil, OR Where furrow irrigation is employed for seed germination, drill granules 2 deep and 2 inches from seed row on water furrow side. For the 7 lb. rate, apply granules in a 1 to 2 inch band in front of the press wheel as the furrow is closing.
At planting plus postemergence (split applications)	Nematodes	14 at planting and 14 at postemergence	9.5 at planting and 9.5 at postemergence	AT PLANTING: Apply granules in a 4 to 6 inch band and immediately work into the soil or cover with soil to a depth of 2 to 4 inches. Plant seed into or above treated zone, OR Where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side. POSTEMERGENCE: Apply granules to both sides of plant row and immediately work into the soil or cover with soil, OR For furrow irrigation, side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 30 days after planting.
Postemergence Do not make more than one 'at-planting' application and two post-emergence applications per crop.	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil, OR For furrow irrigation, side-dress granules 4 to 8 inches to water furrow side of plant row and at furrow depth. Irrigate soon after application.
	Aphids	7 to 14	4.5 to 9.5	Apply as above. A repeat application may be required for continued protection against virus vectors (aphids, leafhoppers).
	Leafminers Leafhoppers	14	9.5	
	Nematodes	14	9.5	Apply as above. Apply within 30 days of planting

BEST AVAILABLE COPY

Crop & Time Of Application	Pests Controlled	Pounds/Acre	Ounces/1000 Feet of Row	Recommended Application
SOYBEANS At planting	Mexican bean beetle Thrips Suppression of three cornered alfalfa hopper	5 to 10	<u>30"-48" row spacing</u> 5.5 to 11	Drill granules 2 to 3 inches below seed line, OR Two to 3 inches to the side of the seed row and 2 to 3 inches deep. Granules can be applied in seed furrow if rate does not exceed 5 pounds per acre.
	Nematodes	10 to 20	11 to 22	Apply granules in a 6 to 8 inch band and immediately work into the soil or cover with soil. Plant seed into treated zone.}
DRY BEANS	Aphids	5 to 7	<u>22-48" row spacing</u> 5.5 to 7.5	Drill granules 2 to 3 inches below seed line, or 2 to 3 inches to side of seed row and 2 to 3 inches deep, OR Granules can be placed in seed furrow if rate does not exceed 5 lbs. per acre.
	Leafhoppers Mexican bean beetle, Mites	7	7.5	
	Nematodes	7	7.5	Apply granules in a 6-inch band and immediately work into the soil or cover with soil. Plant seed into treated zone. OR Where furrow irrigation is employed soon after planting, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
SWEET POTATOES At planting	Nematodes	10 to 20	<u>48" row spacing</u> 15 to 30	Apply granules in a 12-inch band in opened row. Cover immediately with soil by hilling up 8 to 10 inches. Plant in center of treated zone.

SEE PREHARVEST AND GRAZING LIMITATIONS ELSEWHERE IN THIS LABEL



TEMIK® BRAND 15G (GYPSUM) ALDICARB PESTICIDE CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS* FOR TEMIK® BRAND 15G ALDICARB PESTICIDE (GYPSUM) AT 4, 5 & 6 MPH

NOTE: This calibration chart is applicable only to TEMIK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a gypsum carrier. All rates are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Guide provided with tube). Calibration tubes are available from your TEMIK® brand 15G Aldicarb supplier.


TYPE OF GRANULAR APPLICATOR

POUNDS TEMIK® 15G PER ACRE FOR VARIOUS ROW SPACING

ROW SPACING					GANDY GAUGE SETTING			JOHN DEERE (7100) SETTING			NOBLE E-Z FLOW GAUGE SETTING			CLAMP CO 3/8" FEEDER DIAL RPM
					MPH			MPH			MPH			MPH
22"	34"	36"	40"	48"	4	5	6	4	5	6	4	5	6	5
1.8	1.2	1.1	1	.8	7	8	9	3**	3**	4**	2	2	3	3
3.6	2.4	2.2	2	1.7	10	11	12	4**	5**	6**	3	4	5	6
5.5	3.5	3.3	3	2.5	12	13	14	6**	7**	8	5	6	7	8
7.3	4.7	4.4	4	3.3	13	15	16	7**	9	10	6	7	8	11
9.1	5.9	5.5	5	4.2	14	16	18	9	11	13	7	9	10	14
10.9	7.1	6.7	6	5.0	16	18	19	11	13	15	8	10	12	16
12.7	8.2	7.8	7	5.8	17	19	21	12	15	18	9	11	13	18
14.5	9.4	8.9	8	6.7	18	20	22	14	17	20	10	12	14	21
16.4	10.6	10.0	9	7.5	19	21	23	16	19	23	11	13	15	23
18.2	11.8	11.1	10	8.3	20	22	24	17	21	25	12	15	16	26
21.8	14.1	13.3	12	10.0	22	24	26	20	25	28	14	16	19	31
25.5	16.5	15.6	14	11.7	23	26	29	24	28	30	16	18	21	37
29.1	18.8	17.8	16	13.3	25	28	31	26	29	32	17	20	23	42
32.7	21.2	20.0	18	15.0	26	30	33	28	31	33	19	21	24	48
36.4	23.5	22.2	20	16.7	28	31	34	30	32	35	20	23	26	54
43.6	28.2	26.7	24	20.0	31	34	37	32	34	37	22	26	29	65
49.1	31.8	30.0	27	22.5	33	36	40	33	36	38	24	28	31	74

* All settings for one hopper box outlet per row

** For John Deere gauge setting below 8 there will be great variations in flow rate. A gauge setting of 6 is the lowest recommended setting.



TEMK® BRAND 15G (CORNCOB) ALDICARB PESTICIDE CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS* FOR TEMK® BRAND 15G ALDICARB PESTICIDE (CORNCOB) AT 4, 5 & 6 MPH

NOTE: This calibration chart is applicable only to TEMK® brand 15G Aldicarb Pesticide in this container which is formulated for use on a corncob carrier. All rates are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Guide provided with tube). Calibration tubes are available from your TEMK® 15G Aldicarb supplier.


TYPE OF GRANULAR APPLICATOR

POUNDS TEMK® 15G PER ACRE FOR VARIOUS ROW SPACING

ROW SPACING					GANDY GAUGE SETTING			JOHN DEERE (7100) SETTING			NOBLE E-Z FLOW GAUGE SETTING			CLAMP CO 3/8" FEEDER DIAL RPM
22"	34"	36"	40"	48"	MPH			MPH			MPH			MPH
					4	5	6	4	5	6	4	5	6	5
1.8	1.2	1.1	1	.8	14	16	17	8	9	10	5	6	7	4
3.6	2.4	2.2	2	1.7	19	21	22	11	13	14	10	11	13	8
5.5	3.5	3.3	3	2.5	22	24	27	14	16	18	13	15	17	11
7.3	4.7	4.4	4	3.3	25	28	30	17	19	21	16	18	21	15
9.1	5.9	5.5	5	4.2	28	31	33	19	21	23	18	22	24	19
10.9	7.1	6.7	6	5.0	30	33	35	21	23	25	21	24	26	22
12.7	8.2	7.8	7	5.8	32	35	37	22	25	26	23	26	29	26
14.5	9.4	8.9	8	6.7	34	37	39	24	26	28	25	28	31	30
16.4	10.6	10.0	9	7.5	35	38	41	25	27	29	26	30	33	33
18.2	11.8	11.1	10	8.3	37	40	43	26	28	30	28	31	34	37
21.8	14.1	13.3	12	10.0	39	43	47	28	30	33	31	34	37	44
25.5	16.5	15.6	14	11.7	42	46	50	29	32	35	33	36	40	52
29.1	18.8	17.8	16	13.3	44	49	54	31	34	37	35	39	44	59
32.7	21.2	20.0	18	15.0	47	52	59	33	36	38	37	42		67
36.4	23.5	22.2	20	16.7	49	56	66	34	37	40	39	45		76
43.6	28.2	26.7	24	20.0	54	66		37	40	44	44			95
49.1	31.8	30.0	27	22.5	59			38	42					

*All settings for one hopper box outlet per row

**Odd Numbered Housing



11 9 11

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER OR NOT SUCH LIABILITY IS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof outside of the United States.

Rhône-Poulenc Ag Company
Post Office Box 12014, 2 T. W. Alexander Drive
Research Triangle Park, North Carolina 27709

TEMK is the registered trademark of Rhône-Poulenc.

TEMK® 15G /California ONLY (MASTER) Notification 5/24/93.