

SEVIN® brand Granular Carbaryl Insecticide

For Outdoor Home Insecticide Use
For Home Insecticide Use

ACTIVE INGREDIENT: Carbaryl (1-naphthyl N-methylcarbamate) 7.0% by wt.
INERT INGREDIENTS: 93.0% by wt.
E.P.A. Reg. No. 264 - 430 E.P.A. Est. No. 264-MO-02

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

Carbaryl is an N-Methyl Carbamate Insecticide.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN

Carbaryl is an N-methyl carbamate insecticide, which is a cholinesterase inhibitor. Atropine is antidotal. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

Consultation on diagnosis, therapy and analysis can be obtained at all hours by calling the emergency number.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (& DOMESTIC ANIMALS) CAUTION



HARMFUL IF SWALLOWED, ABSORBED THROUGH THE SKIN, INHALED, OR IF IN EYES.

Avoid breathing vapors or dust. Avoid contact with eyes, skin or clothing. Keep out of reach of children and domestic animals.

PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks and household or latex rubber gloves.

Follow manufacturer's instructions for cleaning and maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



SPECIFIC USE DIRECTIONS

TURFGRASS

For control of turfgrass pests, apply granules uniformly to dry turfgrass. For optimum control, follow specific directions for proper application and irrigation.

CROP	PEST	POUNDS OF SEVIN® GRANULAR PER 1000 SQ.FT.	SPECIFIC DIRECTIONS
Lawns, Recreational, and Ornamental Turf Areas	Ants Armyworm Centipedes Chiggers Chinch bugs Cutworms Earwigs Essex skipper European chafer Fall armyworm Fiery skipper Grasshoppers Green June beetle larvae Leafhoppers Lucerne moth Millipedes Mosquitoes (adults) Scarab beetle adults (such as May beetle, June beetle, Japanese beetle, green June beetle) Sod webworm (lawn moths) Sowbugs Spittlebugs Springtails Ticks Yellowstriped armyworm	2.2 to 3.0	For effective control, irrigate after application to remove chemical from granules. Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
	Billbug larvae European crane fly larvae Fleas White grubs (such as Japanese beetle larvae, Chafer beetle larvae, and <i>Phyllophaga</i> spp. larvae, and annual white grubs) Black turfgrass ataenius beetle larvae Hyperodes weevil (bluegrass weevil) larvae	3.0	For effective control of subsurface pests, apply when insects are actively feeding near the soil surface, usually during late March through May, or July through early September, or as recommended by local Agricultural Extension Service Agents. Irrigate thoroughly after application to remove chemical from granules and move it down into the thatch. Repeat applications as necessary up to 2 times per year but not more often than once every 7 days.

RESTRICTIONS AND LIMITATIONS: TURFGRASS

- DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION.


CONTROL OF TICKS WHICH VECTOR LYME DISEASE

For control of juvenile and adult *Ixodes* spp. ticks (Deer tick, Bear tick, and Black legged tick) and *Amblyomma* spp. ticks (Lone star tick). Apply granules at the rate of 2.5 to 3.0 pounds of formulated product per 1000 square feet.

Begin applications in late spring or early summer (May/June). Apply the recommended amount of product uniformly across the entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from granules.

IMPORTED FIRE ANTS

For use as a mound treatment to control imported fire ants, apply 1.5 to 3.0 ounces by weight over and around each fire ant mound. Thoroughly irrigate mound immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and causing migration before chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.



4 7 5

NUISANCE PEST CONTROL


For use in and around flower beds and ornamental plantings and around buildings.

PEST	POUNDS OF SEVIN® GRANULAR PER 1000 SQ.FT.	SPECIFIC DIRECTIONS
Ants Armyworms Ticks Centipedes Crickets Cutworms Earwigs Firebrats Fleas Millipedes Scorpions Silverfish Spiders	3.0	Observe plant response precautions. Apply granules in a 6 foot band around the building, to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules.

NUISANCE PEST CONTROL IN VEGETABLES AND IN SMALL FRUITS AND BERRIES (NOT FOR USE IN CALIFORNIA)

For control of Ants, Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, and Ticks, apply 0.75 pound per 1000 sq. ft. evenly over the soil surface. Water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season. Do not make more than 4 applications per year or apply more often than once every 7 days during the growing season. Do not use on crops other than those listed below. Observe the following preharvest intervals for applications during the growing season:

<u>PHI in Days</u>	<u>Crops</u>
1	Asparagus
2	Sweet Corn
3	Broccoli, Brussel sprouts, Cabbage, Cauliflower, Kohlrabi Fruiting Vegetables (including tomato, eggplant, pepper)
7	Root and Tuber Vegetables (including beets, carrots, parsnips, rutabagas, salsify, turnips, potatoes) Sweet potatoes Small Fruit and Berry crops (including grapes, strawberries, blackberries, dewberries, raspberries, blueberries)
14	Leafy Vegetables (including celery, dandelion, endive, lettuce, parsley, rhubarb, spinach, swiss chard) Chinese cabbage, Collards, Kale, Mustard Peanuts



5 9.5

NOTICE OF DISCLAIMER

BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

SEVIN® is a registered trademark of Rhône-Poulenc.

SEVIN® brand Granular Carbaryl Insecticide For Outdoor Home Insecticide Use (PENDING) Submitted 10/28/97.

