

M 23 264-445

ACCEPTED  
SEP 16 1991  
Director for the Federal Insecticide, Fungicide, and Rodenticide Act  
EPA Reg. No. 264-445

# CHIPCO® RONSTAR® G

## Herbicide

**NOT FOR USE ON HOME LAWNS**

**FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY**

Selective, preemergent herbicide for the control of annual grasses and broadleaf weeds in Turf, Ground Covers, Woody Ornamental Shrubs, Vines, and Trees.

### ACTIVE INGREDIENT:

Oxadiazon [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-Δ<sup>2</sup> 1,3,4-oxdiazolin-5-one] .....2.0%

### INERT INGREDIENT:

98.0%

## KEEP OUT OF REACH OF CHILDREN WARNING

EPA Reg. No. 264-445

EPA Est. No. 33560-TN-1

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call 1-800-334-9745.

### PRECAUTIONARY STATEMENTS:

#### WARNING

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye and skin irritation. Avoid contact with skin, eyes, or clothing. Avoid contamination of feed and food-stuffs. Keep opened containers away from children and animals. Wash clothing and equipment after handling this product.

#### STATEMENT OF PRACTICAL TREATMENT

**IF ON SKIN:** Wash with soap and water.

**IF IN EYES:** Flush with plenty of water. Get medical attention.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

#### GENERAL INFORMATION

CHIPCO® RONSTAR® G is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf and on a wide variety of ground covers, and woody ornamental shrubs, vines, and trees. CHIPCO® RONSTAR® G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply CHIPCO® RONSTAR® G with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

#### TURF

For use in established perennial Bahiagrass, Bluegrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as, fairways, parks, golf courses, and lawns. CHIPCO® RONSTAR® G controls Crabgrass (*Digitaria* spp.), Annual Bluegrass (*Poa*

*annua*), Goosegrass (*Eleusine indica*), Green Foxtail (*Setaria viridis*), Yellow Foxtail (*Setaria glauca*), Bristly Foxtail (*Setaria verticillata*), and Field Sandbur (*Cenchrus incertus*); also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis* spp.), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus officinis*), and Pigweed (*Amaranthus* spp.).

Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. Do not apply to wet turf. If rain is not expected shortly after application, sprinkling turf thoroughly will increase effectiveness. Do not use on Buffalograss in California.

\* Not for use in California.

#### AMOUNT TO USE

**Crabgrass, Goosegrass, Foxtail, and Field Sandbur Control:** For preemergent control, apply 100-200 pounds per acre (2 to 4 pounds active ingredient) or 2 1/4 to 4 1/2 pounds per 1000 sq. ft. of CHIPCO® RONSTAR® G in early spring prior to seed germination. Application made after March or April may not be fully effective.

**ANNUAL BLUEGRASS CONTROL:** For preemergent control, apply 200 pounds per acre (4 pounds active ingredient) or 4 1/2 pounds per 100 sq. ft. of CHIPCO® RONSTAR® G in late summer to early fall or in early spring prior to seed germination.

**Pigweed, Florida Pusley, Oxalis, Stinging Nettle, and Carpetgrass Control:** For preemergent control of these weeds, apply 200 pounds per acre (4 pounds active ingredient) or 4 1/2 pounds per 1000 sq. ft. of CHIPCO® RONSTAR® G in early spring prior to weed seed germination.

**Fall Application for Spring Weed Control in Cool Season Turf:** For preemergent control of crabgrass, goosegrass, and other annual weeds in Bluegrass, Perennial Ryegrass, and Tall Fescue Turf. Apply 150-200 lbs per acre (3 to 4 pounds active ingredient) or 3 3/8 to 4 1/2 pounds per 1000 square feet in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after spring green up.

**Annual Weed Control During Bermudagrass Establishment from Sprigs:** For preemergent control of

**BEST AVAILABLE COPY**

crabgrass, goosegrass and other labeled annual weeds during sprigging, apply 150 pounds per acre (3 pounds active ingredient) or 3 3/8 pounds per 1000 square feet of CHIPCO® RONSTAR® G immediately prior to or immediately after sprigging. Disturbance of the soil surface after CHIPCO® RONSTAR® G application will disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.

For St. Augustinegrass turf, do not exceed the rate of 150 pounds per acre (3 pounds active ingredient) of CHIPCO® RONSTAR® G.

CHIPCO® RONSTAR® G may cause discoloration of Bermuda-grass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

Avoid contact with flowers and shrubs except as recommended elsewhere on this label. Do not apply CHIPCO® RONSTAR® G to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. CHIPCO® RONSTAR® G should not be used on putting greens or tees because of the varying cultural practices used.

**GROUND COVERS, WOODY ORNAMENTAL SHRUBS, VINES, AND TREES APPLICATION:**

Apply CHIPCO® RONSTAR® G anytime during the year, prior to weed seed germination, for weed control. CHIPCO® RONSTAR® G can be applied to both newly transplanted and established ornamentals and ground covers. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of CHIPCO® RONSTAR® G will improve weed control activity. CHIPCO® RONSTAR® G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of CHIPCO® RONSTAR® G. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

**AMOUNT TO USE**

Ground Application - Apply CHIPCO® RONSTAR® G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 1/4 to 4 1/2 pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

CHIPCO® RONSTAR® G at 100 to 200 pounds (2.0 to 4.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broadleaf weeds such as:

**ANNUAL BROADLEAF WEEDS**

- Bittercress (*Cardamine spp.*)
- Carpetweed (*Mollugo verticillata*)
- Common Groundsel (*Senecio vulgaris*)
- Common Purslane (*Portulaca oleracea*)
- \*Evening Primrose (*Oenothera spp.*)
- \*Galinsoga (*Galinsoga spp.*)
- \*\*\*Garden Spurge (*Euphorbia hirta*)
- Golden Ragwort (*Senecio aureus*)
- Lambsquarters (*Chenopodium album*)
- \*\*Liverwort (*Marchantia spp.*)
- \*\*\*Niruri (*Phyllanthus niruri*)
- Pennsylvania Smartweed (*Polygonum pensylvanicum*)
- Petty Spurge (*Euphorbia pepus*)
- \*\*Prostrate Spurge (*Euphorbia supina*)
- Redroot Pigweed (*Amaranthus retroflexus*)
- \*\*\*Sensitive Plant (*Mimosa pudica*)
- \*\*\*Sow Thistle (*Sonchus oleraceus*)
- Speedwell (*Veronica spp.*)
- \*\*\*Spiny Amaranth (*Amaranthus spinosus*)
- Spotted Casuar (*Hypochaeris radicata*)
- \*Swinecress (*Coronopus didymus*)
- Yellow Woodsorrel (*Oxalis stricta*)

**ANNUAL GRASSES**


- Annual Bluegrass (*Poa annua*)
- \*Barnyardgrass (*Echinochloa crus-galli*)
- \*\*\*Bristly Foxtail (*Setaria verticillata*)
- Crabgrass (*Digitaria spp.*)
- Fall Panicum (*Panicum dichotomiflorum*)
- Field Sandbur (*Cenchrus incertus*)
- Green Foxtail (*Setaria viridis*)
- \*\*Goosegrass (*Eleusine indica*)
- \*\*Jungle Ricegrass (*Echinochloa colonum*)
- \*\*Panama Paspalum (*Paspalum fimbriaatum*)
- Yellow Foxtail (*Setaria glauca*)

\*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

\*\*Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds.

\*\*\*Not registered in California.

BEST AVAILABLE COPY



**How To Calibrate Spreaders For Applications Of CHIPCO® RONSTAR®**

	Ground Speed (MPH)	Rate CHIPCO® RONSTAR® G (lb) Per Acre			Effective Swath Width (in feet)
		100	150	200	
<b>CYCLONE</b> Spreader on Wheels Model B. 99 and 100 Electric & PTO. Model B3-PRO	One Trip	3.25	3.75	4.0	6.5
	Two Trips	2.5	3.0	3.25	6.5
	3.0	3	3.4	4	9
<b>EZEE FLOW</b> Model No. 105 Model No. 106 & 116  Model No. 109	-	D	E	-	4.5 or 4 1/2
	3.75	A-2	B-2	-	-
	5.0	B-2	-	C-2	-
	3.5	C-3	-	B-4	-
	5.0	A-4	B-4	C-4	-
<b>GANDY</b> 10S 10T Turf Tender	4	31	37	42	(6, 8, 10, 12, 20 FOOT) 3.5
	4	26	31	39	
	3.0	22	25	27	
<b>LESCO</b> Lesco	3.0	12/32"	14/32"	16/32"	9
<b>LELY</b> Model H Pro 425 RPM (Ring Arm in Hole I or II) Model W (Ring Arm in Hole I to II)	3	2	2.5	3	20
	4	-	3	-	
		4.5	2.5	3	
<b>SCOTT'S</b> Scott Drop Scott R7 -- [Models PR-1, PT-10, PT-11] [Models PF-1, PF-2, PF-3] Scott's Pro Turf (R-8A) Scott Rotary Seymour		3 1/4	4	4 3/4	3
				D	7
	3.0	4.2	4.7	5.7	1.67
	3.0	I	K	L	9
				5A	9
<b>SPYKER</b> Model 24-22		3.8	4.2	4.4/4.5	6
<b>VICON</b>  202-1002	3	18	22	27	20
	4	-	27	30	20
	5	24	-	36	20
	6	27	32	-	20

NOTE: These settings serve as guides only. Other settings may be necessary to conform to the condition of your equipment. for additional information on calibration, contact your Rhône-Poulenc sales representative.

## CROP TOLERANCE

CHIPCO® RONSTAR® G is recommended for use on a wide variety of ornamental species and ground covers in container, field grown, and landscape planting situations. CHIPCO® RONSTAR® G may be used at the recommended rates on the following species.

### ESTABLISHED ORNAMENTALS AND GROUND COVERS

COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME
Abutilon	<i>Abutilon pictum</i>	Citrus, Ornamental*:	
Acacia	<i>Acacia spp</i>	Lime	<i>Citrus aurantifolia</i> Bearss
Agapanthus	<i>Agapanthus africanus</i>	Tangelo	<i>Citrus XTangelo</i> Minneola
Allamanda, Common	<i>Allamanda cathartica</i>	Temple	<i>Citrus Xnobilis</i> Temple
Andromeda	<i>Pieris spp</i>	Cuban Shaddock	<i>Citrus maxima</i>
Aralia	<i>Aralia spinosa</i>	Valencia	<i>Citrus sinensis</i>
Arborvitae	<i>Thuja spp</i>	Coromandel	<i>Asystasia gangetica</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>	Cotoneaster	<i>Cotoneaster spp</i>
Azalea	<i>Rhododendron spp</i>	Crabapple*	<i>Malus spp</i>
Jamboo, Feathery	<i>Bambusa vulgaris</i>	Crapemyrtle	<i>Lagerstroemia indica</i>
Banyan, Weeping	<i>Ficus benghalensis</i>	Chrysanthemum	<i>Chrysanthemum spp</i>
Barberry	<i>Berberis spp</i>	Cypress	<i>Cupressus spp</i>
Beafsteak plant	<i>Acalypha wilkesiana</i>	Cypress, Leyland	<i>Cupressocyparis leylandii</i>
Bearberry	<i>Arctostaphylos uva-ursi</i>	Daphne, Fragrant	<i>Daphne odora</i>
Beech, European	<i>Fagus sylvatica</i>	Deutzia, Slender	<i>Deutzia gracilis</i>
Birch, Paper	<i>Betula papyrifera</i>	Dieffenbachia	<i>Dieffenbachia spp</i>
B. wood, Japanese	<i>Buxus microphylla</i>	Dogwood, Common	<i>Cornus florida</i>
Boxwood, Common	<i>Buxus sempervirens</i>	Dogwood, Redosier	<i>Cornus stolonifera</i>
Bougainvillea	<i>Bougainvillea spectabilis</i>	Dogwood, Kousa	<i>Cornus kousa</i>
Brassaia, Dwarf	<i>Brassaia actinophylla</i>	Douglas fir	<i>Pseudotsuga menziesii</i>
Buddleia	<i>Buddleia davidii</i>	Elm	<i>Ulmus spp</i>
Bugleflower	<i>Ajuga reptans</i>	Escallonia	<i>Escallonia spp</i>
Callistemon	<i>Callistemon lanceolatus</i>	Eucalyptus	<i>Eucalyptus spp</i>
Camellia, Common, Japanese	<i>Camellia japonica</i>	Euryops, Gray Leaved	<i>Euryops pectinatus</i>
Camellia, Sasanqua	<i>Camellia sasanqua</i>	Euonymus, Spreading	<i>Euonymus kiautschovicus</i>
Candytuft	<i>Iberis sempervirens</i>	Euonymus, Wintercreeper	<i>Euonymus fortunei</i>
Carissa	<i>Carissa macrocarpa</i>	Falsecypress, Hinoki	<i>Chaemaecyparis obtusa</i>
Carnation	<i>Dianthus caryophyllus</i>	Falsecypress, Japanese	<i>Chaemaecyparis pichera</i>
Ceanothus	<i>Ceanothus americanus</i>	Flowering Quince	<i>Chaenomeles speciosa</i>
Cedar, Red	<i>Juniperus virginia</i>	Forsythia	<i>Forsythia intermedia</i>
Cedar, Atlas	<i>Cedrus atlantica</i>	Gardenia	<i>Gardenia jasminoides</i>
Cistus	<i>Cistus purpureus</i>	Gazania, Trailing	<i>Gazania uniflora lucicola</i>
		Ginkgo	<i>Ginkgo biloba</i>
		Heath, Spring	<i>Erica carnea</i>
		Heather, Scotch	<i>Calluna vulgaris</i>

**BEST AVAILABLE COPY**

ESTABLISHED ORNAMENTALS AND GROUND COVERS (con't)

COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME
Heliotropium	<i>Heliotropium arborecens</i>	Oak	<i>Quercus spp</i>
Hemlock, Canada	<i>Tsuga canadensis</i>	Ochana	<i>Ochana serrulata</i>
Hibiscus	<i>Hibiscus spp</i>	Ohai	<i>Samanea saman</i>
Honeysuckle	<i>Lonicera spp</i>	Oleander	<i>Nerium oleander</i>
Holly	<i>Ilex spp</i>	Orthosiphon	<i>Orthosiphon stamineus</i>
Ice Plant	<i>Carpobrotus edule</i>	Osmanthus	<i>Osmanthus heterophyllus</i>
Ilima	<i>Sida fallax</i>	Pachysandra	<i>Pachysandra terminalis</i>
Ivy, Algerian	<i>Hedera canariensis</i>	Palm, Bamboo	<i>Rhapi excelsa</i>
Ivy, English	<i>Hedera helix</i>	Palm, Fishtail	<i>Caryota spp</i>
Jasmine, Orange	<i>Murraya paniculata</i>	Palm, Licuala	<i>Licuala Thunb. spp</i>
Jasmine, Primrose	<i>Jasminum mesnyi</i>	Palm, Macarthur	<i>Ptychosperma marcarthurii</i>
Jasmine, Star	<i>Trachelospermum jasminoides</i>	Palm, Manac	<i>Euterpe spp</i>
Jasmine, Showy	<i>Jasminum floridum</i>	Palm, Manila	<i>Veitchia merillii</i>
Juniper	<i>Juniperus spp</i>	Palm, Parlor	<i>Collinia elegans</i>
Justica	<i>Justica brandegeana</i>	Palm, Rattan	<i>Daemonorops spp</i>
Lantana, Trailing	<i>Lantana montevidensis</i>	Palm, Sealing Wax	<i>Cyrtostachyus lakka</i>
Lapalapa	<i>Cheirodendron spp</i>	Palm, Sugar	<i>Arenga saccharifera</i>
Larch, European	<i>Larix decidua</i>	Paperbark Tree	<i>Melaleuca leucadendra</i>
Laurel, English	<i>Prunus laurocerasus</i>	Periwinkle, Common	<i>Vinca minor</i>
Leucothoe, Coast	<i>Leucothoe axillaris</i>	Periwinkle, Large	<i>Vinca major</i>
Leucothoe, Drooping	<i>Leucothoe fontanesia (catesbaei)</i>	Photinia, Frasier's	<i>Photinia fraseri</i>
Lily of the Nile	<i>Agapanthus umbellatus</i>	Poinciana	<i>Delonix regia</i>
Lilac	<i>Syringa spp</i>	Pine	<i>Pinus spp</i>
Loquat*	<i>Eriobotrya japonica</i>	Pittosporum, Japanese	<i>Pittosporum tobira</i>
Magnolia	<i>Magnolia spp</i>	Plum, Natal	<i>Carissa macrocarpa</i>
Mahonia, Grape Holly	<i>Mahonia aquifolium</i>	Podocarpus, Yew	<i>Podocarpus macrophylla</i>
Maple	<i>Acer spp</i>	Pomegranite*	<i>Punica granatum</i>
Metallic Plant	<i>Hemigraphis colorata</i>	Poplar	<i>Populus spp</i>
Mexican Orange	<i>Choisya ternata</i>	Potentilla	<i>Potentilla fruticosa</i>
Money Tree	<i>Pleomele marginata</i>	Potentilla	<i>Potentilla fragiformis</i>
Murraea	<i>Murraea exotica</i>	Privet	<i>Ligustrum spp</i>
Murraea	<i>Murraea paniculata</i>	Protea species	<i>Banksia menziesii</i> <i>Leucadendron discolor</i> <i>Leucospermum cordifolium</i> <i>Protea eximia</i>
Myoporum	<i>Myoporum laetum</i>	Pyracantha	<i>Pyracantha coccinea</i>
Myrtle	<i>Myrtus communis</i>	Raphiolepis	<i>Raphiolepis umbellata</i>
Nandina	<i>Nandina domestica</i>	Redwood, Dawn	<i>Metasequoia glyptostroboides</i>
		Rhododendron	<i>Rhododendron spp</i>
		Rose	<i>Rosa spp</i>

ESTABLISHED ORNAMENTALS AND GROUND COVERS (con't)

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Russian Olive	<i>Eleagnus angustifolia</i>	Syzygium	<i>Syzygium paniculatum</i>
Scotch Broom	<i>Cytisus scoparius</i>	Trailing African Daisy	<i>Osteospermum fruticosum</i>
Sedum	<i>Sedum album</i>	Viburnum	<i>Viburnum spp</i>
Serviceberry, Saskatoon*	<i>Amelanchier alnifolia</i>	Wedelia	<i>Wedelia trilobata</i>
Spruce	<i>Picea spp</i>	Witchhazel, Common	<i>Hamamelis virginiana</i>
St. Johnswort, Golden	<i>Hypericum frondosum</i>	Woodwaxen, Common	<i>Genista tinctoria</i>
St. Johnswort, Gokdencup	<i>Hypericum patulum</i>	Xylosma	<i>Xylosma congestum</i>
Strawberry Tree	<i>Arbutus unedo</i>	Yew	<i>Taxus spp</i>
Sumac, Staghorn	<i>Rhus typhina</i>		

NEWLY TRANSPLANTED ORNAMENTALS AND GROUND COVERS

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Brassia, Dwarf	<i>Brassia actinophylla</i>	Poplar	<i>Populus spp</i>
Camelia	<i>Camelia japonica</i>	Privet	<i>Ligustrum spp</i>
Camation	<i>Dianthus caryophyllus</i>	Protea Species	<i>Banksia menziesii</i> <i>Leucadendron discolor</i> <i>Leucospermum cordifolium</i> <i>Protea eximia</i>
Cedar, Red	<i>Juniperus atlantica</i>	Rhododendron	<i>Rhododendron spp</i>
Coromandel	<i>Asystasia gargentica</i>	Russian Olive	<i>Eleagnus angustifolia</i>
Dogwood, Flowering	<i>Cornus florida</i>	Spruce	<i>Picea spp</i>
Elm	<i>Ulmus spp</i>	Trailing African Daisy	<i>Osteospermum fruticosum</i>
Forsythia	<i>Forsythia intermedia</i>	Wedelia	<i>Wedelia trilobata</i>
Gazania, Trailing	<i>Gazania uniflora leucotana</i>	Yew	<i>Taxus spp</i>
Hackberry	<i>Celtis occidentalis</i>		
Ivy, Algerian	<i>Hedera canariensis</i>		
Ivy, English	<i>Hedera helix</i>		
Hemlock, Canada	<i>Tsuga canadensis</i>		
Holly	<i>Ilex spp</i>		
Ice Plant	<i>Carpobrotus edule</i>		
Juniper	<i>Juniperus spp</i>		
Metallic Plant	<i>Hemigraphis colorata</i>		
Oak	<i>Quercus spp</i>		
Periwinkle, Common	<i>Vinca minor</i>		
Periwinkle, Large	<i>Vinca major</i>		
Pine	<i>Pinus spp</i>		

BEST AVAILABLE COPY

**FOREST NURSERY APPLICATION**

Seedlings of the following species can be treated with CHIPCO® RONSTAR® G Herbicide when at least 4 weeks old:

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Pine, Loblolly	<i>Pinus taeda</i>
Pine, Eastern White	<i>Pinus strobus</i>
Pine, Slash	<i>Pinus caribaea</i>

**SENSITIVE VARIETIES**

Under certain growing conditions, the plant varieties listed below may be sensitive to CHIPCO® RONSTAR® G Herbicide. Growers should evaluate usage on a small scale prior to large scale applications.

<u>Plant Species</u>	<u>Varieties</u>
Andromeda ( <i>Pieris japonica</i> )	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, ( <i>Rhododendron spp</i> ), Leuchtenfuer, Memoire John Hearrens, Hino-Crimson, White Grandeur, Everest, Girard Roberta
Cotoneaster	Eichholz, Himalayan, Shangri-la ( <i>Cotoneaster spp</i> )
Crapemyrtle ( <i>Lagerstroemia indica</i> )	Petite Red Imp, Seminole Bush
Heath, Spring ( <i>Erica carnea</i> )	Mediterranean Pink
Hemlock, Canada ( <i>Tsuga canadensis</i> )	Pendula
Hibiscus ( <i>Hibiscus syriacus</i> )	Red Heart
Rhododendron	Daphnoides, Nova Zembla, ( <i>Rhododendron spp</i> ), Catawba Album
Spruce, Norway ( <i>Picea abies</i> )	Pumila
Spruce, White ( <i>Picea glauca</i> )	Densata
Viburnum ( <i>Viburnum spp</i> )	Davidii

\* For use only on plants that will not bear edible fruit within one year of CHIPCO® RONSTAR® G Herbicide application.

**BEST AVAILABLE COPY**

**STORAGE AND DISPOSAL**

**STORAGE**

Do not contaminate water, food or feed by storage or disposal.  
Do not store near other pesticides or seeds.

**PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL**

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER

WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**NOTICE TO BUYER**

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

**THIS SPECIMEN LABEL IS INTENDED FOR USE ONLY AS A GUIDE IN PROVIDING GENERAL INFORMATION REGARDING THE DIRECTIONS, WARNINGS AND CAUTIONS ASSOCIATED WITH THE USE OF THIS PRODUCT. AS WITH ANY AGRICULTURAL CHEMICAL, ALWAYS FOLLOW THE LABEL INSTRUCTIONS ON THE PACKAGE BEFORE USING.**

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

CHIPCO AND RONSTAR are registered trademarks of Rhône-Poulenc.  
©1988 Rhône-Poulenc Ag Company

CHIPCO RONSTAR G (PENDING) submitted 9/5/89.

Form No. SP89606  
Printed USA -

Made in U.S.A.

