

2 of 6

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to lakes, streams or ponds. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation systems.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Store at temperatures above 20°F.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SUGARCANE

Asulox can be applied to either plant cane or cane grown from stubble. Apply Asulox as a water-mix spray for ground applications. Use 15 to 100 gallons of water per acre, depending on local practice. For aerial application, Asulox should be mixed in 3 to 5 gallons of water per acre, except in Hawaii, where 5 to 10 gallons of water per acre should be used.

Addition of an adjuvant cleared for use on growing crops to the Asulox water-mix spray will improve weed control when environmental conditions are not optimal. Use either a non-ionic surfactant containing a minimum of 80% active ingredient at the rate of 1 to 2 quarts per 100 gallons (0.25 to 0.5% V/V) of water-mix spray or a crop oil concentrate containing 80 to 85% paraffin based petroleum oil and 15 to 20% non-ionic surfactant at the rate of 4 quarts per 100 gallons (1% V/V) of water-mix spray.

The rates of Asulox given below are for broadcast applications. For banded application reduce the rate proportionally to the width of the band according to the following formula:


$$\frac{\text{Band width (Inches)}}{\text{Row width (Inches)}} \times \text{Broadcast rate} = \text{Band rate/acre.}$$

For spot treatments use 8 to 12 pints of Asulox in 100 gallons of water and apply in an amount not to exceed 50 gallons of total spray solution per acre. (Do not exceed 8 pints per acre per treatment.)

Asulox should be used when the weeds are actively growing.

Do not plant crops other than sugarcane in treated area. Cover crops may be planted if plowed under and not grazed.

Do not graze or feed sugarcane fodder and forage to livestock.



OCT 26 1987

356

SUGARCANE

For Single Application per Growing Season:

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|---|--|------------------------------|
| Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>) | Apply when the grass is 12 inches tall or less (addition of surfactant is necessary). | 8 Pints/Acre |
| Johnsongrass (<i>Sorghum halepense</i>) | Apply when the grass is between 18 and 24 inches tall. | |
| Paragrass or Californiagrass (<i>Brachiaria mutica</i> = <i>Panicum purpurascens</i>) | Apply when the grass is 6 to 8 inches tall or less. | |
| Crabgrass (<i>Digitaria spp.</i>) | If treatment is made before the grass reaches seed head formation then the lower rate should be used. If the grass is in early seed head formation then the higher rate should be used. | 6 to 8 Pints/Acre |
| Alexandergrass (<i>Brachiaria plantaginea</i>) Foxtail (<i>Setaria spp.</i>) Goosegrass (<i>Eleusine indica</i>) Broadleaf Panicum (<i>Panicum adspersum</i>) Barnyardgrass (<i>Echinochloa crus-galli</i>) | If treatment is made when the grass is 6 to 8 inches tall or less then the lower rate should be used. If the grass is greater than 8 inches tall then the higher rate should be used. | |
| | If treatment is made when the grass is 6 to 8 inches tall or less then the lower rate should be used. If the grass is greater than 8 inches tall then the higher rate should be used. | |
| | If treatment is made when the grass is 6 to 8 inches tall or less then the lower rate should be used. If the grass is greater than 8 inches tall then the higher rate should be used. | |




Two Applications per Growing Season:

This may be required when initial weed infestations are heavy or reinfestation from seed occurs. Two applications may also be used when treating weed species which germinate at different times during one growing season.

| WEED SPECIES | SPECIAL INSTRUCTIONS | FIRST APPLICATION RATE | SECOND APPLICATION RATE |
|--|--|------------------------------|------------------------------|
| Crabgrass (<i>Digitaria spp.</i>) | At each application the grass should be treated before seed head formation. | 6 to 8 Pints/Acre | 6 to 8 Pints/Acre |
| Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>) | At each application the grass should be 12 inches tall or less. (Addition of surfactant is necessary.) | 8 Pints/Acre | 8 Pints/Acre |
| Johnsongrass (<i>Sorghum halepense</i>) | At each application the grass should be between 18 and 24 inches tall. | 8 Pints/Acre | 4 to 8 Pints/Acre |

Do NOT treat sugarcane within 90 days of harvest.



REFORESTATION AREAS (Site Preparation Areas and Conifer Release Areas) AND CHRISTMAS TREE PLANTINGS

Asulox may be used as a postemergent treatment in Reforestation Areas and Christmas Tree Plantings where Douglas Fir, Grand Fir, Nobel Fir or Scotch Pine are grown. Do not graze or feed foliage from treated areas to livestock.

Asulox should be applied as a water-mix spray. For ground application use a minimum of 20 gallons of solution per acre, and for aerial application use a minimum of 10 gallons of solution per acre. Do not use a wetting agent with Asulox.

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|---|-------------|
| Western Bracken (<i>Pteridium aquilinum</i> var. <i>pubescens</i>) | Apply after bud break and hardening or firming of new tree growth. Bracken should be in full frond prior to treatment | 1 Gal./Acre |

NONCROPLAND

Asulox may be used as a postemergent treatment to control weeds on noncropland areas such as:

- | | |
|------------------------------------|--|
| Boundary fences | Pipeline rights-of-way |
| Ditchbanks | Railroad rights-of-way and yards |
| Fence rows | Storage areas and industrial plant sites |
| Highway and roadside rights-of-way | Utility rights-of-way and yards |
| Lumberyards | Warehouse lots |

A surfactant may be added to the spray solution at 0.25% by volume. (Use an approved non-ionic surfactant.)

Apply Asulox as a water-mix spray for ground applications using 20 to 100 gallons of solution per acre, depending on local practice to control the following weed species:

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|--|----------------------|
| Crabgrass (<i>Digitaria spp.</i>) | Apply before the grass reaches seed head formation. | 1 to 2 Gal./Acre |
| Johnsongrass (<i>Sorghum halepense</i>) | Apply when the grass is 18 inches or taller. Use the higher rate in well established heavy infestations. For spot treatment in Hawaii, use the higher rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre. | |
| Paragrass or Californiagrass (<i>Brachiaria mutica</i> = <i>Panicum purpurascens</i>) | Apply before the grass reaches seed head formation. For spot treatment in Hawaii, use the same rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre. | |
| Western Bracken (<i>Pteridium aquilinum</i> var. <i>pubescens</i>) | Apply when the fern is in full frond. | 7 to 8 Pints/Acre |

TURF

Asulox can be applied on St. Augustinegrass and Tifway 419 Bermudagrass turf. Apply one application per season postemergence to the weeds listed below. Use 20 to 50 gallons of water per acre in the spray solution.

| TURF SPECIES | WEED SPECIES | RATE |
|---|---|--------------|
| St. Augustinegrass Tifway 419 Bermudagrass | Bullgrass (<i>Paspalum supinum</i>) Crabgrass (<i>Digitaria sp.</i>) Goosegrass (<i>Eleusine indica</i>) Sandbur (<i>Cenchrus sp.</i>) | 5 Pints/Acre |

Do not use a surfactant. Do not apply to turf which is under stress or freshly mowed.

ORNAMENTALS


Asulox can be applied as a single, postemergent, broadcast application on the following ornamentals:

- | | |
|-------------------------------|--------------------------------|
| JUNIPERS | YEWS |
| <i>Juniperus andorra</i> | <i>Taxus cuspidata</i> |
| <i>Juniperus chinensis</i> | <i>Taxus media</i> |
| <i>Juniperus conferta</i> | <i>Podocarpus macrophyllus</i> |
| <i>Juniperus horizontalis</i> | |
| <i>Juniperus litoralis</i> | |
| <i>Juniperus sabina</i> | |

Treatment should be made with a minimum of 20 gallons of water per acre. Do not use a surfactant.

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|--|------------|
| Barnyardgrass (<i>Echinochloa crus-galli</i>) Crabgrass (<i>Digitaria sp.</i>) Fall Panicum (<i>Panicum dichotomiflorum</i>) Foxtails (<i>Setaria sp.</i>) Goosegrass (<i>Eleusine indica</i>) Horseweed (Marestail) (<i>Conyza canadensis</i>) | Apply when the weeds are between the stages of early seedling and early seed head formation. | 1 Gal/Acre |

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.



LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. **THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States



PM23

264-447

178

Ms. Karen S. Shearer
Rhone-Poulenc Ag Company
2 T.W. Alexander Drive
P.O. Box 12014
Research Triangle Park, NC 27709

Dear Ms. Shearer:

Subject: Asulam Registration Standard - Revised Labeling
Asulox Herbicide
EPA Registration No. 264-447
Your Submission Dated July 12, 1988

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

The amended labeling supersedes all previously accepted ones.

Sincerely yours,

Richard F. Mountfort
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosure

51094:I:Mountfort:LMF-1:KENCO:10/24/88:12/3/88:rw:VO:aw

CONCURRENCES

| SYMBOL | | | | | | | |
|---------|--|--|--|--|--|--|--|
| SURNAME | | | | | | | |
| DATE | | | | | | | |

2 of 8

ASULOX® HERBICIDE

For Postemergent Weed Control in Sugarcane, Turf, Ornamentals, Christmas Tree Plantings and Noncropland.

Faint regulatory text and a signature.

ACTIVE INGREDIENT:

Sodium salt of asulam(methyl sulfanylcarbamate) *36.2%

INERT INGREDIENTS:.....63.8%

* Equivalent of 33.1% asulam or not less than 3.34 pounds per gallon.

EPA Reg. No. 264-447

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If on Skin: Wash with plenty of soap and water. If irritation persists get medical attention.


If in Eyes: Flush with water for 15 minutes and get medical attention if irritation persists.

See Side Panel for Additional Precautionary Statements.

For **PRODUCT USE** Information Call 1-800-334-9745.
For **EMERGENCY** Information **ONLY** Call 24 Hours A Day 1-800-334-9745.

Rhône-Poulenc Ag Company
P.O. Box 12014, T.W. Alexander Drive
Research Triangle Park, N.C. 27709

CONTENTS:



570

PRECAUTIONARY STATEMENTS
CAUTION
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, lagoons, marshes or potholes). Do not clean equipment or dispose of equipment wash water in a manner that will contaminate water resources.

Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL STATEMENT
STORAGE

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Store at temperatures above 20°F.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse or equivalent. Puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

77

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation systems.

SUGARCANE

ASULOX® can be applied to either plant cane or cane grown from stubble. Apply ASULOX® as a water mix spray for ground applications. Use 15 to 100 gallons of water per acre, depending on local practice. For aerial application, ASULOX® should be mixed in 3 to 5 gallons of water per acre, except in Hawaii, where 5 to 10 gallons of water per acre should be used.

Addition of an adjuvant cleared for use on growing crops to the ASULOX® water mix spray will improve weed control when environmental conditions are not optimal. Use either a non-ionic surfactant containing a minimum of 80% active ingredient at the rate of 1 to 2 quarts per 100 gallons (0.25 to 0.5% V/V) of water mix spray or a crop oil concentrate containing 80 to 85% paraffin based petroleum oil and 15 to 20% non-ionic surfactant at the rate of 4 quarts per 100 gallons (1% V/V) of water mix spray.

The rates of ASULOX® given below are for broadcast applications. For banded application reduce the rate proportionally to the width of the band according to the following formula:

Band width(Inches)/Row width(Inches) x Broadcast rate = Band rate/acre

For spot treatments use 6 to 12 pints of ASULOX® in 100 gallons of water and apply in an amount not to exceed 50 gallons of total spray solution per acre. Do not exceed 8 pints per acre per treatment.

ASULOX® should be used when the weeds are actively growing.

Do not plant crops other than sugarcane in treated area. Cover crops may be planted if plowed under and not grazed.

Do not treat sugarcane within 90 days of harvest.

Do not graze or feed sugarcane fodder and forage to livestock.

Do not rotate with any crop which is not registered for use with Asulam for one year following the last application of this chemical.

5 of 6

For Single Application Per Growing Season:

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|---|-------------------|
| Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>) | Apply when the grass is 12 inches tall or less (addition of surfactant is necessary). | 8 pints/acre |
| Johnsongrass (<i>Sorghum halepense</i>) | Apply when the grass is between 18 to 24 inches tall. | |
| Paragrass or Californiagrass (<i>Bracharia mutica</i> or <i>Panicum purpurascens</i>) | Apply when the grass is 6 to 8 inches tall or less | |
| Crabgrass (<i>Digitaria spp.</i>) | If treatment is made before the grass reaches seed head formation then the lower rate should be used. If the grass is in early seed head formation then the higher rate should be used. | 6 to 8 pints/acre |
| Alexandergrass(<i>Bracharia plantaginea</i>) Foxtail(<i>Setaria spp.</i>) Goosegrass(<i>Eleusine indica</i>) Broadleaf Panicum(<i>Panicum adspersum</i>) Barnyardgrass(<i>Echinochloa crusgalli</i>) | If treatment is made when the grass is 6 to 8 inches tall or less then the lower rate should be used. If the grass is greater than 8 inches tall then the higher rate should be used. If treatment is made when the grass is 6 to 8 inches tall or less then the lower rate should be used. If the grass is greater than 8 inches tall then the higher rate should be used. | |

Two Applications Per Growing Season:

This may be required when initial weed infestations are heavy or reinfestation from seed occurs. Two applications may also be used when treating weed species which germinate at different times during one growing season.

| WEED SPECIES | SPECIAL INSTRUCTIONS | FIRST Application RATE | SECOND Application RATE |
|--|--|------------------------|-------------------------|
| Crabgrass(<i>Digitaria spp.</i>) | At each application the grass should be treated before seed head formation. | 6 TO 8 pints/acre | 6 TO 8 pints/acre |
| Itchgrass or Raoulgrass (<i>Rottboellia exaltata</i>) | At each application the grass should be 12 inches tall or less. Addition of surfactant is necessary. | 8 pints/acre | 8 pints/acre |
| Johnsongrass (<i>Sorghum halepense</i>) | At each application the grass should be between 18 and 24 inches tall. | 8 pints/acre | 4 to 8 pints/acre |

Do not treat sugarcane within 90 days of harvest. Do not graze or feed sugarcane forage and fodder to livestock.

678

CHRISTMAS TREE PLANTINGS

ASULOX® may be used as a postemergent treatment in Christmas Tree Plantings where Douglas Fir, Grand Fir, Nobel Fir or Scotch Pine are grown. Do not graze or feed foliage from treated areas to livestock.

ASULOX® should be applied as a water mix spray. For ground application use a minimum of 20 gallons of solution per acre, and for aerial application use a minimum of 10 gallons of solution per acre. Do not use a wetting agent with ASULOX®.

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|--|-------------|
| Western Bracken (<i>Pteridium aquilinum</i> var. <i>pubescens</i>) | Apply after bud break and hardening or firming of new tree growth. Bracken should be in full frond prior to treatment. | 1 Gal./Acre |

NONCROPLAND

ASULOX® may be used as a postemergent treatment to control weeds on noncropland areas such as:

| | |
|------------------------------------|--|
| Boundary fences | Pipeline rights-of -ways |
| Ditchbanks | Railroad rights-of -way and yards |
| Fence rows | Storage areas and industrial plant sites |
| Highway and roadside rights-of-way | Utility rights-of-way and yards |
| Lumberyards | Warehouse lots |

A surfactant may be added to the spray solution at 0.25% by volume. Use an approved non-ionic surfactant.

Apply ASULOX® as a water mix spray for ground applications using 20 to 100 gallons of solution per acre, depending on local practice to control the following weed species:

1780

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|--|--|-------------------|
| Crabgrass(<i>Digitaria spp.</i>) | Apply before the grass reaches seed head formation. | 1 to 2 Gal./Acre |
| Johnsongrass (<i>Sorghum halepense</i>) | Apply when the grass is 18 inches or taller. Use the higher rate in well established heavy infestations. For spot treatment in Hawaii, use the higher rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre. | |
| Paragrass or Californiagrass (<i>Brachiaria mutica</i> or <i>Panicum purpurascens</i>) | Apply before the grass reaches seed head formation. For spot treatment in Hawaii, use the same rate in 100 gallons of solution and apply an amount not to exceed 50 gallons of total solution per acre. | |
| Western Bracken (<i>Pteridium aquilinum</i> var. <i>pubescens</i>) | Apply when the fern is in full frond. | 7 to 8 pints/acre |

TURF

ASULOX® can be applied on St. Augustinegrass and Tifway 419 Bermudagrass turf. Apply one application per season postemergence to the weeds listed below. Use 20 to 50 gallons of water per acre in the spray solution.

| TURF SPECIES | WEED SPECIES | RATE |
|-------------------------|--|--------------|
| St. Augustinegrass | Bullgrass(<i>Paspalum supinum</i>) Crabgrass(<i>Digitaria spp.</i>) Goosegrass(<i>Eleusine indica</i>) | 5 Pints/acre |
| Tifway 419 Bermudagrass | Sanbur(<i>Cenchrus spp.</i>) | |

Do not use a surfactant. Do not apply to turf which is under stress of freshly mowed.

ORNAMENTALS

ASULOX® can be applied as a single, postemergent, broadcast application on the following ornamentals:

JUNIPERS

Juniperus andorra
Juniperus chinensis
Juniperus conferta
Juniperus horizontalis
Juniperus Litoralis
Juniperus sabina

YEWS

Taxus suspidata
Taxus media
Podocarpus macrophyllus

8 of 8

Treatment should be made with a minimum of 20 gallons of water per acre. Do not use a surfactant.

| WEED SPECIES | SPECIAL INSTRUCTIONS | RATE |
|---|--|--------------|
| Barnyardgrass(<i>Echinochloa crusgalli</i>) Crabgrass(<i>Digitaria spp.</i>) Fall Panicum(<i>Panicum dichotomiflorum</i>) Foxtails(<i>Setaria spp.</i>) Goosegrass(<i>Eleusine indica</i>) Horseweed(Marestail) (<i>Conyza canadensis</i>) | Apply when the weeds are between the stages of early seedling and early seed head formation. | 1 Gal. /Acre |

Lower conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.