


Product Number
3357

rovral
FUNGICIDE



HOW TO USE ROVRAL ON STRAWBERRIES

PEST	APPLICATION	RATE	WHEN AND HOW TO USE
Botrytis crown rot (<i>Botrytis</i> sp.)	Dip	2.0 lbs/100 gallons	Apply Rovral as a 5 minute pre-plant dip of transplants immediately prior to planting.
Gray mold (<i>Botrytis cinerea</i>)	Foliar	1.5 - 2.0 lbs/Acre	<p>A maximum of 6 foliar sprays may be applied at 10 - 14 day intervals. The first foliar application should occur at no later than 10% bloom. Thorough coverage of the fruit is essential so do not apply Rovral in less than 100 gallons per acre.</p> <p>Rovral may be applied up to and including the day of harvest.</p> <p>The higher rate and/or shorter spray interval should be used under severe disease conditions.</p> <p>Rovral should not be used in an alternating spray program with Ronilan.</p>

BEST AVAILABLE COPY

This labeling must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

0287

RHÔNE-POULENC INC.

AGROCHEMICAL DIVISION

Moorestown, New Jersey, N.J. 08052