

MOCAP® Plus 4-2 EC Nematicide-Insecticide submitted to EPA 2/13/89

FRONT PANEL

RESTRICTED USE PESTICIDE

Acute human oral, dermal and inhalation toxicity. For retail sale to, and use by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

1 3/8"

MOCAP® Plus 4-2 EC Nematicide-Insecticide

ACTIVE INGREDIENTS:

Ethoprop: (O-Ethyl S, S-Dipropyl Phosphorodithioate) .....	16.0%
Disulfoton: (O, O-Diethyl S-[2-(ethylthio) ethyl] Phosphorodithioate) .....	23.0%
INERT INGREDIENTS: .....	31.0%

\*Equivalent to 4 pounds per gallon.

\*\*Equivalent to 2 pounds per gallon.

EPA Reg. No. 264-464

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN  
DANGER POISON  
PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Give 1 or 2 glasses of water and induce vomiting by touching back of throat with the finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician, Poison Control Center, or hospital emergency room.

If not breathing, give mouth to mouth artificial respiration.

IN CASE OF SKIN CONTACT: Immediately remove all contaminated clothing, and wash skin with soap and water. Wash clothing before reuse. Discard contaminated shoes.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Call a physician immediately in all cases of suspected poisoning

Persons providing first aid should prevent contamination to themselves.

FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Inside Label

For PRODUCT USE Information Call 1-800-334-9745

For EMERGENCY Information ONLY Call 24 Hours A Day 1-800-334-7577

Rhone-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

MOCAP is a registered trademark of Rhone-Poulenc

ACCEPTED  
with COMMENTS  
to EPA Letter Dated.

AUG 01 1989

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended for the purposes  
registered under EPA Reg. No.

264-464

ARTWORK is PMS 485 + black

**PRECAUTIONARY STATEMENTS** - 10 pt for all headings  
**DANGER**  
**HAZARDS TO HUMAN AND DOMESTIC ANIMALS**

8pt for all text

Fatal if swallowed, inhaled, or absorbed through skin. Rapidly absorbed through skin. Wear water proof protective clothing, rubber gloves and goggles. In case of contact, wash immediately with water.

Do not breathe fumes or spray mist. Wear an AO R-6058 respirator with an R-58 cartridge, or equivalent, for protection during field handling and field exposure. After use, wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all clothing with soap and hot water before reuse. Discard contaminated shoes. Do not contaminate feed or food.

Do not store in or around the home.

**SYMPTOMS OF POISONING:** Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

**NOTE TO PHYSICIAN**

This product is a cholinesterase inhibitor. Severe symptoms and signs include diarrhea, pinpoint and non-reactive pupils, respiratory difficulty, pulmonary edema, cyanosis, loss of sphincter control, convulsions, and coma. Solvent presents aspiration hazard. Gastric lavage may be indicated if material was taken internally. Support respiration as needed. Measures should include removal of secretions, maintenance of patent airway, and if necessary, artificial ventilation.

If cyanosis is absent, give ATROPINE 2-4 mg intravenously (0.5 mg/kg for children). Repeat Atropine at 5 to 10 minute intervals until atropinization occurs (dry, flushed skin, tachycardia, pupillary dilation), and maintain for 48 hours. If cyanotic, give initial atropine intramuscularly and start measures to improve ventilation.

Start 2-PAM at the same time. Give 1 to 2 gm 2-PAM (20 to 40 mg/kg for children) in 100cc saline over 15 to 30 minutes. If pulmonary edema is present, give intravenously slowly as a 5% solution in water over 5 minutes.

A second dose may be given after 1 hour if muscle weakness persists. Additional doses may be given cautiously for persistent muscle weakness. May be given by intramuscular or subcutaneous routes if intravenous administration is not feasible.

In case of skin contact, wash patient with soap and water followed by wash with 95% ethyl alcohol. Keep patient under constant observation for 24 to 36 hours. Symptoms may persist for 1 month.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift and run off from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

Do not apply within 140 feet of inland freshwater habitats.

Along the Atlantic seaboard, do not apply within 800 feet of brackish water habitats.

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**BEST AVAILABLE COPY**

**BEE CAUTION**

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning. Contact your Cooperative Agricultural Extension Service or your local Rhone-Poulenc Ag Company representative for further information.

**DIRECTIONS FOR USE**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

Do not apply in Long Island, New York.

Do not apply MOCAP® Plus 4-2 EC Nematicide-Insecticide through any type of irrigation system.

**REENTRY STATEMENT**

Reentry into treated fields is prohibited within 24 hours of application. State or local use and reentry regulations may be more restrictive than those stated on this label. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not allow reentry into treated fields within 24 hours of application, unless appropriate protective clothing is worn. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers and given in a language customarily understood by the workers.

Oral and written warnings must include the STATEMENT OF PRACTICAL TREATMENT and notification that unprotected persons may enter treated fields or area only after chemical has been mixed in the soil.

**EQUIPMENT:** MOCAP spray mixtures may affect some plastic materials after prolonged use. Spray tubing, hoses and gaskets should be inspected frequently, particularly black-colored materials. Replace any material which is softened or swollen. Polyethylene, polypropylene, nylon, and teflon materials are recommended. Polyvinyl chloride (PVC) material is not recommended.

**MIXING:** Add the recommended amount to the water in the spray tank and mix well. Use enough water in the sprayer (usually 20 to 25 gallons) to thoroughly and evenly cover the area to be treated. Agitate frequently during use.

**INCORPORATION INFORMATION**

MOCAP nematicide-insecticide can be incorporated with a rotary tiller, rotary hoe, spring tooth harrow, by double-discing or with other equipment for mixing the chemical with the soil to the depth recommended. Follow any specific incorporating directions given under the crop heading.


If moisture is applied immediately after application to carry the treatment into the soil, only shallow incorporation (1/4 to 1/2 inch) with mechanical equipment such as drag chains is required.

**TOBACCO NEMATODES CONTROLLED**

- Dagger nematode (Xiphinema spp.)
- Lesion nematode (Pratylenchus spp.)
- Root-knot nematode (Meloidogyne hapla, Meloidogyne incognita)
- Stunt nematode (Tylenchorhynchus spp.)

**TOBACCO INSECTS CONTROLLED**

- Wireworm (Conoderus spp.)
- Tobacco flea beetle (Epicrix hirtipennis)
- Tobacco aphid (Myzus nicotianae)



**TOBACCO**

**FOR CONTROL OF NEMATODES, FLEA BEETLES, APHIDS, AND WIREWORMS**

**Row Treatment:** Use 1 1/2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide per acre (42 inch row spacing) or 15.4 fluid ounces per 1000 linear feet of row. Apply after fertilizer has been listed or bedded. Make application in a band 18 inches wide on the row from 1 week before planting to at-planting time. Mix MOCAP® Plus 4-2 EC with the top 2 to 4 inches of soil right after application. Use a rotary tiller, rotary hoe, or other cultivating equipment that will do a good job of mixing MOCAP® Plus 4-2 EC with the soil.

**FOR CONTROL OF NEMATODES, FLEA BEETLES, APHIDS, AND WIREWORMS**

**Broadcast Treatment:** Apply 1 1/2 to 2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide per acre from 1 week before planting to at-planting time. Mix with the top 2 to 4 inches of soil right after application. Use double gang disc harrow, or other equipment that will do a good job of mixing MOCAP® Plus 4-2 EC with the soil. After mixing, apply fertilizer and, for best results, shape a wide bed with a tractor equipped with 4 disc hillers and wide sweeps for running out the middles. Be sure that only the treated soil (top 2 to 4 inches) is used to make beds. Use the higher rate in fields where nematode injury to tobacco has been especially severe, or where the soil is sandy.

**FOR CONTROL OF NEMATODES, APHIDS, FLEA BEETLES, WIREWORMS, BLACK SHANK AND BLUE MOLD.**

**Broadcast Treatment In Tank Mix With Ridomil® 2EC:** Use 1 1/2 to 2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide mixed with 2 to 4 quarts of Ridomil 2EC per acre. Mix the treatment with the top 2 to 4 inches of soil right after application then form beds. Read both product labels for specific use and rate information. Follow all precautions, directions, and restrictions on the labels.

Rhone-Poulenc Ag Company does not recommend the use of MOCAP® Plus 4-2 EC Nematicide-Insecticide on tobacco in Florida.

Do not plant any food crop in rotation after a field treatment with disulfoton unless it is a registered use for disulfoton.

**STORAGE AND DISPOSAL**

**BEST AVAILABLE COPY**

**STORAGE**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL**

Triple rinse or equivalent. Then dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; that the directions, warnings, and other statements on this label are based upon responsible evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

**RESTRICTED USE PESTICIDE**

Acute human oral, dermal and inhalation toxicity. For retail sale to, and use by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

**MOCAP® Plus 4-2 EC Nematicide-Insecticide**

**ACTIVE INGREDIENTS:**

Ethoprop: (O-Ethyl S, S-Dipropyl Phosphorodithioate) .....	46.0%
Disulfoton: (O, O-Diethyl S-[2-(ethylthio) ethyl] Phosphorodithioate) .....	23.0%
<b>INERT INGREDIENTS:</b> .....	31.0%

\*Equivalent to 4 pounds per gallon.  
\*\*Equivalent to 2 pounds per gallon.

EPA Reg. No. 264-464

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER POISON**  
**PELIGRO**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Give 1 or 2 glasses of water and induce vomiting by touching back of throat with the finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician, Poison Control Center, or hospital emergency room.

If not breathing, give mouth to mouth artificial respiration.

**IN CASE OF SKIN CONTACT:** Immediately remove all contaminated clothing, and wash skin with soap and water. Wash clothing before reuse. Discard contaminated shoes.

**IF IN EYES:** Flush eyes with plenty of water for at least 15 minutes. Call a physician immediately in all cases of suspected poisoning

Persons providing first aid should prevent contamination to themselves.

**FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Inside Label**

For PRODUCT USE Information Call 1-800-334-9745

For EMERGENCY Information ONLY Call 24 Hours A Day 1-800-334-7577

Rhone-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

MOCAP is a registered trademark of Rhone-Poulenc



**PRECAUTIONARY STATEMENTS**

**DANGER**

**HAZARDS TO HUMAN AND DOMESTIC ANIMALS**

Fatal if swallowed, inhaled, or absorbed through skin. Rapidly absorbed through skin. Wear water proof protective clothing, rubber gloves and goggles. In case of contact, wash immediately with water.

Do not breathe fumes or spray mist. Wear an AO R-6058 respirator with an R-58 cartridge, or equivalent, for protection during field handling and field exposure. After use, wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all clothing with soap and hot water before reuse. Discard contaminated shoes. Do not contaminate feed or food.

Do not store in or around the home.

**SYMPTOMS OF POISONING:** Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

**NOTE TO PHYSICIAN**

This product is a cholinesterase inhibitor. Severe symptoms and signs include diarrhea, pinpoint and non-reactive pupils, respiratory difficulty, pulmonary edema, cyanosis, loss of sphincter control, convulsions, and coma. Solvent presents aspiration hazard. Gastric lavage may be indicated if material was taken internally. Support respiration as needed. Measures should include removal of secretions, maintenance of patent airway, and if necessary, artificial ventilation.

If cyanosis is absent, give ATROPINE 2-4 mg intravenously (0.5 mg/kg for children). Repeat Atropine at 5 to 10 minute intervals until atropinization occurs (dry, flushed skin, tachycardia, pupillary dilation), and maintain for 48 hours. If cyanotic, give initial atropine intramuscularly and start measures to improve ventilation.

Start 2-PAM at the same time. Give 1 to 2 gm 2-PAM (20 to 40 mg/kg for children) in 100cc saline over 15 to 30 minutes. If pulmonary edema is present, give intravenously slowly as a 5% solution in water over 5 minutes.

A second dose may be given after 1 hour if muscle weakness persists. Additional doses may be given cautiously for persistent muscle weakness. May be given by intramuscular or subcutaneous routes if intravenous administration is not feasible.

In case of skin contact, wash patient with soap and water followed by wash with 95% ethyl alcohol. Keep patient under constant observation for 24 to 36 hours. Symptoms may persist for 1 month.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift and run off from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

Do not apply within 140 feet of inland freshwater habitats.

Along the Atlantic seaboard, do not apply with 800 feet of brackish water habitats.

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**BEE CAUTION**

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning. Contact your Cooperative Agricultural Extension Service or your local Rhone-Poulenc Ag Company representative for further information.

**DIRECTIONS FOR USE**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

**BEST AVAILABLE COPY**

Do not apply in Long Island, New York.

Do not apply MOCAP® Plus 4-2 EC Nematicide-Insecticide through any type of irrigation system.

**REENTRY STATEMENT**

Reentry into treated fields is prohibited within 24 hours of application. State or local use and reentry regulations may be more restrictive than those stated on this label. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not allow reentry into treated fields within 24 hours of application, unless appropriate protective clothing is worn. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers and given in a language customarily understood by the workers.

Oral and written warnings must include the STATEMENT OF PRACTICAL TREATMENT and notification that unprotected persons may enter treated fields only after chemical has been mixed in the soil.

**EQUIPMENT:** MOCAP spray mixtures may affect some plastic materials after prolonged use. Spray tubing, hoses and gaskets should be inspected frequently, particularly black-colored materials. Replace any material which is softened or swollen. Polyethylene, polypropylene, nylon, and teflon materials are recommended. Polyvinyl chloride (PVC) material is not recommended.

**MIXING:** Add the recommended amount to the water in the spray tank and mix well. Use enough water in the sprayer (usually 20 to 25 gallons) to thoroughly and evenly cover the area to be treated. Agitate frequently during use.

**INCORPORATION INFORMATION**

MOCAP nematicide-insecticide can be incorporated with a rotary tiller, rotary hoe, spring tooth harrow, by double-discing or with other equipment for mixing the chemical with the soil to the depth recommended. Follow any specific incorporating directions given under the crop heading.

If moisture is applied immediately after application to carry the treatment into the soil, only shallow incorporation (1/4 to 1/2 inch) with mechanical equipment such as drag chains is required.

**TOBACCO NEMATODES CONTROLLED**

Dagger nematode	(Xiphinema spp.)
Lesion nematode	(Pratylenchus spp.)
Root-knot nematode	(Meloidogyne hapla, Meloidogyne incognita)
Stunt nematode	(Tylenchorhynchus spp.)

**TOBACCO INSECTS CONTROLLED**

Wireworm	(Conoderus spp.)
Tobacco leaf beetle	(Epirix hirtipennis)
Tobacco aphid	(Myzus nicotianae)

**BEST AVAILABLE COPY**

## TOBACCO

### FOR CONTROL OF NEMATODES, FLEA BEETLES, APHIDS, AND WIREWORMS

**Row Treatment:** Use 1 1/2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide per acre (42 inch row spacing) or 15.4 fluid ounces per 1000 linear feet of row. Apply after fertilizer has been listed or bedded. Make application in a band 18 inches wide on the row from 1 week before planting to at-planting time. Mix MOCAP® Plus 4-2 EC with the top 2 to 4 inches of soil right after application. Use a rotary tiller, rotary hoe, or other cultivating equipment that will do a good job of mixing MOCAP® Plus 4-2 EC with the soil.

### FOR CONTROL OF NEMATODES, FLEA BEETLES, APHIDS, AND WIREWORMS

**Broadcast Treatment:** Apply 1 1/2 to 2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide per acre from 1 week before planting to at-planting time. Mix with the top 2 to 4 inches of soil right after application. Use double gang disc harrow, or other equipment that will do a good job of mixing MOCAP® Plus 4-2 EC with the soil. After mixing, apply fertilizer and, for best results, shape a wide bed with a tractor equipped with 4 disc hillers and wide sweeps for running out the middles. Be sure that only the treated soil (top 2 to 4 inches) is used to make beds. Use the higher rate in fields where nematode injury to tobacco has been especially severe, or where the soil is sandy.

### FOR CONTROL OF NEMATODES, APHIDS, FLEA BEETLES, WIREWORMS, BLACK SHANK AND BLUE MOLD.

**Broadcast Treatment In Tank Mix With Ridomil® 2EC:** Use 1 1/2 to 2 gallons of MOCAP® Plus 4-2 EC Nematicide-Insecticide mixed with 2 to 4 quarts of Ridomil 2EC per acre. Mix the treatment with the top 2 to 4 inches of soil right after application then form beds. Read both product labels for specific use and rate information. Follow all precautions, directions, and restrictions on the labels.

Rhone-Poulenc Ag Company does not recommend the use of MOCAP® Plus 4-2 EC Nematicide-Insecticide on tobacco in Florida.

Do not plant any food crop in rotation after a field treatment with disulfoton unless it is a registered use for disulfoton.

## STORAGE AND DISPOSAL

### STORAGE

Do not contaminate water, food, or feed by storage or disposal.

### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL

Triple rinse or equivalent. Then dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; that the directions, warnings, and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

**BEST AVAILABLE COPY**