

PM 75 2-1-77

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons directly under their supervision and only for those uses covered by the Certified Applicator's certification. See precautionary statements for reasons this product is classified Restricted Use.

BUCTRIL + atrazine HERBICIDE

POSTEMERGENT HERBICIDE FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN CORN AND SORGHUM.

ACTIVE INGREDIENT: Octanoic acid ester of bromoxynil* (3,5-dibromo-4-hydroxybenzotrile).....15.74%
atrazine** (2-chloro-4-ethylamino-6-isopropylamino-S-triazine).....21.62%

INERT INGREDIENTS:.....62.64%

*Product contains bromoxynil octanoate equivalent to 10.81% of bromoxynil or 1.0 pound of bromoxynil per gallon.

**Product contains 2.0 pounds of atrazine per gallon.

E.P.A. Reg. No. 264-477

E.P.A. Est. No. 264-MO-02

KEEP OUT OF REACH OF CHILDREN

CAUTION

SHAKE WELL BEFORE USING

ACCEPTED

FEB 21 1977

Division of Pesticide Control, Department of Agriculture
Washington, D.C. 20250
EPA Reg. No. 264-477

For PRODUCT USE Information Call 1-800-334-9745.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

PRECAUTIONARY STATEMENTS

CAUTION - RESTRICTED USE

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Use of this product may be hazardous to your health. Exposure to this product has been shown to cause birth defects in laboratory animals. Women of childbearing age should be particularly careful when handling this product to avoid ingestion and skin contact.

Harmful if swallowed or absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

Users of this product are encouraged to participate in Rhône-Poulenc's educational program dealing with potential hazards and safer use of this product. Further information on this program can be obtained by contacting Rhône-Poulenc Ag Company at 1-800-334-9745.

ALL PERSONS HANDLING THE CONCENTRATE MUST OBSERVE THE FOLLOWING REQUIREMENTS.

Wear clean cotton (or cloth) coveralls that cover all parts of the body except the head, hands, and feet. The coveralls must be worn over a long sleeve shirt and long pants. Wear clean nitrile gloves, a chemical resistant apron, goggles or face shield, and chemical resistant shoes, shoe coverings, or boots. If you use a mechanical transfer system for all mixing and loading operations, use of a chemical resistant apron is optional.

If you will handle a total of 30 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinse directly to the mixing or spray tank.

ALL PERSONS APPLYING THIS PRODUCT, OR REPAIRING OR CLEANING EQUIPMENT USED WITH THIS PRODUCT, MUST OBSERVE THE FOLLOWING REQUIREMENTS.

When applying this product from a tractor (unless the tractor has a completely enclosed cab), or when repairing or cleaning equipment used with the product, you must wear clean nitrile gloves and clean cotton (or cloth) coveralls that cover all parts of the body except the head, hands, and feet. Coveralls must be worn over a long sleeve shirt and long pants.

All applicators (except for pilots) and all persons repairing or cleaning equipment used with this product must wear chemical resistant shoes, shoe coverings, or boots.

Application from a tractor with a completely enclosed cab or aerial application is required whenever this product is applied to 180 or more acres in a day. To avoid contamination, coveralls and gloves worn when handling the concentrate must be removed prior to entering an enclosed cab or cockpit. When applying from a tractor with an enclosed cab, clean coveralls and clean nitrile gloves must be kept inside the cab, and must be worn when exiting the cab to perform in-field maintenance or repair.

To reduce exposure to residues, wash the spray rig, tractor, and all other equipment used to handle or apply this product with water daily or before using the equipment for any other purpose.

APPLICATION BY CHEMIGATION must be done by fixed pipe, overhead sprinkler systems or hand moved pipe. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

DURING AERIAL APPLICATION, human flaggers are prohibited unless in enclosed vehicles. Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, hospitals, shopping areas, etc.)

GENERAL REQUIREMENTS FOR ALL USE

IMPORTANT! Before removing gloves or starting a new work operation, rinse the outside of the gloves thoroughly with water. Always remove gloves and wash your hands and face with soap and water before smoking, eating, drinking, or toileting.

AFTER WORK. Gloves must be discarded at the end of each day. Take off all clothes and shoes. Shower with soap and water. Do not wear or re-use contaminated clothing. Wash protective equipment with water after each use. Protective clothing worn during use must be laundered separately from other household articles.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention. Do not induce vomiting, contains petroleum distillates. Do not give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with water for 15 minutes. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and fish. Use with care when applying to areas frequented by wildlife or adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or Oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers and given in a language customarily understood by the workers. Oral and Written warnings must include the STATEMENT OF PRACTICAL TREATMENT and REENTRY STATEMENTS as they appear on the label.

GENERAL INFORMATION

BUCTRIL® + atrazine contains the equivalent of 1 pound per gallon of octanoic acid ester of bromoxynil and 2 pounds per gallon of atrazine.

BUCTRIL® + atrazine is a selective postemergence herbicide for control of important broadleaf weeds infesting field corn, popcorn, and sorghum. Optimum weed control is obtained when **BUCTRIL® + atrazine** is applied to actively growing weed seedlings. **BUCTRIL® + atrazine** is primarily a contact herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control. **BUCTRIL® + atrazine** has limited residual activity. However, dependent on weather conditions following application, subsequent flushes of weeds may not be controlled by the initial treatment.

Occasional transitory leaf burn may occur. The temporary leaf burn is similar to that seen with liquid fertilizer. Because the activity of **BUCTRIL® + atrazine** is not systemic, recovery of the crop is generally rapid with no lasting effect.

MIXING, LOADING AND HANDLING INSTRUCTIONS

2.5 Gallon Containers

It is strongly recommended that special care be taken in mixing and loading this product. Hands should be placed on the container in such a way as to avoid possible drip or splash. Correct procedures for mixing and loading are provided in Rhône-Poulenc Ag Company's Educational Program.

30 Gallon and Bulk Containers

If you will handle a total of 30 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not

presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinseate directly to the mixing or spray tank.

BUCTRIL® + ATRAZINE ALONE: Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of **BUCTRIL® + atrazine**. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIXTURES: **BUCTRIL® + atrazine** can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply **BUCTRIL® + atrazine** in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. Add the **BUCTRIL® + atrazine** first and mix thoroughly. Add the other product to the tank while maintaining agitation and add water to the spray tank to the desired level. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

If tank mixing with products other than those listed within each crop section, a compatibility test is recommended to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture with **BUCTRIL® + atrazine**.

SPRAYABLE LIQUID FERTILIZERS AND SPRAY ADDITIVES

BUCTRIL® + atrazine can be applied in combination with sprayable liquid fertilizer or spray additives such as surfactants or crop oil concentrate. When tankmixing with liquid fertilizer always add the fertilizer to the spray tank first and agitate thoroughly before adding **BUCTRIL® + atrazine**. Always predetermine the compatibility with liquid fertilizer by mixing small proportional quantities in advance. Agitation must be maintained during filling and application operations to ensure that **BUCTRIL® + atrazine** is evenly mixed with the fertilizer.

CAUTION: Fertilizers and spray additives can increase foliage leaf burn when applied with **BUCTRIL® + atrazine**. Do not apply fertilizers or spray additives with **BUCTRIL® + atrazine** if leaf burn is a major concern due to environmental conditions, crop or variety sensitivity to **BUCTRIL® + atrazine**.

APPLICATION PROCEDURES

BUCTRIL® + atrazine can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment.

GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a minimum spray pressure of 30 psi at the nozzle tips are recommended. Other nozzle types that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop(3) nozzles are not recommended as weed control with **BUCTRIL® + atrazine** may be reduced. In general a minimum spray volume of 10 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment a maximum speed of 10 mph is suggested if field conditions cause excessive boom movement during application and subsequent poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas.

When weed infestations are heavy, use of higher spray volumes and spray pressure will be helpful in obtaining uniform weed coverage. When corn or grain sorghum are large enough to interfere with the spray pattern, drop nozzles should be used to obtain uniform weed coverage. Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended. A minimum spray volume of 3 GPA may be used if crop canopy and weed density allow adequate spray coverage at that gallonage.

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement. Off target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.

SPRINKLER IRRIGATION APPLICATION

BUCTRIL® + atrazine Herbicide can be applied through sprinkler irrigation systems to field corn, popcorn, and sorghum.

Apply **BUCTRIL® + atrazine** Herbicide through sprinkler systems including center pivot, lateral move, side (wheel) roll solid set or hand move irrigation systems only. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours

After chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle. Do not apply this product through any other type of irrigation system.

SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH AUTOMATED SPRINKLER IRRIGATION SYSTEM.

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Agitation is recommended in the pesticide supply tank when applying the **BUCTRIL® + atrazine Herbicide**.
9. **BUCTRIL® + atrazine Herbicide** should be applied continuously for the duration of the water application with center pivot and continuous lateral move systems. Application of **BUCTRIL® + atrazine Herbicide** should be made during the last 30-45 minutes of the irrigation set with other overhead sprinkler systems.
10. For best performance, set the sprinkler system to deliver approximately 0.5 inch or less of water per acre.
11. Remove scale, pesticide residues and other foreign matter from the supply tank and entire injector system. Flush with clean water.
12. If **BUCTRIL® + atrazine Herbicide** is diluted in the supply tank, fill the tank with half of the water amount desired, add the **BUCTRIL® + atrazine** and then add remaining water amount with agitation. Always dilute with at least 4 parts of water to 1 part of **BUCTRIL® + atrazine**.
13. Start the sprinklers and then inject **BUCTRIL® + atrazine Herbicide** into the irrigation line. **BUCTRIL® + atrazine** should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to insure adequate mixing. Refer to the **BUCTRIL® + atrazine Herbicide** label for detailed information on application rates and timings.

CHEMIGATION USER PRECAUTIONS

Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils.

Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.

Allow sufficient time for pesticide to be flushed through all the lines and nozzles before turning off irrigation water.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.


Do not connect an irrigation system used for pesticide application to a public water system.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CULTIVATION


When properly utilized, timely cultivations may aid overall weed control efforts as well as crop growth. However, cultivation BEFORE or DURING **BUCTRIL® + atrazine** applications may place target weeds under stress, resulting in erratic weed control. Whenever **BUCTRIL® + atrazine** is being utilized in an overall weed control program, plan to postpone any anticipated cultivations until 5-7 days after application to ensure best performance.



CORN (FIELD AND POP) AND SORGHUM (GRAIN AND FORAGE)

BUCTRIL® + ATRAZINE RECOMMENDATIONS

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUCTRIL® + atrazine	1 1/2 - 2 pints/A	<p>Apply to corn between the 3 leaf stage but before corn is 30 inches tall.</p> <p>Apply to sorghum between the 3 leaf stage and prior to preboot stage (growth stage 4)</p>	See APPLICATION RATE TABLE for list of weeds and corresponding maximum stage of growth that are controlled by BUCTRIL® + atrazine Herbicide at 1 1/2, 2 or 3 pints/A.
	3 pints/A	<p>Apply to corn between the 4 leaf stage but before corn is 30 inches tall.</p> <p>Apply to sorghum between the 4 leaf stage and prior to preboot to boot stage (growth stage 4)</p>	
	Chemigation 3 pints/A	<p>Apply to corn between the 3 leaf stage but before corn is 30 inches tall through automated sprinkler irrigation systems.</p> <p>Apply to sorghum between the 3 leaf stage and prior to preboot stage (growth stage 4) through automated sprinkler irrigation systems.</p>	Apply to broadleaf weeds up to the 4 leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.




APPLICATION RATE TABLE FOR CORN AND SORGHUM

WEED SPECIES	BUCTREL® + ATHAZINE RATE					
	1 1/2 Pints/A		2 Pints/A		3 Pints/A	
	MAXIMUM LEAF STAGE (INCHES)	MAXIMUM WEED HEIGHT (INCHES)	MAXIMUM LEAF STAGE (INCHES)	MAXIMUM WEED HEIGHT (INCHES)	MAXIMUM LEAF STAGE (INCHES)	MAXIMUM WEED HEIGHT (INCHES)
Black nightshade (Solanum nigrum)	4	4	6	6	6	6
Buffalobur (Solanum rostratum)	4	4	6	4	6	4
Burcucumber (Sicyos angulatus)	-	-	4	4	4	4
Common cocklebur (Xanthium pensylvanicum)	6	8	8	10	10	12
Common Lambquarters (Chenopodium album)	-	6	-	10	-	12
Common ragweed (Ambrosia artemisiifolia)	6	4	8	6	8	6
Eastern black nightshade (Solanum ptyzanthum)	4	4	6	6	6	6
Entleaf morningglory (Ipomoea hederacea)	-	-	4	3	4	3
Giant ragweed (Ambrosia trifida)	4	6	6	8	6	8
Hemp sesbania (Sesbania eschola)	4	4	4	4	4	4
Hyleaf morningglory (Ipomoea hederacea)	3	3	4	4	4	4
Jimsonweed (Datura stramonium)	4	4	6	6	6	6
Kochia (Kochia scoparia)	-	-	-	2	-	4
Ladysthumb (Polygonum persicaria)	4	4	6	6	8	8
Palmleaf morningglory (Ipomoea wrightii)	-	-	4	3	4	3
Pennsylvania smartweed (Polygonum pensylvanicum)	4	4	6	6	8	8
Prickly sida (Sida spinosa)	-	-	4	1	6	2
Puncturevine (Tribulus terrestris)	-	-	-	-	6	-
Triple morningglory (Ipomoea muricata)	-	-	2	3	2	3
Redroot pigweed ¹ (Amaranthus retroflexus)	4	2	6	4	6	6
Small flower morningglory (Jacquemontia tamnifolia)	-	-	4	3	4	3
Smooth pigweed ¹ (Amaranthus hybridus)	4	2	4	2	6	4
Spiny pigweed ¹ (Amaranthus spinosus)	4	2	6	4	8	6
Sunflower (Helianthus annuus)	6	8	8	10	10	12
Tall morningglory (Ipomoea purpurea)	3	3	4	4	4	4
Tall waterhemp ¹ (Amaranthus tuberculatus)	4	2	6	4	8	6
Toothed spurge (Euphorbia dentata)	-	-	4	4	4	4
Velvetleaf (Abutilon theophrasti)	4	3	6	5	8	6
Vernon mallow (Hibiscus trionum)	4	2	4	2	4	2
Wild Buckwheat (Polygonum convolvulus)	6	8	8	10	10	12
Wild mustard (Sinapis arvensis)	4	4	4	4	4	4

WEED SUPPRESSED

Canada Thistle (Cirsium arvense)	Not Recommended	8 Inch To Bud Stage	8 Inch To Bud Stage
BUCTREL® Herbicide + Athazine suppresses the growth by burning down the growth. Regrowth may occur.			

¹ Pigweeds (Amaranthus spp.) present in the field to be treated have been identified as triazine resistant biotypes, use BUCTREL® + atrazine Herbicide only at 3 pints/A. Control of pigweeds in the high plains areas of Texas and Oklahoma may not be satisfactory with BUCTREL® + atrazine Herbicide. Repeat applications may be necessary to achieve satisfactory control in these areas. Applications should be made when pigweeds do not exceed the 4 leaf stage and 2 inches in height.




BUCTRIL® + ATRAZINE TANK MIXTURE RECOMMENDATIONS

		APPLICATION TIMING AND SPECIFIC COMMENTS	
PRODUCT	RATE	CROP	WEEDS
BUCTRIL® + atrazine + Banvel ¹	1 1/2 - 2 pints/A + 1/8 - 1/4 pint/A	Apply to corn between the 3 leaf stage and before corn is 30 inches tall. Apply to sorghum between the 3 leaf stage and before sorghum is 15 inches tall. Do not apply in the boot stage. Use drop nozzles if crop is taller than 8 inches.	All weeds controlled by BUCTRIL® + atrazine at recommended rates of application plus improved control of pigweed, kochia, field bindweed suppression, and Canada thistle burndown.
	3 pints/A + 1/8 - 1/4 pint/A	Apply to corn between the 4 leaf stage and before corn is 30 inches tall. Apply to sorghum between the 4 leaf stage and before sorghum is 15 inches tall. Do not apply in the boot stage. Use drop nozzles if crop is taller than 8 inches.	
BUCTRIL® + atrazine 2,4-D (such as Weedone ² and Weedar ² brand)	1 1/2 - 2 pints/A + 1/16 - 1/4 lb ai/A	Apply to corn between the 3 leaf stage and before corn is 30 inches tall. Use drop nozzles if crop is taller than 8 inches. Apply to sorghum between the 3 leaf stage and before sorghum is 15 inches tall. Do not apply in the boot stage. Use drop nozzles if crop is taller than 8 inches.	All weeds controlled by BUCTRIL® + atrazine at recommended rates of application plus improved control of devil's claw, kochia, field bindweed suppression, and Canada thistle burndown.
	3 pints/A + 1/16 - 1/4 lb ai/A	Apply to corn between the 4 leaf stage and before corn is 30 inches tall. Use drop nozzles if crop is taller than 8 inches. Apply to sorghum between the 4 leaf stage and before sorghum is 15 inches tall. Do not apply in the boot stage. Use drop nozzles if crop is taller than 8 inches.	

Restrictions and Precautions: Corn (Field and Pop) and Sorghum (Grain and Forage)

- Postemergence applications of **BUCTRIL® + atrazine Herbicide** will not adequately control grasses. Therefore, it is recommended that a suitable grass control program be used to provide any required grass control.
- Addition of a spray additive or mixture with liquid fertilizers may cause excessive crop leaf burn.
- Do not use **BUCTRIL® + atrazine** on sorghum grown in sandy or loamy sand soils as excessive crop injury may occur.
- Seed corn producers should consult the respective seed corn company regarding tolerance of seed products of hybrid lines to **BUCTRIL® + atrazine Herbicide**.
- Do not cut crop for feed or graze within 30 days after application.
- A second application of **BUCTRIL® + atrazine Herbicide** may be applied if a new flush of weeds occur following the first application. The total cumulative rate should not exceed 6 pints/A per season.
- Do not use **BUCTRIL® + atrazine Herbicide** on any crop other than stated on this label.
- Application to grain sorghum growing under stress caused by minor element deficiency or to grain sorghum growing on highly calcareous soil may result in crop injury.
- Special care should be taken when using **BUCTRIL® + atrazine** and Banvel or 2,4-D tank mixtures to avoid off-target drift to sensitive crops.
- Tank mixtures with 2,4-D or Banvel can cause stalk brittleness to corn or sorghum. Winds or cultivation may cause breakage while crop is brittle.



CROP ROTATION

BUCTREL® + atrazine Herbicide contains atrazine at 1/4 lb active per pint of product. Due to the residual activity of the atrazine, injury to crops planted in treated fields the following season may occur. Consult your local extension service on the potential for atrazine carryover injury to rotational crops for your soil types and weather conditions encountered following application. If in doubt about your specific situation, a soil test prior to planting rotation crops is recommended to determine the amount of atrazine remaining and its potential to cause crop injury to the intended crop.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Do not store near fertilizers or seeds. Store at temperatures above 32° F. If exposed to freezing temperatures, store at temperatures above 55° F for 24 hours or until completely thawed. Shake well before using.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Metal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and Local authorities.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; that the directions, warnings, and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. **THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE, WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

¹ Registered trademark of Sandoz Crop Protection Corporation.

² Registered trademark of Rhône-Poulenc Ag Company.

³ Registered trademark of Delavan Corporation.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

BUCTREL is a registered trademark of Rhône-Poulenc.

BUCTREL + atrazine Herbicide (PENDING) SUBMITTED 12/89.

