



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 14 1992

Danielle A. Larochelle
RHONE-POULENC AG COMPANY
BOX 12014
RESEARCH TRIANGLE PARK, NC 27709

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 08/09/93 In Response to PR Notice 93-7
EPA Reg. No. 264-481
CHIPCO 26019

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

CHIPCO[®] brand 26019 Fungicide

30F9

A Fungicide For The Prevention And Control Of Certain Diseases Of Turfgrass And Ornamentals.

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide).....50.0%

INERT INGREDIENT:.....50.0%

EPA Reg. No. 264-481

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IN CASE OF CONTACT: Wash skin with soap and water.

FOR EYES: Flush with water for 15 minutes and get medical attention.

ACCEPTED
with COMMENTS
In EPA Letter Dated

JAN 1 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

264-481

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

TURF

In order to assure maximum crop tolerance and disease control, follow recommendations on this label and all the precautions and limitations of the package label.

CHIPCO® brand 26019 is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, home lawns, and institutional areas where fine turf is grown.

BEST AVAILABLE COPY

When used in conjunction with good turf management practices, CHIPCO® brand 26019 Fungicide is effective in controlling the following diseases:

50F9

Spring, Summer And Fall Disease

Dollar Spot, Brown Patch, Fusarium Blight, and **Necrotic Ring Spot on all common turfgrasses. Helminthosporium Leaf Spot, Corticium Red Thread and Melting Out are controlled on Bluegrass, Bentgrass, Perennial Ryegrass and Bermudagrass.

Winter Diseases: Fusarium Patch (Pink Snow Mold) and Gray Snow Mold

Apply the recommended rates as indicated in the table in 1 to 10 gallons of water per 1000 square feet. Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

** Not currently registered in California.

TURF

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

DISEASE	INTERVAL OF APPLICATIONS	RATE OZ. PRODUCT/ 1000 FT ²
Dollar Spot (Lanzia spp. and Moellerodiscus spp.) Brown Patch (Rhizoctonia solani) Helminthosporium: Leaf Spot and Melting Out (Drechslera spp.)	Greens and Tees: Repeat at 14 to 21 day intervals as long as required. Fairways and Other Turf Areas: Repeat at 22 to 28 day intervals as long as required.	1.5 to 2.0 NOTE: On Fairways, for Dollar Spot control, use 1 oz./1000 ft ²
Fusarium Blight (Fusarium spp.) **Necrotic Ring Spot (Leptosphaeria korrae)	Use only preventative foliar applications when conditions first become favorable for disease development. Two additional applications should be made at 2 to 3 week intervals.	4.0
Fusarium Patch (Microdochium nivalis) Pacific Northwest Only (West of the Cascade Mountains)	Repeat at 14 to 21 day intervals as long as required.	2.0 to 4.0
Gray Snow Mold (Typhula spp.) Pink Snow Mold (Microdochium nivalis)	One application before first permanent snow cover. If possible, another application during a mid-winter thaw.	2.0 to 4.0
Corticium Red Thread (Laetisaria fuciformis)	Use as a preventative every 14 days as long as required.	2.0

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases.

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24 hour waiting period following treatment is preferred.

Do not mix with any sticker, extender, or wetting agent.

Do not graze animals on treated turf.

Do not feed clippings from treated turf to livestock or poultry.

** Not currently registered in California.

TANK MIXTURES**

** Not currently registered in California.

ADDITIONAL DISEASE CONTROL

Gray Snow Mold: For control of Gray Snow Mold (Typhula spp.) in areas where continuous snow cover occurs, apply CHIPCO® brand 26019 Fungicide at 2 to 4 ounces of product per 1000 square feet with Daconil® 2787 Flowable Fungicide at 8 fluid ounces per 1000 square feet or Daconil 2787® WG at 4 1/2 ounces per 1000 square feet. Application must be made in autumn before snow covers occurs. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 square feet). Use the higher rate if the turf remains frozen prior to snow cover. For optimal control, reapply this treatment if a winter thaw and loss of snow cover occurs.

Pythium Blight: If turf is threatened by additional diseases, CHIPCO® brand 26019 Fungicide is compatible with most commonly used fungicides. Pythium blight can be controlled by the tank-mixing of CHIPCO® ALIETTE® brand Fungicide with CHIPCO® brand 26019 Fungicide. If a tank mixture is used, follow label directions for the use of that product and apply at the rate recommended for control of the target disease organism.

BEST AVAILABLE COPY

FIELD, LANDSCAPE AND GREENHOUSE ORNAMENTALS AND *CONIFER NURSERIES

CHIPCO® brand 26019 Fungicide is a broad spectrum fungicide that may be applied safely to a wide range of ornamental flowering and foliage plants, either as a foliar spray, drench or dip. Please read specific instructions and use only as directed.

RECOMMENDED FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL

CHIPCO® brand 26019 Fungicide is recommended for use on a wide variety of container and field grown flowering and foliage ornamentals as follows :

DISEASES

- | | |
|--------------------------------------|---|
| 1. Aerial Web Blight | (Rhizoctonia sp.) |
| 2. Alternaria Leaf Blight | (Alternaria euphorbiae) |
| 3. Alternaria Leaf Spot | (Alternaria panax, Alternaria tenuissima) |
| 4. Botrytis Blight | (Botrytis sp.) |
| 5. Fusarium Leaf Spot | (Fusarium moniliforme) |
| 6. Helminthosporium Leaf Spot | (Helminthosporium sp.) |
| 7. Rhizoctonia stem and root rot | (Rhizoctonia sp.) |
| 8. Ink Spot | (Drechslera iridis) |
| 9. Tulip Fire | (Botrytis tulipae) |
| 10. Alternaria Leaf Blight | (Alternaria zinniae) |
| 11. Ray Blight | (Ascochyta chrysanthami) |
| 12. Fusarium Corn rot | (Fusarium oxysporum) |
| 13. *Daffodil Leaf Scorch | (Stagnospora curtissi) |
| 14. *Blossom Blight | (Monilinia fructicola) |
| 15. *Botrytis Storage Rot | (Botrytis sp.) |
| 16. *Cylindrocladium Blight and Wilt | (Cylindrocladium scoparium) |

* Not currently registered in California.

PLANT TOLERANCE: Plant tolerances to CHIPCO® brand 26019 Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to CHIPCO® brand 26019 Fungicide. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

ORNAMENTALS

- | | | |
|---------------------------------|-----------------------------|--------------------------------|
| Ageratum (1 to 7) | Dieffenbachia (1 to 7) | Pansy (1 to 7) |
| Ajuga (1 to 7) | Dizygotheca (1 to 7) | Peach (ornamental) (1 to 7) |
| Almond (ornamental) (1 to 7) | Dogwood (1 to 7) | Peperomia (1 to 7) |
| Alyssum (1 to 7) | Dracena (1 to 7) | Periwinkle (1 to 7) |
| Andromeda (1 to 7) | English Ivy (1 to 7) | Philodendron (1 to 7) |
| Aphelandra (1 to 7) | Episcia (1 to 7) | Phlox (1 to 7) |
| Artemisia (1 to 7) | Euonymous (1 to 7) | Pilea (1 to 7) |
| Aster (1 to 7) | Ficus (1 to 7) | Pine (1 to 7) |
| Azalea (1 to 7, 16) | Forsythia (1 to 7) | Pitosporum (1 to 7) |
| Boxwood (1 to 7) | Gazania (1 to 7) | Plum (ornamental) (1 to 7, 14) |
| Cactus (1 to 7) | Geranium (1 to 7) | Poinsettia (1 to 7) |
| Calendula (1 to 7) | Gladiolus (1 to 7, 12) | Poppy (1 to 7) |
| Carnation (1 to 7) | Gloxinia (1 to 7) | Pothos* (1 to 6) |
| Cherry (ornamental) (1 to 7) | Gypsophila (1 to 7) | Primrose (1 to 7) |
| Chrysanthemum (1 to 7, 11) | Hawthorn (1 to 7) | Privet (1 to 7) |
| Cirreraria (1 to 7) | Holly (1 to 7) | Protea (1 to 7) |
| **Cistena Plum (1 to 7, 14) | Hoya (1 to 7) | Pyracantha (1 to 7) |
| Coleus (1 to 7) | Hydrangea (1 to 7) | Rhododendron (1 to 7, 16) |
| Columbine (1 to 7) | Impatiens* (1 to 7) | Rose Tree of China (1 to 7) |
| Coral Bells (Heuchera) (1 to 7) | Iris (1 to 8) | Rose (1 to 7, 15) |
| Crape Myrtle (1 to 7) | Juniper (1 to 7) | Salvia (1 to 7) |
| Crassula (1 to 7) | Kalanchoe (1 to 7) | Schefflera (1 to 7) |
| Croton (1 to 7) | Lillies (1 to 7) | Snapdragon (1 to 7) |
| Cyclamen (1 to 7) | Lipstick vine (1 to 7) | Statice (1 to 7) |
| **Daffodils (1 to 7, 13) | (Aeschynanthus) | Tree Ivy (1 to 7) |
| Dahlia (1 to 7) | Marigold (1 to 7) | Tulip (1 to 7, 9) |
| Delphinium (1 to 7) | Monarda (Bee Balm) (1 to 7) | Viburnum (1 to 7) |
| Deutzia (1 to 7) | Pachysandra (1 to 7) | Violet (1 to 7) |
| Dianthus (1 to 7) | Palm (1 to 7) | Zinnia (1 to 7, 10) |

** Not currently registered in California.

*NOTE: Do not use CHIPCO® brand 26019 Fungicide as a soil drench on Impatiens, and Pothos.

Do not use CHIPCO® brand 26019 Fungicide on Spathiphyllum.

BEST AVAILABLE COPY

Fungicide as a foliar spray to run-off, at the following rates and intervals, when conditions become favorable for disease development.

DISEASE	POUNDS PRODUCT/100 GALS.	INTERVAL OF APPLICATION	USE DIRECTIONS
Aerial Web Blight (<i>Rhizoctonia sp.</i>)	1.0 to 2.0	7 to 14 Days	Spray until wet using no more than 400 gallons of spray per acre.
Alternaria Leaf Blight (<i>Alternaria zinniae</i>)			
Alternaria Leaf Blight (<i>Alternaria euphorbiae</i>)			
Alternaria Leaf Spot (<i>Alternaria panax</i>) (<i>Alternaria tenissima</i>)			
Botrytis Blight (<i>Botrytis sp.</i>)			
Fusarium Leaf Spot (<i>Fusarium moniliforme</i>)			
Helminthosporium Leaf Spot (<i>Helminthosporium sp.</i>)			
Ink Spot (<i>Drechslera iridis</i>)			
Ray Blight (<i>Ascochyta chrysanthami</i>)			
Tulip Fire (<i>Botrytis tulipae</i>)			
*Daffodil Leaf Scorch (<i>Stagnospora curtissi</i>)			
*Blossom Blight (<i>Monilinia fructicola</i>)			

* Not currently registered in California.

Under severe disease pressure, use the highest recommended rate and/or the shortest spray interval.

NOTE: A resin based surfactant should be added to foliar sprays for use on iris, lilies, and tulips.

HOW TO USE CHIPCO® BRAND 26019 FUNGICIDE AS A DRENCH

Apply CHIPCO® brand 26019 Fungicide as a drench at seeding and/or after transplanting for *Rhizoctonia* control at the following rates and interval:

DISEASE	PRODUCT/100 GALS.	INTERVAL OF APPLICATION
Rhizoctonia Stem and Root Rot (<i>Rhizoctonia spp.</i>)	6 1/2 ounces Apply 1 to 2 pints of solution per square foot.	14 Days

NOTE: Do not use CHIPCO® brand 26019 Fungicide as a drench on impatiens, and pothos.

Do not use CHIPCO® brand 26019 Fungicide on *Spathiphyllum*.

BEST AVAILABLE COPY

HOW TO USE CHIPCO® BRAND 26019 FUNGICIDE AS A DIP

PLANT SPECIES	DISEASE	POUNDS PRODUCT 100 GALLONS	DIP DURATION	DIRECTIONS
Rose	*Botrytis Storage Rot (<i>Botrytis sp.</i>)	1.0	5 Minutes	Dip bare root roses prior to cold storage.
Azalea and Rhododendron	*Cylindrocladium Blight and Wilt (<i>Cylindrocladium scoparium</i>)	1.0	5 Minutes	Dip cuttings prior to planting.
Gladiolus	Fusarium Corm Rot (<i>Fusarium oxysporum</i>)	2.0	5 Minutes	Dip corms prior to storage.

*Not currently registered in California.

DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS

Do not use through sprinkler irrigation systems in California.

Apply this product only through sprinkler irrigation systems including center pivot and solid set. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of CHIPCO® brand 26019 Fungicide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of CHIPCO® brand 26019 Fungicide, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of CHIPCO® brand 26019 Fungicide per 1 to 4 gallons of water are recommended) Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of CHIPCO® brand 26019 Fungicide into the irrigation water line so as to deliver the desired rate per acre. The suspension of CHIPCO® brand 26019 Fungicide should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with CHIPCO® brand 26019 Fungicide has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

BEST AVAILABLE COPY

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warning and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

CHIPCO is a registered trademark of Rhône-Poulenc.

CHIPCO® 26019 Fungicide (PENDING) Submitted ___/___/93.

BEST AVAILABLE COPY