

APP 11 1994

Ms. Lizbeth R. Simila
Rhone-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Dr.
Research Triangle Park, NC 27709

Dear Ms. Simila:

Subject: CHIPCO® WEEDONE® DPC Amine Herbicide
EPA Registration No. 264-519
Application Dated April 1, 1994, Request for
Label Amendments, To Revise the Forestry Site
Preparation Directions and To Add WPS
Required Labeling in Response To Agency
Guidance for WPS Labeling Revisions for 2,4-D
Products, Dated September 8, 1993

The subject proposed labeling amendments have been reviewed and found acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended provided that you:

1. Revise the Environmental Hazards section by adding the statement "This product is toxic to aquatic invertebrates."
2. Submit five (5) printed copies of the final printed label before releasing the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

E.Wilson:Diskette ABC10: 04-12-94

BEST AVAILABLE COPY

CHIPCO® WEEDONE® DPC

S. Wilson
RL

Amine Herbicide

For Professional Weed Control in Ornamental Turf and Woody Plants in Non-Crop Areas. For Sale to and Use By Professional or Commercial Personnel Only

ACTIVE INGREDIENTS:

2-(2,4-Dichlorophenoxy)propionic acid, dimethylamine salt* 22.0%
2,4-Dichlorophenoxyacetic acid, trisopropanolamine salt** 34.5%

INERT INGREDIENTS: 43.5%

*2-(2,4-Dichlorophenoxy)propionic acid equivalent 18.5% by weight or 1.69 pounds per gallon.

**2,4-Dichlorophenoxyacetic acid equivalent 18.5% by weight or 1.82 pounds per gallon.

*Isomer specific by AOAC Method No. 6-D01-5

EPA Reg. No. 264-519

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For **PRODUCT USE** Information Call 1-800-334-9745

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or poison control center. Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting.

IF ON SKIN: Wash skin with plenty of soap and water and remove contaminated clothing. Get medical attention. Wash clothing separately before reuse.

NOTE TO PHYSICIAN

No specific antidote is available. Probable mucosal damage may contraindicate the use of gastric lavage. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

BEST AVAILABLE COPY PRECAUTIONARY STATEMENTS

DANGER

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

NOTE: (1) For containers over 1 gallon, but less than 5 gallons, mixers and loaders who do not use a mechanical transfer system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the required PPE. (2) For containers of 5 gallons or more, a mechanical transfer system (probe and pump) must be used for transfer of contents of the container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed by the mechanical system. This mechanical system must be used in a manner that meets the requirements listed in the Worker Protection Standard for agricultural pesticides [40 CFR 170.240(d)(4)]. The handler PPE requirements may be reduced or modified as specific in the WPS.

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE (personal protective equipment) may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D or 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D or 2,4-DP pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

CLEANING OF EQUIPMENT: Do not use the same spray equipment for other purposes unless thoroughly cleaned. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from well and other water sources.

Do not contaminate water used for irrigation or domestic purposes (except as specifically recommended on this label) especially in areas where grapes, cotton, tomatoes or other susceptible plants are grown. Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops especially grapes, tomatoes, tobacco and cotton.

Do not apply when weather conditions favor drift from treated areas. Do not apply CHIPCO® WEEDONE® DPC Amine Herbicide directly to, or permit to drift onto cotton, okra, grapes tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by CHIPCO® WEEDONE® DPC Amine Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions, a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion, consult with local weather services before making an application.

Use coarse sprays to minimize drift. Do not apply with hollow cone-type insecticide or other nozzles that produce fine spray droplets. Drift from aerial or ground application may be reduced by: (1) applying as near to the target as possible in order to obtain coverage; (2) by increasing the volume of spray mix per acre; (3) by decreasing the pounds of pressure at the nozzle tips; and (4) by using nozzles which produce a coarse spray pattern; (5) by not applying when wind is blowing toward susceptible crops or valuable plants.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in any manner inconsistent with its labeling. Read entire label before using product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specified instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks and protective eye wear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to ornamental turf uses (golf courses, cemeteries, parks and other turf grass areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application.

Do not enter treatment areas until sprays have dried.

STORAGE & DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL

Open dumping is prohibited. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL

Do not reuse empty containers. Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by state or local authorities, by burning. If container is burned, stay out of smoke.

GENERAL CAUTIONS AND RESTRICTIONS

Do not apply CHIPCO® WEEDONE® DPC Amine Herbicide through any type of irrigation system.

Do not use in or near a greenhouse.

Avoid treatment when turf is stressed, as from drought or high summer temperatures (above 90° F).

IMPORTANT: Do not use CHIPCO® WEEDONE® DPC Amine Herbicide to control weeds in flower or vegetable beds, in shrub or ornamental plantings, or on lippia or dichondra lawns as severe injury may result.

A separate sprayer should be kept for killing weeds because 2,4-D and related chemicals are difficult to clean from equipment. Spray when the air is calm to avoid spray drift that might injure nearby desirable plants.

MIXING LIQUID FERTILIZERS WITH AMINE FORMULATIONS:

CHIPCO® WEEDONE® DPC Amine Herbicide may be combined with most liquid fertilizers. However, under certain conditions (such as prolonged standing, especially under low temperatures) insoluble salts may form and cause nozzle clogging and spotty or poor weed control. If in doubt about compatibility or solubility, test a small sample in the dilution ratio planned for application.

To prepare the spray CHIPCO® WEEDONE® DPC Amine Herbicide must first be premixed with water. For liquid nitrogen use a mix of 1 part CHIPCO® WEEDONE® DPC Amine Herbicide plus 4 parts of water; For other fertilizers use 1 part of CHIPCO® WEEDONE® DPC Amine Herbicide with 50 to 60 parts of water. Add the premix to the fertilizer while maintaining continuous agitation during both the mixing and spraying operations. Apply the same day it is prepared.

GENERAL INFORMATION

This material forms a true solution in water. It does not require additional agitation after the initial thorough mixing, and does not volatilize under common spraying conditions.

Note: Local conditions and application regulations vary and may affect use of this herbicide. Consult local agricultural experiment station or extension service weed specialists and state regulatory agencies for recommendations in your area.

WEED CONTROL IN ORNAMENTAL TURF

Use CHIPCO® WEEDONE® DPC Amine Herbicide on lawns, golf courses, athletic fields, parks and other similar ornamental turf areas.

CHIPCO® WEEDONE® DPC Amine Herbicide for professional weed control in ornamental turf is also effective on poison ivy, honeysuckle, brambles and many noxious perennial weeds in rough areas such as along driveways, roadsides, fencerows and drainage ditchbanks.

Apply CHIPCO® WEEDONE® DPC Amine Herbicide for professional weed control in ornamental turf any time broadleaf weeds are growing actively. Dandelion, ground ivy and clovers respond best to treatment in fall or in spring before flower heads develop. To control wild garlic, spray it in early spring and late fall for two consecutive treatments. Winter weeds such chickweed and henbit should be treated in very early spring. Summer weeds such as knotweed and spotted spurge are most easily controlled while they are small. In areas with extended growing seasons, treatment in both spring and fall may be needed to control more resistant species.

For established turf, fall treatment fits into a good management program. Adequate fertilization and proper mowing should be combined with chemical weed control to thicken the turf after weeds have died and thus discourage more weeds from invading.

On turf to be over-seeded in the fall, treatment should be scheduled for at least 4 weeks before the seeding date. A good rain (1/4 inch or more) or deep watering should occur during this period.

Fall-seeded turf may be treated the following spring. Spring-seeded turf may be treated after the grasses have germinated and been cut at least twice, generally 6 to 10 weeks after seeding, depending on species and growing conditions.

WOODY PLANT CONTROL IN NON-CROP AREAS

CHIPCO® WEEDONE® DPC Amine Herbicide is specifically designed for utilities and other industrial users to control woody plants.

CHIPCO® WEEDONE® DPC Amine Herbicide controls many noxious perennial weeds on non-cropped land, such as along highways, drainage ditchbanks and fence rows.

BEST AVAILABLE COPY

WEEDS CONTROLLED BY CHIPCO® WEEDONE® DPC AMINE HERBICIDE FOR PROFESSIONAL ORNAMENTAL TURF WEED CONTROL:

- | | | |
|------------------------------|--|--------------------|
| bindweed | hawkweed | purslane |
| black medic | healall | ragweed |
| brambles | henbit | sheep sorrell |
| buckhorn plantain | honeysuckle | shepherdspurse |
| bull thistle | jimsonweed | smartweeds |
| burdock | knotweed | soliva |
| buttercups | kochia | sowthistle |
| Canada thistle | kudzu | speedwell (annual) |
| Carolina geranium | lambquarters | spurge |
| chickweeds(common, mouseear) | little starwort | sumac |
| chicory | mallow | verbain |
| clovers | momingglory | vetch |
| cocklebur | mustards | violet |
| cudweed | oxalis (yellow woodsorrel) | wild aster |
| dandelion | pennywort | wild blackberry |
| dock | peppergrass | wild carrot |
| evening primrose | pigweed | wild garlic |
| false dandelion | plantains (narrow, buckhorn;broadleaf) | wild geranium |
| fleabane | poison ivy | wild lettuce |
| Florida pusley | wild raspberry | wild onion |
| frenchweed poison oak | poorjoe | wild radish |
| ground ivy | povertyweed | yarrow |

This product is not effective on perennial veronicas or on weed grasses.

WOODY PLANTS CONTROLLED BY CHIPCO® WEEDONE® DPC AMINE HERBICIDE IN NON-CROP AREAS:

- | | |
|---------------|-------------------|
| Alder | Multiflora Rose |
| Ash | Oaks |
| Aspen | Osage Orange |
| Bigleaf Maple | Pine |
| Birch | Poison Ivy |
| Blackberry | Poison Oak |
| Blackjack Oak | Poplar |
| Black Locust | Red Elder |
| Box elder | Red Elm |
| Brambles | Salmonberry |
| Buckbrush | Sand sagebrush |
| Ceanothus | Sand Shinnery Oak |
| Chamise | Serviceberry |
| Coffeeberry | Snowberry |
| Currant | Spruce |
| Elderberry | Sumac |
| Fir | Sycamore |
| Gooseberry | Tulip Poplar |
| Greenbrier | Wild Cherry |
| Hemlock | Wild Grape |
| Honeysuckle | Willow |
| Locust | Winged Elm |
| Manzanita | Yerba Santa |

BEST AVAILABLE COPY

6078

SPECIFIC USE DIRECTIONS

ORNAMENTAL TURF

To control most annual and perennial broadleaf weeds in ornamental turf, apply CHIPCO® WEEDONE® DPC Amine Herbicide professional ornamental turf weed control at the rate of 3 to 4 pints per acre (1 to 1.5 ozs. per 1000 sq. ft.) in enough water for good wetting (25 to 100 gallons per acre).

Treatment at this rate may injure bentgrass, St. Augustinegrass, centipedegrass, carpetgrass and newly seeded turf. If it is necessary to control weeds in these grasses, use half the rate recommended (1.5 to 2 pints per acre or 1/2 to 3/4 oz. per 1000 sq. ft.) and make a second half-rate treatment 2 to 3 weeks later. Do not use CHIPCO® WEEDONE® DPC Amine Herbicide on bentgrass greens or tees.

Resistant weeds such bindweed, oxalis(yellow woodsorrel), sheep (red) sorrel, soliva, violets and yarrow may require up to 6 pints per acre (2.25 ozs. per sq. ft.) and should be retreated whenever new growth appears.

The maximum number of broadcast applications per treatment site is 2 per year. This does not exclude spot treatments.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until sprays have dried.

WOODY PLANT CONTROL IN NON-CROP AREAS

FOLIAGE STEM TREATMENT

This is the standard method for high volume sprays along fencerows, highways and rights-of-ways. Use it as a "first spray" on thick brush composed of mixed species.

Apply spray to both foliage and stems of all plants from the time foliage is fully developed until plants begin to dormant. For effective control all leaves, stems and suckers must be thoroughly wet to ground line. Some regrowth may be expected on resistant species, such as ash, maple, oak and persimmon.

Add 1 to 1 1/2 gallons of CHIPCO® WEEDONE® DPC Amine Herbicide to 100 gallons of water. Mix thoroughly before spraying. Use 200 to 600 gallons of spray per acre depending on height and density of brush.

FRESH CUT SURFACE TREATMENT

Stumps: This treatment may be used anytime of the year, however, it is more effective when applied as a quickly as possible after trees are cut. The whole stump is to be sprayed, especially exposed roots and bark. A complete soaking is essential for effective control.

This procedure should be used after original removal. It is the initial procedure in a chemical brush control operation on newly denuded highways and rights-of-way. The spray is most effective and profitable on stumps 3 to 4 inches and larger.

Mix 3 to 4 gallons of CHIPCO® WEEDONE® DPC Amine Herbicide in 100 gallons of water. Applications should be made with a low volume sprayer using a solid cone shaped nozzle with medium openings.

Concentrate Stump Treatment: For small (up to 3 inches diameter) stems, cut them as close to the ground as possible and apply undiluted CHIPCO® WEEDONE® DPC Amine Herbicide directly from the can to the surface of the freshly cut stump.

Frill: By using an ax, overlapping V-shaped indentations in a continuous ring are cut around the trunk near its base. The bark is to be cut through, but the chips are not to be removed. This is the advised treatment for trees 5 to 6 inches in diameter and larger. Newly cut frills can be treated anytime of the year. Mix 3 to 4 gallons of CHIPCO® WEEDONE® DPC Amine Herbicide in 100 gallons of water. Apply as much of the solution as the frills can accommodate without carelessly dissipating the chemical.

SPOT TREATMENT

For spot spraying with backpack sprayer mix 1 cup (8 ozs.) CHIPCO® WEEDONE® DPC Amine Herbicide with 5 gallons of water. Wet brush stems and foliage thoroughly.

FENCE ROW APPLICATION

To control mixed brush, perennial and annual broadleaf weeds use one of the application methods such the foliage stem method described on this label. Some regrowth may be expected on resistant species such as ash, maple, oak and persimmon.

FORESTRY SITE PREPARATION

To reduce competition from mixed trees and brush before planting forest trees. Apply 1 gallon CHIPCO® WEEDONE® DPC Amine Herbicide for easy to control species such as aspen, hazel, alder, sassafras, sumac, black locust, willow and other similar species. Apply 2 gallons CHIPCO® WEEDONE® DPC Amine Herbicide to control more difficult species such as oaks, sweet gum, black gum, hickory and tulip poplar.

Mix with water to make 10 gallons of total spray solution when applying as an aerial spray or 20 gallons of total solution when applying as a ground spray. Do not apply to established plantations as this spray mix will injure planted conifers.

BEST AVAILABLE COPY

PINE RELEASE

To control hardwood brush and release Northern conifers such as red pine, Jack pine, white pine and white spruce.

Apply 2 to 4 quarts CHIPCO® WEEDONE® DPC Amine Herbicide as a broadcast spray in 10 gallons of water per acre. Make applications in mid summer after pine height growth is complete and the conifer buds are set.

CHIPCO® WEEDONE® DPC Amine Herbicide will control aspen, birch, cherry, alder, hazel, oaks and similar species. CHIPCO® WEEDONE® DPC Amine Herbicide will not, at these rates, provide satisfactory control of red maple, sugar maple or ash.

AERIAL OR GROUND APPLICATION TO CONTROL MIXED BRUSH INCLUDING CONIFERS AND ROOT SUCKERING SPECIES ON UTILITY, RAILROAD AND HIGHWAY RIGHTS-OF-WAY, FENCEROWS AND OTHER NON-CROP AREAS

CHIPCO® WEEDONE® DPC Amine Herbicide and Tordon Mixtures

Where red maple, conifers and root-suckering species such as sassafras, sumac, black locust or persimmon are the major problem, use 1 to 2 1/2 gallons CHIPCO® WEEDONE® DPC Amine Herbicide in combination with 1 to 2 1/2 gallons of Tordon 101 herbicide mixture or in combination with 3 to 6 pints of Tordon K herbicide. Dilute with water and apply in a total spray volume of 15 to 50 gallons per acre. Use lower rates to control brush on sandy soils or coarse soils with susceptible species such as aspen, cherry, willow and sumac. Use higher rates to control brush on sandy soils or coarse soils or fine soils or rocky terrain with more resistant species such as maple, oak, pine and red cedar. Aerial applications of the tank mixture should be made only with a helicopter mounted MICROFOIL® applicator or an equipment system providing equivalent drift control. Do not use these tank mixtures on drainage ditchbanks or for site preparation in forest plantations. Observe all restrictions, precautions and limitations on the labeling of each product used in tank mixtures.

**CHIPCO® WEEDONE® DPC Amine Herbicide and Banvel Herbicide Mixtures
HIGH VOLUME STEM FOLIAGE SPRAY**

Mix 1 to 1 1/2 gallons CHIPCO® WEEDONE® DPC Amine Herbicide with 1 to 2 quarts Banvel 4WS per 100 gallons total spray mix. Use the low rate for easy to kill brush such as alder, aspen, cherry, sycamore, tulip poplar and willows. Use the high rate to control species such as oak, ash, elm, pines, spruce and fir.

LOW VOLUME STEM FOLIAGE SPRAY- Aerial or Ground

Mix 2 to 3 gallons of CHIPCO® WEEDONE® DPC Amine Herbicide with 1 to 1 1/2 gallons of Banvel herbicide in 15 to 50 gallons of water per acre to control mixed brush including conifers and ash. Observe all restrictions, precautions and limitations on the labeling of each product used in tank mixtures.

8078

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

)
)

)
)

BEST AVAILABLE COPY

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

CHIPCO and WEEDONE are registered trademarks of RHONE-POULENC.
Tordon is a trademark of Dow Chemical Company.
Banvel is a trademark of Sandoz Crop Protection Company.
©1990 Rhône-Poulenc Ag Company
Made in U.S.A.

CHIPCO® WEEDONE® DPC Amine Herbicide (PENDING) Submitted 4/1/94.