

Reg # 264-530

PM-9

19 16

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 264-530	DATE OF ISSUANCE APR 07 1993
	TERM OF ISSUANCE Until Reregistration	
	NAME OF PESTICIDE PRODUCT Larvin Brand DF WSP Thiodicarb Insecticide/ Ovicide	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Rhone-Poulenc Ag Company
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, NC 27709

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c)(5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the label changes listed below before you release the product for shipment:
 - a. Add the phrase EPA Registration No. "264-530" to your label.
 - b. Include application rates for cotton and soybeans in the table titled Number of Water Soluble Packets to Use.
 - c. Submit a revised Confidential Statement of Formula (CSF) deleting "& others" from inert ingredient suppliers' names. A different supplier's formulation may or may not be acceptable, and therefore this phrase is not acceptable on the CSF. As discussed with you over the phone we are crossing this out from the CSF we currently have to make it acceptable. We have also added the date 2/16/93 column 21 of the CSF.

continued on next page

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL: **DOUGLAS H. EDWARDS**

DATE: **APR 07 1993**

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide & Rodenticide Branch
Registration Division (H7505C)

Enclosure

3816

LARVIN® brand DF WSP Thiodicarb

Insecticide In Water Soluble Packaging

ACTIVE INGREDIENT:
(THIODICARB) Dimethyl N, N' [thiobis [(methylimino) carbonyloxy]] bis [ethanimidothic ate] 80% by wt.
INERT INGREDIENTS..... 20% by wt.

Package Size: 40 pounds of product per box packaged as : 2.50 pound packets, 16 per box or
1.25 pound packets, 32 per box

EPA Reg. No. 264-LGN **EPA Est. No. 264-MO-02**

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: If patient is conscious and alert, give 2 to 3 glasses of water or milk to drink. If available, give one tablespoon of Syrup of Ipecac to induce vomiting. If vomiting does not occur within 20 minutes of the first dose, follow with a second and last dose of 1 teaspoon. Alternatively, induce vomiting by touching back of throat with finger. Do not make an unconscious person vomit. **GET MEDICAL ATTENTION.**

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. **GET MEDICAL ATTENTION.**

IF ON SKIN: Immediately flush skin with plenty of water while removing contaminated clothing and shoes. Wash clothes separately before reuse.

IF INHALED: Remove to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. **GET MEDICAL ATTENTION.**

GENERAL

Contact a physician immediately in all cases of suspected poisoning. If breathing stops, establish an airway, start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.** If poisoning is suspected in animals, contact a veterinarian.

NOTE TO PHYSICIAN

All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product contains a methyl carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. LARVIN® poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent over atropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are **NOT** recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Rhône-Poulenc Ag Company.

Consultation on therapy can be obtained at all hours by calling the Rhône-Poulenc emergency number: 1-800-334-7577

RECEIVED
Rhône-Poulenc
Letter Dated
APR - 7 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
264-530

49/16

PRECAUTIONARY STATEMENTS

WARNING HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be harmful or fatal if swallowed or inhaled. Illness may result rapidly following overexposure to this product. May be harmful or fatal if swallowed or inhaled. May be harmful if absorbed through the skin. May be irritating to eyes and may cause mild skin sensitization. Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Keep away from domestic animals.

Wear long-sleeved clothing, full-length trousers, head covering and protective gloves while using this product. Remove contaminated clothing daily and wash before reuse. Wash exposed skin thoroughly with soap and water after use and before eating or using tobacco. Discontinue use if allergic reaction occurs. Do not eat or use tobacco while using this product.

IN SEVERE CASES, CONVULSIONS, UNCONSCIOUSNESS AND RESPIRATORY FAILURE MAY OCCUR.

SIGNS AND SYMPTOMS OF OVEREXPOSURE

Abdominal cramps	Diarrhea
Excessive sweating	Muscle tremors
Pinpoint eye pupils	Vomiting
Blurred vision	Difficult breathing
Headache	Nausea
Salivation	Weakness
	Watery eyes

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, invertebrates and other wildlife. Do not apply directly to lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from area treated. This product is toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the treatment area.

GENERAL INFORMATION

LARVIN® brand DF WSP Thiodicarb Insecticide is a dry flowable concentrate (water dispersable granule) formulation that readily disperses in water for spraying by ground or air equipment. Directions on this label are based on tests and field experience relating to effectiveness, impact on environment and residues remaining in food and feed. READ THE ENTIRE LABEL AND OBSERVE ALL LABEL DIRECTIONS AND PRECAUTIONS BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not use this product except as directed on this label or on other Rhône-Poulenc Ag Company supplemental labeling for this product. Applications at higher rates or at more frequent intervals than is stated on the label may result in illegal residues in crops, meat and milk.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENT, STATEMENT OF PRACTICAL TREATMENT, and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (WARNING), area treated with LARVIN brand DF WSP Thiodicarb Insecticide, date of application, appropriate protective clothing, appropriate reentry interval, if applicable.



STORAGE AND DISPOSAL

STORAGE

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115°F (46°C). NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

If container is damaged, before clean-up, put on long-sleeved shirt, full-length trousers, head covering and protective gloves.

PESTICIDE DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Contamination with this product will render water, food or feed unfit for human or animal consumption. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Do not reuse outer bag. Dispose of outer bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACCIDENTS: In case of a major spill of LARVIN® brand DF WSP Thiodicarb Insecticide TELEPHONE (24 HOURS A DAY) IN THE U.S.A. 1-800-334-7577.

In case of minor spills, follow all precautions indicated above and cleanup immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

GENERAL DIRECTIONS

HOW TO USE LARVIN® BRAND DF WSP

Remove oil, rust, scale, pesticide residues and other foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent slotted strainers in spray system. To prepare for spraying, fill tank to approximately 1/2 the needed volume of water and start agitation. Open outer plastic bag and drop the water soluble packet(s) into the tank. Agitate to ensure thorough mixing for a minimum of 2 minutes or until bags and product are completely dispersed, while filling tank with remaining water. Do not store spray mixture for prolonged periods. If tank mixes are to be used, LARVIN® brand DF WSP Thiodicarb Insecticide must be fully dispersed in water first, followed by the addition of the intended tank-mix material.

COMPATIBILITY

Thiodicarb Insecticide when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using LARVIN® brand DF WSP Thiodicarb Insecticide in any tank mixture the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. THIS PRODUCT IS NOT COMPATIBLE WITH NON-EMULSIFIED VEGETABLE OR CROP OIL DILUENTS. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.**

NOTE: Be sure the water soluble packet is completely dissolved in the spray tank before adding any other ingredients. Thiodicarb Insecticide is hydrolytically sensitive to degradation of active ingredient by strong acids, strong bases, and certain heavy metal oxides and salts of certain fungicides. Degradation can cause reduced effectiveness, plant phytotoxicity and increased handling hazards. **DO NOT USE THIS PRODUCT IN SPRAY MIXTURES WITH THE ABOVE TYPE OF PRODUCTS INCLUDING MIXTURES WITH Maneb, Mancozeb (except Manzate 200 and Dithane M-45 wettable powders where mixtures are used immediately after mixing), Copper Count N-8, and Bordeaux mixture. DO NOT ADD LARVIN® BRAND DF WSP THIODICARB INSECTICIDE TO WATER WITH pH VALUES BELOW 3.0 OR ABOVE 8.5. If necessary, water should be buffered within this range before adding LARVIN® brand DF WSP Thiodicarb Insecticide.**

DIRECTIONS FOR AERIAL OR GROUND SPRAY APPLICATION

APPLICATION TIMING

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, Professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

APPLICATION INSTRUCTIONS

Apply a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground unless otherwise directed under crop specific directions. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground. For best results, it is important to obtain thorough and uniform spray coverage of the plant. Repeat application as needed to maintain control. Use higher dosage rates for heavy infestations, large larvae, or dense foliage. The specific length of residual control depends on environmental factors, plant growth, disease rate, and degree of insect infestation. Although LARVIN® has some contact activity, larvae not feeding on treated foliage will not be controlled. When banding, determine the amount of chemical to use per acre by dividing the band width by the row width and multiplying by the appropriate broadcast rate.

To clean the sprayer after use, drain and flush with water. Use rinsings on crop according to label instructions. Dispose of in an approved manner (See STORAGE AND DISPOSAL).

RECOMMENDED USES FOR BROCCOLI, CABBAGE AND CAULIFLOWER

SPRAY VOLUME FOR BROCCOLI, CABBAGE AND CAULIFLOWER: Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® BRAND DF WSP		
COLE CROPS (BROCCOLI, CABBAGE, CAULIFLOWER)	Alfalfa Looper	0.5	0.63	Spray as needed. As insect populations increase and/or new plant growth is rapid, the spray interval should be shorter. Use higher dosage rates for heavier infestations or large larvae. Use lower rates for light to moderate infestations. NOTE: Regional differences have been noted in the susceptibility of certain strains of diamondback moth to thiodicarb. If local experience indicates inadequate control, use an alternate pesticide.	Do not exceed 6.0 pounds of active ingredient (7.5 pounds of this product) per acre per season. Do not apply less than 7 days before harvest.
	Beet Armyworm	to	to		
	Cabbage Flea Beetle	0.75	0.94		
	Imported Cabbageworm				
	Cabbage looper	0.75	0.94		
	Diamondback moth	to 1.0	to 1.25		
	Cutworms	See "SPECIAL DIRECTIONS ON CUTWORMS"			

RECOMMENDED USES FOR COTTON

INTEGRATED PEST MANAGEMENT (IPM) USE OF THIS PRODUCT

LARVIN® Thiodicarb insecticide has both contact and residual ovicidal activity which can be effectively utilized in IPM programs (See ovicide section). LARVIN® Thiodicarb insecticide has been shown to leave substantial populations of beneficial insects and spiders after use. The lower rates allow for maximum beneficial survival and faster rebound of beneficial populations. Larvicidal and ovicidal control coupled with retention of beneficial insects and spiders can offer significant benefits to those producers utilizing integrated pest control programs.


SPRAY VOLUME FOR COTTON

LARVIN® brand DF WSP Thiodicarb insecticide should be applied in a minimum finished spray volume of 2 gallons per acre by aircraft and 5 gallons per acre by ground equipment. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum spray volume of 10 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® BRAND DF WSP		
COTTON	Ovicide: Cotton bollworm Tobacco budworm	0.125 to 0.25	0.16 to 0.31	Apply on a 3 to 5 day schedule when moth flights begin or sufficient eggs are present. Thorough spray coverage is necessary as eggs must be contacted by LARVIN® for activity to occur. Eggs laid on untreated foliage or not contacted by LARVIN® sprays will not be controlled. If significant number of larvae survive, use higher recommended LARVIN® rates or 0.125 to 0.25 lbs ai/A of LARVIN® in tank mixture combinations with organophosphates, pyrethroids or <i>Bacillus thuringiensis</i> such as DPEL® ES registered for use on cotton at their label recommended rates. Observe all restrictions and precautions which appear on the labels of all products to be tank mixed with LARVIN®	Do not allow livestock to graze fields. Do not apply less than 28 days before harvest. In Arizona, apply before bolls begin to open and do not exceed 1.8 pounds per acre per season.

RECOMMENDED USES FOR COTTON (CONTINUED)


SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® BRAND DF WSP		
COTTON (cont)	Larvicide: Cotton bollworm Tobacco budworm	0.6 to 0.9	0.75 to 1.12	In this rate range, LARVIN® is a direct contact and residual contact ovicide, as well as an effective larvicide. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Use the higher rate when insect larvae populations are heavier or when infestations are predominantly large larvae (>1/4 inch in size).	Do not allow livestock to graze fields. Do not apply less than 28 days before harvest. In Arizona, apply before bolls begin to open and do not exceed 1.8 pounds per acre per season.
	Larvicide (Cont.): Armyworms (such as: Beet, Fall, etc.) Cabbage looper Cotton leaf perforator Cotton leafworm Soybean looper Fleahopper (Suppression) Pink bollworm (Suppression) Plant bugs (Suppression)	0.6 to 0.9	0.75 to 1.12	As pest pressure increases, reduce intervals between applications and/or increase dosage rate as indicated in specific directions below. A rate of 0.75 pounds (0.60 lbs. active) should be used when: For low to moderate insect populations and when maximum survival of beneficial insects and spiders is desired. Rates of 0.94 to 1.12 pounds (0.75 to 0.90 lbs. active) should be used when: Insect populations are heavy or infestation is predominantly large larvae. Although LARVIN® has some contact activity, larvae must feed on treated foliage to be controlled. Applications for suppression of Pink bollworm should be timed for the presence of adult moths.	
	Boll weevil (Suppression)	0.6 to 0.9	0.75 to 1.12	LARVIN® will reduce boll weevil populations comparable to most pyrethroids. Light Boll Weevil Populations: When spraying LARVIN® on a 5 to 7 day worm control schedule, control of boll weevil will usually be adequate to maintain boll weevil populations below established economic thresholds. Moderate to Heavy Boll Weevil Populations: When spraying LARVIN® on a worm control schedule, application of a registered boll weevil control product and/or tank mix of LARVIN® plus a registered boll weevil product should be used. A 3 to 5 day interval between sprays may be required to maintain boll weevil control under heavy infestations.	
	Cutworms			See "SPECIAL DIRECTIONS ON CUTWORMS"	



RECOMMENDED USES FOR LEAFY VEGETABLES

SPRAY VOLUME FOR LEAFY VEGETABLES: Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® BRAND DF WSP		
LEAFY VEGETABLES (Amaranth, arugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible leaved, garland), cress (garden, upland), dandelion, dock, endive, Florence fennel, lettuce (head, leaf), orach, parsley, purslane (garden, winter), rhubarb, spinach(leaf, fine, new zealand), Swiss chard	Alfalfa looper	0.5 to 0.75	0.63	Spray as needed. Use the higher dosage rates for heavier infestations of larger larvae. Use the lower rates for light to moderate infestation.	Do not exceed 1.5 pounds of active ingredient (1.9 pounds of LARVIN® brand DF WSP) per acre per season. Do not apply less than 14 days before harvest.
	Amyworms (Such as: Beet, Fall, Southern) Heliothis zea (Bollworm, Corn earworm, Tomato fruitworm)		to 0.94		
	Cabbage looper	0.75	0.94		
Cutworms		See "SPECIAL DIRECTIONS ON CUTWORMS"			



RECOMMENDED USES FOR SOYBEANS (STANDARD AND LOW SPRAY VOLUMES)

STANDARD SPRAY VOLUME FOR SOYBEANS

Apply in a minimum finished spray volume of 2 gallons per acre by air or 5 gallons per acre by ground. Under extreme pest populations or dense foliage, use a minimum spray volume of 5 gallons per acre by air and a minimum of 10 gallons per acre by ground.

LOW SPRAY VOLUME FOR SOYBEANS


Apply in a minimum finished spray volume of 1 gallon per acre by aircraft. Only use the low volume aerial application (i. e. minimum of 1 GPA) where that spray volume will provide adequate coverage of the crop.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVINE BRAND DF WSP		
SOYBEAN (Standard Spray Volume Applications)	Podworms (Bollworm, Corn earworm, Tobacco budworm)	0.25 to 0.40	0.31 to 0.50	Use the lower rates for low to moderate populations and maximum protection of beneficials. For loopers, use low rate (0.45) for newly hatched worms, moderate rate (0.60) for young worms, and use high rates (0.75) for large worms. Use at least 5 gallons of finished spray material per acre if applied by air.	Do not feed forage, hay or straw to livestock. Do not apply less than 28 days before harvest.
	Amyworms (Such as: Beet, Fall, Southern, Yellow-striped, etc.) Green cloverworm Velvetbean caterpillar Woollybear caterpillar				
	Soybean looper Cabbage looper Mexican bean beetle Bean leaf beetle Stinkbugs (Suppression) Three-Cornered alfalfa hopper (Suppression)	0.45 to 0.75	0.56 to 0.94		
	Cutworms	See "SPECIAL DIRECTIONS ON CUTWORMS"			
SOYBEAN (Low Spray Volume Applications)	Podworms (Bollworm, Corn earworm, Tobacco budworm) Green Cloverworm Velvetbean caterpillar Amyworms (such as Fall, Beet, Southern, etc.) Woollybear caterpillar	0.25 to 0.40	0.31 to 0.50	For best results, use the lower rates for low to moderate populations and maximum protection of beneficials. When applying at 1 gallon of spray volume per acre by air, use 1 quart of emulsified crop or vegetable oil (e.g., emulsified corn, soybean or cottonseed oil). Water alone may be used as a diluent for spray volumes of 2 gallons or higher per acre. TANK MIX DIRECTIONS FOR EMULSIFIED CROP OR VEGETABLE OILS: Fill spray tank with approximately 3/8 of the water required for the spray mixture. Add the required amount of LARVINE BRAND DF WSP to the spray tank with the agitator running. Then add the desired amount of emulsified crop or vegetable oil. Continue agitation while adding the remainder of water.	FOR USE ONLY IN MARYLAND, VIRGINIA, NORTH CAROLINA, DELAWARE, SOUTH CAROLINA, GEORGIA, ALABAMA, MISSISSIPPI, LOUISIANA, ARKANSAS AND MISSOURI. Do not feed forage, hay, or straw to livestock. Do not apply less than 28 days before harvest.

RECOMMENDED USES FOR SWEET CORN

SPRAY VOLUME FOR SWEET CORN: Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground. For aerial applications, only water may be used as a diluent and total spray volume must be at least 2 gallons per acre.

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVIN® BRAND DF WSP		
SWEET CORN (Florida Fresh Market Only)	Corn earworm European corn borer Armyworms (Such as Fall, Beet, etc). Western bean cutworm	0.5 to 0.75	0.63 to 0.94	<p>Whorl Feeders: Spray as needed. As insect populations increase and/or when new plant growth is rapid, the spray interval should be shorter. For best results, use spray volumes of 15 gallons or more per acre applied by ground equipment using solid or hollow cone nozzles directed into the plant whorl.</p> <p>Insects Attacking Silks And Ears: Apply at 1 to 7 day intervals starting when silks first appear and continuing until harvest, if necessary. As insect populations increase and/or silk growth is rapid, the spray interval should be shorter. Use the higher dosage rate and apply at 1 to 2 day intervals following periods of heavy moth flights. Follow local recommendations for proper timing.</p> <p>Foliage Feeders: Use at least 2 gallons finished spray volume per acre by air or 10 gallons per acre by ground.</p>	After silk initiation, do not exceed 7.5 pounds of active ingredient (9.38 pounds of this product) per acre per season. Do not allow livestock to graze treated field. Do not feed treated corn silage (green plant) or fodder to livestock. Ears may be harvested on day of last application.
	Fall armyworm	0.25 to 0.50	0.31 to 0.63	<p>Low dosage/high frequency management program for Tassel, silk, and ear protection only: Start LARVIN® applications when 1% of plants have initiated tassel stage and spray every day until all plants are completely tasseled (minimum of 5 consecutive sprays). Subsequently, maintain a 1 to 2 day spray schedule for protection of silk and ears until harvest. Use a minimum of 15 gallons of finished spray volume per acre by ground or 5 gallons per acre by air.</p> <p>For best results, whorl stage infestations of fall armyworm should be controlled before onset of tassel stage and use of this LARVIN® recommendation.</p>	
	Cutworms	See "SPECIAL DIRECTIONS ON CUTWORMS"			




11 9 16

RECOMMENDED USES FOR CUTWORMS

SPECIAL DIRECTIONS ON CUTWORMS

SPRAY VOLUME FOR CUTWORMS: Apply in a minimum finished spray volume of 3 gallons per acre by air or 15 gallons per acre by ground.


SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVINE BRAND DF WSP		
COTTON SOYBEANS SWEET CORN (Florida, Fresh Market Only) CABBAGE BROCCOLI CAULIFLOWER LEAFY VEGETABLE	Cutworms	0.5 to 0.75	0.63 to 0.94	Postemergence Rescue Treatments: For best results, apply as a broadcast spray when cutworms are actively feeding. If cutworms are feeding below the ground, control will be more difficult. Light cultivation with a rotary hoe before spraying may enhance control. If banding, use a minimum width of 10 inches over the row. When banding, determine the amount of chemical to use per acre, by dividing the band width by the row width and multiplying by the appropriate broadcast rate.	Refer to each specific crop for use restrictions.



RECOMMENDED USES FOR ORNAMENTALS AND TREES


For dilute-spray ground applications to ornamentals (such as: Trees, Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants and Non-bearing fruits): Apply the specified dosage per 100 gallons of water. For concentrate-spray ground applications, apply the specified dosage per acre in sufficient spray volume to provide thorough coverage. For aerial application to ornamental plants, shrubs and trees (except forests), apply the specified dosage per acre in a minimum total spray volume of 5 gallons to provide thorough coverage. BEFORE MAKING TREATMENTS REFER TO "PLANT TOLERANCE INFORMATION" LISTED ON THIS LABEL.

SITE	PEST	POUNDS OF LARVING BRAND DF WSP PER 100 GALS SPRAY/ OR PER ACRE	SPECIFIC DIRECTIONS	USE RESTRICTIONS
<p>ORNAMENTALS such as: Trees (except forests), Shrubs, Herbaceous bedding plants, Herbaceous greenhouse plants, Non-bearing fruit trees (Citrus: Grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats and hybrids;</p> <p>Nut Crops: almonds, filberts, pecans, walnuts;</p> <p>Pome Fruits: apples, pears, loquats, crabapple, oriental pears, quince;</p> <p>Stone Fruits: peaches, apricots, nectarines, plums, prunes and cherries)</p>	<p>Armyworms Bagworm Cankeworms (such as: spring, fall) Gypsy moth Leafrollers (such as: omnivorous, fruittree) Spruce budworms Webworms (such as: fall, mimosa) Tent caterpillars</p>	<p>0.63 to 0.94</p>	<p>See "PLANT TOLERANCE INFORMATION" and "APPLICATION" sections of this label.</p> <p>Apply when pests appear and thereafter as needed not to exceed 6 applications per site per season.</p>	<p>Do not use on fruit trees or vines that will bear edible fruit within one year.</p>



RECOMMENDED USES FOR NON-CROP AREAS

SITE	PEST	DOSAGE PER ACRE		SPECIFIC DIRECTIONS	USE RESTRICTIONS
		Pounds Active	Pounds LARVING BRAND DF WSP		
NON-CROP AREAS (Wastelands, Roadsides, Rights-of-Way, Grassways, Shelter belts, and Fencerows)	Armyworms	0.40 to 0.76	0.50 to 0.95	Treat non-crop areas where pests congregate to reduce infestation levels or to prevent movement into croplands. Apply when pests appear. Make additional applications as needed, not to exceed 4.56 lbs. of active ingredient per site per season.	Do not allow livestock to graze treated areas. Do not use treated plants as food or feed.
	Bagworm				
	Budworms				
	Cankerworms (such as: spring, fall)				
	Cutworms				
	Gypsy moth				
	Heliothis spp. (such as: corn earworm)				
	Leafrollers (such as: omnivorous, fruittree)				
	Loopers				
	Oakworms				
	Sawflies				
	Skippers				
	Tent caterpillars				
Webworms (such as: fall, mimosa)					




ORNAMENTAL PLANT TOLERANCE INFORMATION

LARVIN® brand Thiodicarb insecticide has been tested on many species of ornamental plants. Results indicate plants listed below are tolerant to the higher rates of this product. Horticultural varieties of many plants differ significantly in sensitivity to pesticides. Before making applications to plants not listed, it is recommended that a small portion of the crop be treated to determine its tolerance to this product. Plants stressed from heat, moisture or other factors are usually more sensitive to sprays. Phytotoxic reactions can be minimized by spraying when stress is minimal such as early mornings or late evenings.

ORNAMENTAL PLANTS TOLERANT TO THIODICARB INSECTICIDE

TREES	SHRUBS	HERBACEOUS GREENHOUSE PLANTS		HERBACEOUS REDDING PLANTS
Apple, (Crab) Birch, (Weeping, White) Honeylocust, (Sunburst) Maple, (Sugar) Pine, (Scotch) Spruce, (Norway)	Arborvitae Aucuba Azalea Boxwood Camellia Elaeagnus Euonymus Fatsia, (Japanese) Gardenia Hawthorn Holly, (Japanese) Holly, (Yaupon) Juniper Lantana Ligustrum Lilac Mahonia Osmanthus Pittosporum Pyracantha Rhododendron Rose, (Hybrid Tea) Yew	<i>Acalypha hispida</i> <i>Acalypha wilkesiana</i> <i>Aechmea lueddemanniana</i> <i>Alternanthera versicolor</i> Aluminum Plant <i>Begonia semperflorens</i> Cactus <i>Ceropegia woodii</i> <i>Chamaedorea elegans</i> <i>Chlorophytum comosum</i> Dieffenbachia <i>Dracaena sanderana</i> Echeveria Fern <i>Ficus diversifolia</i> Gloxinia <i>Gynura sarmetosa</i> <i>Hechtia tellandsioides</i> Hoya <i>Iboza riparia</i> Ivy, (Swedish)	Ixora Jade Plant <i>Kalanchoe daigremontiana</i> <i>Maranta leuconeura erythroneura</i> <i>Nephtrolepis exaltata</i> <i>Peperomia magnoliifolia</i> <i>Peperomia obtusifolia</i> <i>Philodendron oxycardium</i> Piggyback Plant <i>Pilea grandis</i> Poinsettia Pseuderanthemum <i>Rhoeo spathacea</i> <i>Sansevieria trifascia</i> Senecio <i>Spathyiphyllum wallisii</i> <i>Syngonium podophyllum</i> Torenia <i>Zebrina pendula</i> <i>Zygocactus truncata</i>	Calendula Canna Chrysanthemum Coleus Geranium Hosta Impatiens Lantana Periwinkle Petunia Salvia Snapdragon Verbena Violet



Number of Water Soluble Packets to Use

Acres	1.25 lb Packets (1.0 lb active)				2.50 lb Packets (2.0 lbs active)			
	Typical Use Rates in Pounds of Active Ingredient per Acre							
	0.25	0.50	0.75	1.00	0.25	0.50	0.75	1.00
	Packets Needed for Given Acres							
4	1	2	3	4		1		2
8	2	4	6	8	1	2	3	4
12	3	6	9	12		3		6
16	4	8	12	16	2	4	6	8
20	5	10	15	20		5		10
24	6	12	18	24	3	6	9	12
28	7	14	21	28		7		14
32	8	16	24	32	4	8	12	16
36	9	18	27	36		9		18
40	10	20	30	40	5	10	15	20
60	15	30	45	60		15		30
80	20	40	60	80	10	20	30	40
100	25	50	75	100		25		50
120	30	60	90	120	15	30	45	60
140	35	70	105	140		35		70
160	40	80	120	160	20	40	60	80

Estimate use requirements to the closest number of packets without exceeding labeled use rate for crop. Do not split packets.


Suggested methods for determination of number of packets needed:

1. Use table above. For example, to spray a 24 acre field at a rate of 0.75 pounds active would require 18 - 1.25 lb. packets or 9 - 2.5 lb. packets.
2. Calculate number of packets. To calculate number of packets, multiply the number of acres to be sprayed by the desired pounds active to be used per acre then divide by the active ingredient weight of the packet size you are using.

$$\text{Number of packets to be used} = \frac{(\text{acres to be sprayed} \times \text{desired pounds active per acre})}{(\text{packet size in pounds active})}$$

For example, to spray a 22 acre field at a rate of 0.50 pounds active using the 1.25 lb. packets (1.0 lb. active packet) would require $(22 \times 0.50) / 1.0$ or 11 packets. To spray the same field at the same rate using the larger 2.5 lb. packets (2.0 lb. active) would require $(22 \times 0.50) / 2.0$ or 5.5 packets. Because splitting of packets is not recommended, the applicator should use either 5 or 6 - 2.5 lb. packets depending on pest pressure and label limitations for the crop being sprayed. Do not exceed the labeled use rate for the crop being sprayed.

3. Mix packet sizes. For example, to spray a 20 acre field at a rate of 0.75 pounds active using the larger 2.5 lb. packets (2.0 lb. active) would require $(20 \times 0.75) / 2.0$ or 7.5 packets. Because splitting of packets is not recommended, the applicator could use 7 - 2.5 lb. packets plus 1 - 1.25 lb. packet.



16916

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experiences. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO THE BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

LARVIN is a registered trademark of Rhône-Poulenc.
U.S. Patent No.: 4,382,957

LARVIN® brand DF WSP Thiodicarb Insecticide 80% Dry Flowable (PENDING) Submitted 2/16/93.

