

US ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (75-767)  
WASHINGTON, DC 20460

EPA REGISTRATION NO.  
264-541

DATE OF ISSUANCE  
JAN 26 1994

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

TERM OF ISSUANCE  
NAME OF PESTICIDE PRODUCT  
Mocap Gel Nematicide/Insecticide

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Rhone-Poulenc Ag Company  
P. O. Box 12014  
2 T. W. Alexander Dr.  
Research Triangle Park, NC 27709

544 2636 <sup>165</sup>/<sub>38</sub>  
545 2279 <sup>166</sup>/<sub>18</sub>

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the labeling changes listed below before you release the product for shipment:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 264-541.

b. Under the "Personal Protective Equipment (PPE)" section the following changes must be made:

1. Remove the statement, "applicators and other handlers must wear a chemical-resistant protective suit" and replace it with a statement that reads:

"Applicators and other handlers must wear coveralls over long-sleeved shirt and long pants".

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

RAF

JAN 26 1994

MADDEN DSK #4 1/25/94

2. Immediately after the statement that reads "chemical resistant headgear for overhead exposure" add a statement that reads:

"chemical-resistant apron when cleaning equipment, mixing, or loading".

- c. Add the following statements to the end of the "User Safety Recommendations" section:

"Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

- d. Revise the section "State Restrictions" so that it reads:

"Rhone-Poulenc does not recommend the use of Mocap gel on tobacco in Florida."

- e. In accordance with PR Notice 82-1, under the application directions for tank mixing Di-syston 8 Liquid Systemic Insecticide and Mocap Gel add a statement that reads:

"in accordance with the more (most) restrictive of label limitations and precautions."

4. Submit a revised Confidential Statement of Formula (CSF) with the word "example" removed. You must identify the actual inert to be used to manufacture this product. Each trade name should have its specific chemical component along with the CAS number.

5. The Precautionary Review Section of the Registration Support Branch has not completed a review of the acute toxicology data submitted to support the registration of this product. The Precautionary Statements and Statements of Practical Treatment currently on the label are derived from a similar product (EPA Reg. No. 264-458). These statements are subject to change once the review of the acute toxicology data is complete.

6. Submit one copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

399

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

- ) Enclosure: 1) Stamped label  
2) A-79 Enclosure
- )

4 2 9

# RESTRICTED USE PESTICIDE

ACUTE DERMAL TOXICITY

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

## MOCAP GEL Nematicide-Insecticide

### GEL-TEC™ Water Soluble Paks

ACTIVE INGREDIENTS: Ethoprop: (O-Ethyl S, S-Dipropyl Phosphorodithioate)..... 68.2%

INERT INGREDIENTS: .....31.8%

(Contains 6 pounds active ingredient per gallon, each water soluble packet contains 1/3 gallon with 2 pounds active ingredient)

EPA Reg. No. 264-LUR

EPA Est. No. 264-GA-001



**KEEP OUT OF REACH OF CHILDREN  
DANGER POISON  
PELIGRO**



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-334-9745

### STATEMENT OF PRACTICAL TREATMENT

Ethoprop is an organophosphate.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Contains petroleum distillates. Avoid aspiration to lungs.

**IF INHALED:** Remove to fresh air and apply artificial respiration if needed.

**IF ON SKIN:** Remove all contaminated clothing and wash skin with soap and water. Get medical attention.

**IF IN EYES:** Flush with plenty of water for at least 15 minutes. Call a physician immediately.

Persons providing first aid should prevent contamination of themselves.

#### NOTE TO PHYSICIAN

This product is a cholinesterase inhibitor. Severe symptoms and signs include diarrhea, pinpoint and non-reactive pupils, respiratory difficulty, pulmonary edema, cyanosis, loss of sphincter control, convulsions, and coma.

Support respiration as needed. Measures should include removal of secretions, maintenance of a patent airway, and if necessary, artificial ventilation.

If cyanosis is absent, give ATROPINE 2-4 mg intravenously (0.05 mg/kg for children). Repeat atropine at 5-10 minute intervals until atropinization occurs (dry, flushed skin, tachycardia, pupillary dilatation), and maintain for 48 hours.

If cyanotic, give initial atropine intramuscularly and start measures to improve ventilation.

Start 2-PAM (PROTOPAM, Ayerst) at the same time. Give 1-2 grams PROTOPAM (20-40 mg/kg for children) in 100cc saline over 15-30 minutes. If pulmonary edema is present, give intravenously slowly as a 5% solution in water over a period of at least 5 minutes.

A second dose may given after one hour if muscle weakness persists. Additional doses may be given cautiously for persistent muscle weakness. May be given by intramuscular or subcutaneous routes if intravenous administration is not feasible.

In case of skin contact, wash patient with soap and water followed by wash with 95% ethyl alcohol. Keep patient under constant observation for 24-36 hours. Symptoms may persist for one month.

The use of theophylline, ~~barbiturates~~, phenothiazines, reserpine, and succinyl choline is contraindicated.

with COMMENTS  
in EPA Letter Dated

JAN 26 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

**BEST AVAILABLE COPY**

599

# PRECAUTIONARY STATEMENTS

## **DANGER HAZARD TO HUMANS AND DOMESTIC ANIMALS**

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get into eyes, on skin, or on clothing. Do not breathe fumes or spray mist. Do not contaminate feed or food.

**SYMPTOMS OF POISONING:** Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

## **PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

Do not store in or around the home.

## **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear a chemical-resistant protective suit, chemical-resistant gloves such as barrier laminate, butyl rubber (> 14 mils), nitrile rubber (> 14 mils) or viton (> 14 mils), chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure and a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## **User Safety Recommendations**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

## **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic organisms (fish and invertebrates) and extremely toxic to birds. Cover or disc spill areas. Birds in treated areas may be killed. For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be dangerous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment wash waters.

Do not apply within 140 feet of inland freshwater habitats.

Along the Atlantic seaboard, do not apply within 800 feet of brackish water habitats.

Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply in Long Island, New York.

**BEST AVAILABLE COPY**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and expectations pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

The 48 hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

**EXCEPTION:** If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is a chemical-resistant protective suit, chemical-resistant gloves, such as barrier laminate, butyl rubber (> 14 mils), nitrile rubber (> 14 mils), or viton (> 14 mils), chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### STORAGE AND DISPOSAL

#### STORAGE

Do not contaminate water, food, or feed by storage or disposal. Do not store in or around the home.

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Dispose of uncontaminated outer bags and boxes in a sanitary landfill, or by other procedures approved by state and local authorities.

### GENERAL INFORMATION

MOCAP® GEL is formulated as a gelatinous mixture of ethoprop containing the equivalent of 6 pounds of ethoprop per gallon of formulated product. MOCAP® GEL is packaged in 1/3 gallon water soluble paks. Each water soluble pack contains 2 pounds of ethoprop active ingredient.

### CAUTIONS AND RESTRICTIONS

Use MOCAP® GEL Brand Nematicide-Insecticide only in accordance with label directions, warnings, and cautions. **DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING, AS ANY RESIDUES REMAINING MAY BE ILLEGAL OR HARMFUL. DO NOT STORE OR USE IN OR AROUND THE HOME OR HOME GARDEN.**

### APPLICATION STATEMENTS

- Calibrate and adjust application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. For any leftover material, see instruction for "Storage and Disposal" on this label.
- Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

#### STATE RESTRICTIONS

- Do not apply on tobacco in Florida.

### MIXING, LOADING, AND HANDLING INSTRUCTIONS

**GENERAL HANDLING INSTRUCTIONS:** Each MOCAP® GEL water soluble bag comes sealed in a clear plastic outer sleeve. Do not break the seal until the time of use. To use, open the end of the outer sleeve by tearing along the printed solid black line. Do not use a knife or other sharp object to break the seal. Once the outer protective sleeve is open, the water soluble bag inside can be dropped into the spray tank at the appropriate time without touching it.

- MOCAP® GEL spray mixture may affect some plastic materials after prolonged use. Spray tubing hoses and gaskets should be inspected frequently, particularly black-colored materials. Replace any material which is softened or swollen. Polyethylene, polypropylene, nylon, and Teflon materials are recommended. Polyvinyl chloride (PVC) materials are not recommended.

**MOCAP® GEL ALONE:** Begin filling the spray tank with clean water. When bottom of spray tank is covered with water to a depth of 6-8 inches, add the recommended number of MOCAP® GEL soluble bags. Avoid dropping the soluble bags into the tank in such a way that the bags end up covering the recirculation and outlet holes. After adding bags, continue to fill tank with water to achieve total spray volume desired. Allow 3-4 minutes after dropping bags into the tank for bags to dissolve before starting agitation. Maintain sufficient agitation to ensure a uniform spray mixture during application.

**BEST AVAILABLE COPY**

**MOCAP® GEL TANK-MIXTURES:** Begin filling the spray tank with clean water. When bottom of spray tank is covered with water to a depth of 6-8 inches, add the recommended number of MOCAP® GEL soluble bags. After adding bags, continue to fill tank with water to achieve total spray volume desired. Allow 3-4 minutes after dropping bags into the tank for bags to dissolve before starting agitation. Add the recommended amount of wettable powder, soluble powder, flowable, EC or soluble liquid product to be tank mixed to the spray tank. Maintain sufficient agitation to ensure a uniform spray mixture during application.

• If MOCAP® GEL is to be tank mixed with liquid fertilizers, physical compatibility should be tested prior to mixing. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing.

**HANDLING PRECAUTIONS**

- Do not allow the empty outer sleeve or the tear-strip to fall into the spray tank as they are not water soluble and may clog the sprayer.
- Do not handle the inner bag with wet gloves as the bag will begin to dissolve on contact with even a small amount of water.
- Do not open the inner bags or attempt to remove the gel from them. The gel is not designed to be re-measured or subdivided.
- The PVA bag should not be opened and partial bag not used for a tank mix.
- The entire contents of the water soluble bag must be used.

**INCORPORATION STATEMENT**

MOCAP® GEL can be incorporated with a rotary tiller, rotary hoe, spring-tooth harrow, by double-discing or with other equipment for mixing the chemical with the soil to the recommended depth.

**GROUND APPLICATION**

Use sufficient total spray volume to thoroughly and evenly cover the area to be treated.

**REQUIREMENTS FOR IRRIGATION SYSTEMS**

Apply this product only through sprinkler systems including: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Wear personal protective equipment as defined in the PPE section for applicators and other handlers when making adjustments or repairs on the chemigation system when MOCAP® EC is in the irrigation water.

**POSTING OF AREAS TO BE CHEMIGATED IS REQUIRED WHEN:**

1. Any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities.
2. When the chemigated area is open to the public, posting must conform to the following requirements: Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area toward the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

\*This sign is in addition to any sign posted to comply with Worker Protection Standard.

**EQUIPMENT AND OPERATIONS REQUIREMENTS:**

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.



Do not apply when wind speed favors drift beyond the area intended for treatment.

The chemical injection tank should be cleaned before application to remove all fertilizer, pesticide and other foreign matter. Add MOCAP® GEL paks as described in the instructions for mixing, loading, and handling. The MOCAP® GEL water mixture in the injection tank should be under constant agitation. Carefully calibrate the system before and during application. The irrigation system should provide uniform water flow and should have no leaks.

For sprinkler irrigation systems, the MOCAP® GEL water mixture should be injected continuously during water application in 7,000 to 28,000 gallons of water per acre.

### SPECIFIC USE DIRECTIONS

Where a range of rates is recommended, use the higher rate if pest infestations are expected to be severe. To determine your infestation level, consult local or state recommendations for sampling methods and economic thresholds. Where both banded and broadcast applications are allowed, refer to the specific crop directions for the proper use of each application method. Use sufficient total spray volume for thorough and uniform application.

## TOBACCO

APPLICATION TIMING	PESTS CONTROLLED	RATE OF MOCAP® GEL		APPLICATION DIRECTIONS
		BANDED WATER SOLUBLE PAKS PER 4000 ROW FEET	BROADCAST WATER SOLUBLE PAKS PER ACRE	
from one week before transplanting to transplanting time	southern root-knot nematode wireworms cutworms flea beetle larvae mole crickets	1 to 2	3 to 6	Band: Apply in a band 18 to 24 inches wide on the row after the fertilizer has been listed or bedded. Broadcast: Use a double gang disc harrow or other cultivating equipment that will do a good job of mixing MOCAP® GEL with the soil. After mixing, apply fertilizer and, for best results, shape a wide bed with a tractor equipped with 4 disc hillers and wide sweeps for running out the middles. Be sure that only the treated soil (top 2 to 4 inches) is used to make beds.
	flea beetles	1 to 2	3 to 6	Add 1/2 gallon (4 pints) per acre of Di-Syston® 8 Liquid Systemic Insecticide to the spray tank containing MOCAP® GEL mixture for nematode and wireworm control (see above). Do not make more than one field application per crop per season regardless of the formulation of Di-Syston used.
from two weeks before transplanting to transplanting time	cutworms flea beetle larvae wireworms	1 PAK PER 12,000 ROW FEET	1	Broadcast: Broadcast material uniformly on top of the soil and incorporate to a depth of 2 to 4 inches. Use a disc harrow or other cultivation equipment that will do a good job of mixing MOCAP® GEL with the soil. Band: Apply to pre-formed bed at bed reshaping or reforming at or just before transplanting in a band 18 to 24 inches wide. Make this application to the final bed surface and incorporate into the top 2 to 6 inches of soil into which the crop will be transplanted.

Note: Use the higher rates in fields where nematode infestations have been especially severe or where soil is sandy. To determine your infestation level, consult local or state recommendations for sampling methods and economic thresholds. Mix with the top 2 to 4 inches of soil immediately after application.

Do not use MOCAP® on tobacco in Florida.

**BEST AVAILABLE COPY**

## LIMITED WARRANTY AND DISCLAIMER

999

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

### NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside the United States.

Rhône-Poulenc Ag Company  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709

MOCAP is a registered trademark of Rhône-Poulenc.

Di-Syston is a registered trademark of Miles Corporation.

MCCAP® GEL Nematicide-Insecticide (PENDING) Registered 10/25/93.

