



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

264-542

Date of Issuance:

Term of Issuance:

Unconditional

Name of Pesticide Product:

Chipco Aliette WSP
Brand Fungicide

NOTICE OF PESTICIDE:

x ☐ Registration
☐ Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Rhone-Poulenc AG Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5) provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Add the phrase "EPA Registration No. 264-542" to your label.

b. You have an Asterisk "*" after the brand names "Fore*" and "Daconil 2767*" but no corresponding asterisk with brand names on the last page of the label. Also, the asterisk does not appear every time the brand name is listed on the label.

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

A stamped copy of the label is enclosed for your records.

BEST AVAILABLE COPY

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Signature of Approving Official:

151

Date:

CHIPCO^(K) ALIETTE^(K) WSP brand Fungicide

ACTIVE INGREDIENT: Aluminum tris (O-ethyl phosphonate).....80.0%

INERT INGREDIENTS:20.0%

E. P. A. REG. NO. 264-XXX

E. P. A. EST. NO.

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-800-334-9745.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye injury. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Do not reuse outer sleeve. Dispose of outer sleeve in the trash

GENERAL HANDLING INSTRUCTIONS: Each CHIPCO® ALIETTE® WSP Fungicide water soluble packet comes sealed in a clear plastic outer sleeve. Do not break the seal until the time of use. To use, open the end of the outer sleeve by tearing along the printed solid black line. Do not use a knife or other sharp object to break the seal. Once the outer protective sleeve is open, the water soluble packet inside can be dropped into the spray tank at the appropriate time without touching it.

HANDLING PRECAUTIONS

- Do not allow the empty outer sleeve or the tear-strip to fall into the spray tank as they are not water soluble and may clog the sprayer.
- Do not handle the inner packet with wet gloves as the packets will begin to dissolve on contact with even a small amount of water.
- Do not open the inner packets or attempt to remove the contents from them. The water soluble packet is not designed to be remeasured or subdivided.
- The PVA packets should not be opened and partial packets not used for a tankmix.
- The entire contents of the water soluble packet must be used.

DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including microjet, drip, solid set and center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of CHIPCO® ALIETTE® WSP in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of CHIPCO® ALIETTE® WSP, and then the remaining volume of water. Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of CHIPCO® ALIETTE® WSP into the irrigation water line so as to deliver the desired rate per acre. The suspension of CHIPCO® ALIETTE® WSP should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: For Microjet and Drip Irrigation Systems: When treatment with CHIPCO® ALIETTE® WSP has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours. **For Solid Set and Center Pivot Irrigation Systems:** When treatment with CHIPCO® ALIETTE® WSP has been completed, further field irrigation over the treated area should be avoided until foliage is dry to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

ORNAMENTALS AND BEDDING PLANTS

Use CHIPCO® ALIETTE® WSP Fungicide on ornamentals and bedding plants grown in field nursery, greenhouse, landscaping and conifer nursery situations, for control of diseases caused by Pythium and Phytophthora. CHIPCO® ALIETTE® WSP can also be used on ornamentals for the control of downy mildew and fire blight and for the suppression of bacterial blight caused by certain pathovars of *Xanthomonas campestris*. Applications should be made prior to disease development and should be made in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed recommended rates or apply more frequently than at specified intervals or plant injury may occur.

BEST AVAILABLE COPY

ORNAMENTALS

FOLIAR APPLICATIONS to plants such as Aglaonema, Aphelandra, Azalea, Bougainvillea, Boxwood, *Cattelya skinneri*, Cissus, Dieffenbachia, Hibiscus, Juniper, Leather-leaf Fern, Pittosporum, Philodendron, Pothos, Rhododendron, Spathiphyllum and Taxus media.

Mix 2 to 5 pounds of product (2 to 5 WS packets) with 100 gallons of water and spray to wet using no more than 400 gallons of spray per acre. Repeat as necessary but do not exceed one application every 30 days.

DRENCH APPLICATIONS to plants such as Aphelandra, Azalea, Boxwood, Cissus, Dieffenbachia, Japanese Holly, Juniper, Monterey Pine, Philodendron, Pieris, Pittosporum, Rhododendron, Schefflera, Spathiphyllum, and Taxus media:

Mix 1 to 2 pounds of product (1 to 2 WS packets) with 200 gallons of water and apply to 800 sq. ft. (equivalent to 2 pt. solution per sq. ft.) Repeat as necessary but do not exceed one application every 30 days.

SOIL INCORPORATION to plants such as Rhododendron, Azalea, or Pieris for control of Phytophthora species only:

Mix 1 to 2 pounds of product (1 to 2 WS packets) per cubic yard of soil mix immediately before potting. If conditions are favorable for disease development, applications as a foliar spray or drench should be made as necessary but do not exceed one application every 30 days. Use soil incorporation only with well-rooted plants.

BEDDING PLANTS

FOLIAR APPLICATIONS to plants such as Begonia, Pansy, Vinca, Marigold, Zinnia, Petunia, Geranium, and Impatiens:

Mix 2 to 4 pounds of product (2 to 4 WS packets) with 100 gallons of water and spray to wet using no more than 400 gallons of water per acre. Repeat as necessary but do not exceed one application every 30 days.

FOR USE ON CONIFERS IN NURSERIES TO PREVENT PHYTOPHTHORA ROOT ROT

DIP TREATMENTS to conifers such as Douglas firs, spruce and pines. Dip transplants in 3 pounds of product (3 WS packets) per 100 gallons to thoroughly wet plant and root mass. Dip immediately before transplanting. When making dip applications, wear water proof gloves, protective eyewear, coveralls, shoes and socks.

FOLIAR APPLICATIONS to conifers such as Douglas firs, spruce and pines. Mix 3 to 5.0 pounds of product (3 to 5 WS packets) with 100 gallons of water and spray to wet using no more than 400 gallons of water per acre. Do not exceed one application every 30 days.

DOWNY MILDEW CONTROL IN ROSES

FOLIAR APPLICATIONS TO ROSES: field, container, landscape and mini varieties to control downy mildew (*Peronospora sparsa*). Applications should be made in conjunction with a disease sanitation program to reduce the spread of the disease to uninfected plants.

Mix 3 pounds of product (3 WS packets) with 100 gallons of water and spray to wet using no more than 400 gallons per acre. Repeat as necessary but do not exceed one application every 7 days.

FIRE BLIGHT SUPPRESSION

FOLIAR APPLICATIONS to plants such as ornamental pear, pyracantha and hawthorne. Applications should be made in conjunction with a strict sanitation program to reduce the spread of the disease to uninfected plants.

Mix 3 pounds of the product (3 WS packets) with 100 gallons of water and spray to wet using not more than 400 gallons of water per acre. CHIPCO ALIETTE works solely as a preventive treatment. Begin spray treatments at pre-bloom stage and continue at 7 day intervals until bloom period ends. Do not exceed one application every 7 days.

BACTERIAL BLIGHT SUPPRESSION

FOLIAR APPLICATIONS to plants such as English ivy, schefflera, anthurium, dieffenbachia, spathiphyllum, syngonium, and ficus for the suppression of the *Xanthomonas campestris* pathovars *hederae*, *dieffenbachiae*, *syngonii*, and *fici*. Applications should be made in conjunction with a disease sanitation program to reduce the spread of the disease to uninfected plants.

Mix 2 to 4 lbs of product (2 to 4 WS packets) with 100 gallons of water and spray to wet using no more than 400 gallons of spray per acre. Repeat as necessary but do not exceed one application on every 7 days.

Refer to compatibility statements concerning use of coppers or other compounds.

PLANT TOLERANCE

Plant tolerances to CHIPCO[®] ALIETTE[®] WSP Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to CHIPCO[®] ALIETTE[®] WSP Fungicide. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

BEST AVAILABLE COPY

ORNAMENTAL AND BEDDING PLANT TANK MIXTURES

Additional Disease Control: If ornamentals are threatened by additional diseases, CHIPCO® ALIETTE® WSP is compatible with most commonly used fungicides. For control of diseases such as Aerial Web Blight, Alternaria Leaf Blight, Botrytis Blight, Fusarium Leaf Spot, Helminthosporium Leaf Spot, and Ray Blight, CHIPCO® ALIETTE® WSP can be tank-mixed with CHIPCO® 26019 brand fungicide. Consult the CHIPCO® 26019 label for a complete listing of diseases controlled and for application rates.

If a tank-mix with other fungicides is used, follow label directions for the use of that product and apply at the recommended rate for control of the target disease organism. CHIPCO® ALIETTE® WSP is not compatible with the flowable formulation of Daconil 2787. These products should not be combined for application.

COMPATIBILITY

CHIPCO® ALIETTE® WSP may not be compatible with foliar fertilizers or with copper based fungicides. To avoid undesirable copper phytotoxicity, do not apply CHIPCO® ALIETTE® WSP and copper based compounds at less than 7 day intervals. Do not mix CHIPCO® ALIETTE® WSP with any sticker, extender or wetting agent.

To determine the physical compatibility of CHIPCO® ALIETTE® WSP with any other product, use a small container to mix a small amount (e.g. 1 pint) of spray solution, containing all ingredients in the same order and ratio as the anticipated use. If any indication of physical incompatibility develop, do not use this mixture for spraying. Indication of incompatibility usually appear within 5-15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

CHIPCO® ALIETTE® WSP is not physically compatible with flowable formulations of Daconil 2787 or Fore fungicides as a tank mixture. These products should not be combined for application.

TURF

CHIPCO® ALIETTE® WSP is a systemic fungicide which may be used in a seasonal program for the control of Pythium diseases, such as blight and root rot, and Yellow tuft on common turfgrasses on golf courses, sod farms and other turf areas. Apply as a foliar spray, using 1 to 5 gallons of water per 1,000 sq. ft., as indicated in the table below. Apply with a properly calibrated sprayer.

DISEASE	INTERVAL OF APPLICATIONS	RATE OZ. PRODUCT/1,000 FT ²	NUMBER OF WS PACKETS PER ACRE
Pythium diseases, Yellow Tuft	14 days	4.0	11
	21 days	8.0	22

Begin preventive applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations.

TURF TANK MIXTURES

Summer Stress Complex: For control of summer stress complex caused by a complex of *Pythium* and *Rhizoctonia* diseases, CHIPCO® ALIETTE® WSP can be tank-mixed with Fore WP Fungicide on common turfgrasses on golf courses, sod farms, industrial or municipal turf areas, and professional applications to residential lawns.

PRODUCT	DISEASE	APPLICATION INTERVAL	RATE OZ. PRODUCT PER 1,000 FT ²	NUMBER OF WS PACKETS PER ACRE
CHIPCO® ALIETTE® WSP + Fore WP	Summer Stress Complex (<i>Pythium</i> and <i>Rhizoctonia</i> spp.)	14 days	4.0 to 8.0 + 4.0 to 8.0	11 to 22

Apply as a foliar spray using 1 to 5 gallons of water per 1,000 sq.ft. Apply with a properly calibrated sprayer. Begin preventative applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations. Do not mix CHIPCO® ALIETTE® WSP with the flowable formulation of Fore Fungicide as physical incompatibility of these products may result.

Additional Disease Control: If turf is threatened by additional diseases, CHIPCO® ALIETTE® WSP is compatible with most commonly used fungicides. For control of diseases such as Dollar Spot, Brown Patch, Helminthosporium Leaf Spot, and Necrotic Ring Spot, CHIPCO® ALIETTE® WSP can be tank-mixed with CHIPCO® 26019 or CHIPCO® 26019 FLO brand fungicides. Consult the CHIPCO® 26019 and CHIPCO® 26019 FLO labels for a complete listing of diseases controlled and for application rates.

If a tank-mix with other fungicides is used, follow label directions for the use of that product and apply at the recommended rate for control of the target disease organism. CHIPCO® ALIETTE® WSP is not compatible with the flowable formulations of Daconil 2787 or Fore Fungicide. These products should not be combined for application.

BEST AVAILABLE COPY

19

2

CHIPCO® ALIETTE® WSP is not physically compatible with flowable formulations of Daconil 2787 or Fore fungicides as a tank mixture. These products should not be combined for application.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

ACCEPTED
MAY 10 1968
LESLIE L. BROWN

264-542

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

ALLETTE and CHIPCO are registered trademarks of Rhône-Poulenc.
Daconil 2787 is a registered trademark of Fermenta Corporation.
Fore is a registered trademark of Rohm and Haas Company.

CHIPCO® ALIETTE® WSP Fungicide (PENDING) Submitted 6/3/94.

BEST AVAILABLE COPY