

PM 23

264-600

9-21-99

1 of 12



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

264-600

Date of Issuance:

SEP 21 1999

Date of Expiration:

NOV 1 2001

NOTICE OF PESTICIDE:
 X Registration
 — Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Balance 4SC Herbicide

Name and Address of Registrant (include ZIP Code):

Rhone-Poulenc Ag Company
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 264-600".
 - b. The statements "Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse" may be deleted from the Precautionary Statements, since they also appear in the Safety User Recommendations box.
 - c. Revise the unit of measure in the first statement in the section for Restrictions and Precautions for Use on Field Corn to specify fluid ounces, so that the statement reads "Do not apply more than 4.5 fluid ounces of Balance Herbicide per acre in one season."

Signature of Approving Official:

Date:

SEP 21 1999

EPA Form 8570-6 DK: 305-7546: HB/PM23

7505C
D. KENNY
9/21/99

3. Submit all data required as a condition of registration for use of isoxaflutole on field corn, as stated in the Notice of Pesticide Registration for Technical Isoxaflutole (EPA Registration No. 264-566).

4. This registration is time-limited and will terminate on November 1, 2001 unless renewed by EPA. This registration will permit sale and use only in the states of Arkansas, Colorado, Indiana, Illinois, Iowa, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, Tennessee, Texas (those counties to the north of or bisected by I-20), and Wyoming.

5. By September 1, 2000 or four months after receipt of data, whichever is later, EPA will make a determination as to whether the available information supports expansion of the geographical areas in which this product may be used, or warrants the addition or removal of label restrictions or mitigation measures.

6. During the term of this conditional registration, Rhone-Poulenc will not support efforts, in states where use has not been authorized, to obtain emergency exemptions for use of this product under FIFRA Sec. 18 or additional uses to meet special local needs under FIFRA Sec. 24(c). The scope of this provision will be revised to conform to any decisions reached under provision #5 as to the states where use is authorized.

7. Product registrations of other registrants which include use of isoxaflutole will be required to meet the same label restrictions and limitations on support of FIFRA Sec. 18 and FIFRA Sec. 24(c) requests as EPA has imposed under this Notice of Pesticide Registration or as subsequently modified.

9. Prior to November 1, 2001, EPA will make a determination whether to renew or deny this registration based on the available information. All information EPA is to consider in support of renewal of this registration must be submitted by August, 2001. Data submitted in support of this registration after that date will not be considered "available information" for purposes of a decision on renewal of this registration. If EPA does not make a determination by November 1, 2001, either to extend or deny this registration, EPA's inaction will constitute a denial of application for registration.

10. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

3 of 12

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

4/12

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible non-target plants.

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.

BALANCE® 4SC Herbicide

For weed control in field corn in the states of: Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, Tennessee, Texas (those counties north of or bisected by I-20), and Wyoming. In the states of Colorado, Kansas and Missouri, there may be more restrictive conditions on the use of this product. You should check with your state regulatory authority prior to use.

ACTIVE INGREDIENT:

Isoxaflutole* [5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethylbenzoyl) isoxazole]..... 40.5%

INERT INGREDIENTS: 59.5%

*Product contains 4.0 pounds of isoxaflutole per gallon.

E.P.A. Reg. No. 264-XXX

E.P.A. Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-888-4RP-AGRO (1-888-477-2476)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting. Get medical attention.

IF ON SKIN: Wash skin with plenty of soap and water. Get medical attention.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NOTE TO PHYSICIAN: No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eye wear. When mixing/loading or cleaning equipment, wear a chemical resistant apron in addition to the other required PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ACCEPTED
with COMMENTS
In EPA Letter Dated
SEP 21 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
264-600

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water, is coveralls over long-sleeved shirt and long pants, waterproof gloves, socks plus chemical resistant footwear and protective eye wear.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry secured storage area.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

BALANCE® 4SC Herbicide is formulated as a soluble concentrate of isoxaflutole at a concentration of 4 pounds of active ingredient, isoxaflutole, per gallon.

BALANCE® 4SC is a selective herbicide for control of important broadleaf and grass weeds infesting field corn when used as a preplant (surface-applied or incorporated) or preemergence herbicide.

BALANCE® 4SC is effective in controlling triazine or ALS resistant populations of weed species which are listed in the "Weed Species Control" tables below on this label.

Seed corn inbreds vary in their response to BALANCE® 4SC Herbicide. Consult your seed company for advice before using BALANCE® on seed corn inbreds.

Environmental extremes may increase the potential for crop response.

Do not apply this product through any type of irrigation system.

Do not apply this product using aerial application equipment.

Do not use flood irrigation to apply or incorporate this product.

For use in the states of: Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, Tennessee, Texas (those counties north of or bisected by I-20), and Wyoming. In the states of Colorado, Kansas and Missouri, there may be more restrictive conditions on the use of this product. You should check with your state regulatory authority prior to use.

MIXING INSTRUCTIONS

BALANCE® 4SC Alone: BALANCE® 4SC can be mixed and applied with water or liquid fertilizer. Fill the spray tank ¼ to ½ of the required volume of water or liquid fertilizer prior to the addition of BALANCE® 4SC. Add the proper amount of BALANCE® 4SC, then add the rest of the water or liquid fertilizer to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

Tank Mixtures: BALANCE® 4SC can be applied in tank mixture with many other pesticides. Refer to the specific crop section for rate recommendations and other restrictions. To apply BALANCE® 4SC in mixture with another product, fill the spray tank ½ to ¾ full with water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add the recommended amount of BALANCE® 4SC and add water to the spray tank to the desired level. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water. Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

Sprayer Cleanup: To avoid injury or exposure to non-target crops, thoroughly clean all mixing and spray equipment on approved rinse pad or on the field site where an approved crop is to be grown. Do this immediately after application and prior to use on crops other than those registered for use on this label.

TANK MIXTURES

BALANCE® 4SC Herbicide can be applied in tank mixture with many other pesticides registered for use on approved crops. Refer to specific crop section for rate recommendations and other restrictions.

COMPATIBILITY

If BALANCE® 4SC Herbicide is to be tank mixed with liquid fertilizers or other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read and follow the label of each tank-mix product used for precautionary statements, directions for use, geographic and other restrictions.

APPLICATION PROCEDURES

APPLICATION TIMING

BALANCE® 4SC may be used in either conventional, conservation tillage, or no-till crop management systems and may be applied either preplant, preplant incorporated or preemergence for use in field corn production. Do not apply after corn emerges or crop injury may occur.

Preplant Surface-Applied: BALANCE® 4SC alone and certain BALANCE® 4SC tank mixtures may be applied up to 14 days before planting field corn. BALANCE® 4SC may be applied up to 30 days prior to planting when used on Roundup Ready® or Liberty Link™ corn where a sequential application of glyphosate or Liberty™ is planned. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be reduced.

Preplant Incorporated: BALANCE® 4SC alone and certain BALANCE® 4SC tank mixtures may be applied up to 14 days before planting field corn. BALANCE® 4SC may be applied up to 30 days prior to planting when used on Roundup Ready® or Liberty Link™ corn where a sequential application of glyphosate or Liberty™ is planned. Apply to the soil and incorporate in the top two inches of soil before planting using a finishing disc harrow, field cultivator with rolling baskets or similar implement capable of providing uniform two inch incorporation.

BALANCE® 4SC treatments are most effective in controlling weeds when adequate rainfall is received within 14 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain occurs, use shallow tillage such as rotary hoe to lightly incorporate BALANCE® 4SC and make certain corn seeds are below the tilled area. If treated soil is moved during tillage practices in such a way that the herbicide barrier is no longer intact, weeds may emerge in those areas from which treated soil has been moved.

Preemergence: Apply BALANCE® 4SC during planting (behind the planter after furrow closure) or after planting, but before weeds or crop emerge. Failure to thoroughly close and firm the seed furrow may allow herbicide to directly contact the seed which can cause injury.

Preplant/Preemergence Burndown: When weeds are present at the time of treatment, a tank mixture of BALANCE® 4SC Herbicide with crop oil concentrate or methylated seed oil is recommended for burndown of labeled weeds less than 3" in height. When weeds are greater than 3" in height or weeds not controlled by BALANCE® 4SC Herbicide are present, the addition of a burndown herbicide (e.g., Gramoxone® Extra, Roundup Ultra™, or WEEDONE® 638) is recommended. If Giant Ragweed or Pennsylvania Smartweed are present at the time of application, the addition of 1 to 1 ½ pounds of atrazine will improve control. Observe directions for use and precautions and restrictions on the label of the burndown herbicide.

GROUND APPLICATION

Apply BALANCE® 4SC alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre. Do not apply with hollow cone nozzles or with other application equipment which does not provide uniform coverage. Use sprayers that provide accurate and uniform application.

Do not exceed the maximum labeled rate for any soil type. Spray overlaps produce areas of over application and increase the potential for crop response.

With the exception of minimum or no-tillage systems, plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL AND FURROW FIRMED.**

In minimum or no-tillage systems, **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL AND SOIL FIRMED.** The use of no-till planters under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if BALANCE® 4SC contacts the germinating corn seed. Check equipment to ensure good seed coverage.

SPECIFIC USE DIRECTIONS

BALANCE® 4SC HERBICIDE APPLIED ALONE

APPLICATION TIMING	AMOUNT OF BALANCE® 4SC PER ACRE			
	Medium and Fine Soils including Loam, Silt loam, Silt, Silty clay loam, Silty clay, Clay, Clay loam, Sandy clay loams and Sandy clays		Coarse Soils including Sands, Sandy loams and Loamy sands	
	Conventional tillage	No-till	Conventional tillage	No-till
PREPLANT (SURFACE-APPLIED or INCORPORATED) 8 to 14** days prior to planting	3.75 to 4.5 fluid ounces	4.5 fluid ounces	2.25 to 3.0 fluid ounces*	2.25 to 3.0 fluid ounces*
PREPLANT (SURFACE-APPLIED or INCORPORATED) 0 to 7 days prior to planting	3.0 to 3.75 fluid ounces	3.0 to 3.75 fluid ounces	1.875 fluid ounces*	1.875 fluid ounces*
PREEMERGENCE	3.0 to 3.75 fluid ounces	3.0 to 3.75 fluid ounces	1.875 fluid ounces*	1.875 fluid ounces*

* If grass pressure is moderate to severe, tank mixing with a pre-emergence grass herbicide is recommended.

** BALANCE® 4SC may be applied up to 30 days prior to planting when used on Roundup Ready® or Liberty Link™ corn where a sequential application of glyphosate or Liberty™ is planned.

RESTRICTIONS AND PRECAUTIONS FOR USE ON FIELD CORN

Do not apply more than 4.5 ounces of BALANCE® Herbicide per acre in one season.


Do not rotate to other crops within 6 months after application.

In the States of CO, KS, KY, MO, and TN, if the water table (i.e, level of saturation) is less than 25 feet below the ground surface, do not use on loamy sand or sand surface soil and subsoils with an average organic matter (in the upper 12 inches) of less than 2% by weight.

In the States of IA, IL, IN, MT, ND, NE, OH, SD, and WY, if the water table (i.e, level of saturation) is less than 25 feet below the ground surface, do not use on sandy loam, loamy sand or sand surface soils and subsoils with an average organic matter (in the upper 12 inches) of less than 2% by weight.

TANK MIX COMBINATIONS

Tank mix combinations may be used in either conventional, conservation tillage, or no-till cropping systems and be applied at the same timings as BALANCE® 4SC Herbicide unless otherwise specified in the tank mix label. Multiple tank mixtures are allowed unless otherwise specified by the respective product labels. Check all tank mix product labels for proper rates and compatibilities for multiple tank mixes.



TANK MIX USE DIRECTIONS

APPLICATION TIMING	AMOUNT OF BALANCE® 4SC PER ACRE			
	Medium and Fine Soils including Loam, Silt loam, Silt, Silty clay loam, Silty clay, Clay, Clay loam, Sandy clay loams and Sandy clays		Coarse Soils including Sands, Sandy loams and Loamy sands	
	Conventional tillage	No-till	Conventional tillage	No-till
PREPLANT (SURFACE-APPLIED or INCORPORATED) 8 to 14 days prior to planting	3.75 to 4.0 fluid ounces	4.5 fluid ounces	2.25 to 3.0 fluid ounces	2.25 to 3.0 fluid ounces
PREPLANT (SURFACE-APPLIED or INCORPORATED) 0 to 7 days prior to planting	2.25 to 3.0 fluid ounces	2.25 to 3.0 fluid ounces	1.5 to 1.875 fluid ounces	1.5 to 1.875 fluid ounces
PREEMERGENCE	2.25 to 3.0 fluid ounces	2.25 to 3.0 fluid ounces	1.5 to 1.875 fluid ounces	1.5 to 1.875 fluid ounces

THIS PRODUCT MAY BE TANK-MIXED WITH THESE HERBICIDES FOR CONTROL OF CERTAIN BROADLEAF AND GRASS WEEDS IN CORN:

TANK MIX PARTNERS NOT CONTAINING ATRAZINE

Apply the following tank mix partners at one half to full rate based on the product's allowable use rate for specific soil types and/or organic matter content. Tank mixes with BALANCE® 4SC Herbicide are not limited to the tank mix partners mentioned in this table. Refer and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

- AXIOM® (½ to ¾ of labeled rate)
- DUAL®/DUAL® III/ DUAL® II MAGNUM
- FRONTIER®
- HARNESS®
- LASSO® / MICRO-TECH® / PARTNER®
- PROWL® (Preemergence application only)
- SURPASS®
- TOPNOTCH®
- SIMAZINE

TANK MIX PARTNERS CONTAINING ATRAZINE

Apply the following tank mix partners at one half to full rate but not to exceed 1.5 pounds of atrazine active ingredient per acre based on the product's allowable use rate for specific soil types and/or organic matter content. Tank mixes with BALANCE® 4SC Herbicide are not limited to the tank mix partners mentioned in this table. Refer and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

- | | | |
|---|---------------|--|
| Atrazine 4L | HARNESS® Xtra | |
| Atrazine 90 WG | SURPASS® 100 | |
| BICEP® II / BICEP® II Lite/ BICEP® II Magnum/ BICEP® II Magnum Lite | | |
| EXTRAZINE® | | |
| FULTIME® | | |
| GUARDSMAN® | | |

100412

**BROADLEAF WEEDS
CONTROLLED BY BALANCE® 4SC ALONE AND IN TANK MIXTURES FOR FIELD CORN**

Tank mixes with BALANCE® 4SC Herbicide are not limited to the tank mix partners mentioned in this table. Refer and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

Broadleaf Weeds (C = Weeds Controlled)	BALANCE® Alone	BALANCE® plus Atrazine	BALANCE® plus BICEP II/ BICEP II Lite/ BICEP II Magnum/ BICEP II Magnum Lite	BALANCE® plus DUAL/ DUAL II/ DUAL II Magnum	BALANCE® plus FRONTIER	BALANCE® plus HARNESS/ SURPASS/ TOPNOTCH	BALANCE® plus HARNESS Xtra/ SURPASS 100/ FULLTIME	BALANCE® plus MICRO- TECH/ PARTNER/ LASSO	BALANCE® plus GUARDSMAN
Amaranth, Palmer	C	C	C	C	C	C	C	C	C
Buffalobur	C	C	C	C	C	C	C	C	C
Burcucumber	C	C	C	C	C	C	C	C	C
Carpetweed				C		C	C	C	C
Chamomile spp.	C	C	C	C	C	C	C	C	C
Chickweed, common	C	C	C	C	C	C	C	C	C
Cocklebur*		C	C				C		C
Dandelion (seedling)	C	C	C	C	C	C	C	C	C
Groundcherry (seedling)		C	C				C		C
Galinsoga	C	C	C	C	C	C	C	C	C
Jimsonweed	C	C	C	C	C	C	C	C	C
Kochia	C	C	C	C	C	C	C	C	C
Lambsquarters, common	C	C	C	C	C	C	C	C	C
Mallow, Venice	C	C	C	C	C	C	C	C	C
Marestail	C	C	C	C	C	C	C	C	C
Morningglory, annual*		C	C				C		C
Wild mustard	C	C	C	C	C	C	C	C	C
Nightshade, black	C	C	C	C	C	C	C	C	C
Nightshade, eastern black	C	C	C	C	C	C	C	C	C
Nightshade, hairy						C	C	C	C
Pennycress, field	C	C	C	C	C	C	C	C	C
Plantain, broadleaf	C	C	C	C	C	C	C	C	C
Pigweed, prostrate	C	C	C	C	C	C	C	C	C
Pigweed, redroot	C	C	C	C	C	C	C	C	C
Pigweed, smooth	C	C	C	C	C	C	C	C	C
Purslane, common	C	C	C	C	C	C	C	C	C
Radish, wild	C	C	C	C	C	C	C	C	C
Ragweed, common	C	C	C	C	C	C	C	C	C
Ragweed, giant*	C	C	C	C	C	C	C	C	C
Shepherds-purse	C	C	C	C	C	C	C	C	C
Smartweed, Pennsylvania	C	C	C	C	C	C	C	C	C
Spurge, toothed	C	C	C	C	C	C	C	C	C
Sunflower, wild*		C	C				C		C
Velvetleaf	C	C	C	C	C	C	C	C	C
Waterhemp, common	C	C	C	C	C	C	C	C	C
Waterhemp, tall	C	C	C	C	C	C	C	C	C

* These weeds may require a postemergence application of BUCTRIL® or other appropriate postemergence herbicides for control of late season escapes.

GRASS WEEDS CONTROLLED BY BALANCE® 4SC ALONE AND IN TANK MIXTURES FOR FIELD CORN

Tank mixes with BALANCE® 4SC Herbicide are not limited to the tank mix partners mentioned in this table. Refer and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

Grassy Weeds (C = Weeds Controlled)	BALANCE® Alone	BALANCE® plus Atrazine	BALANCE® plus BICEP II/ BICEP II Lite/ Magnum/ BICEP II Magnum Lite	BALANCE® plus DUAL/ DUAL II/ Magnum	BALANCE® plus FRONTIER	BALANCE® plus HARNESS/ SURPASS/ TOPNOTCH	BALANCE® plus HARNESS Xtra/ SURPASS 100/ FULLTIME	BALANCE® plus MICRO- TECH/ PARTNER/ LASSO	BALANCE® plus GUARDSMAN
Barnyardgrass	C	C	C	C	C	C	C	C	C
Crabgrass, large	C	C	C	C	C	C	C	C	C
Crabgrass, smooth	C	C	C	C	C	C	C	C	C
Cupgrass, southwestern			C	C	C				C
Cupgrass, woolly *	C	C	C	C	C	C	C	C	C
Foxtail, bristly	C	C	C	C	C	C	C	C	C
Foxtail, giant	C	C	C	C	C	C	C	C	C
Foxtail, green	C	C	C	C	C	C	C	C	C
Foxtail, robust purple	C	C	C	C	C	C	C	C	C
Foxtail, robust white	C	C	C	C	C	C	C	C	C
Foxtail, yellow		C	C	C	C	C	C	C	C
Goosegrass	C	C	C	C	C	C	C	C	C
Johnsongrass, seedling	C	C	C	C	C	C	C	C	C
Panicum, fall	C	C	C	C	C	C	C	C	C
Proso millet, wild*	C	C	C	C	C	C	C	C	C
Rice, red				C		C		C	C
Sandbur, field*	C	C	C	C	C	C	C	C	C
Signalgrass, broadleaf*	C	C	C	C	C	C	C	C	C
Witchgrass			C	C	C	C	C	C	C

* These weeds may require the addition of a pre-emerge grass herbicide tank-mix partner, or an appropriate post-emergence herbicide application for control of late season escapes. BALANCE® Herbicide should be used as part of an integrated control program.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

NET CONTENTS:

BALANCE and BUCTRIL are registered trademarks of Rhône-Poulenc.

Weedone is a registered trademark of NuFarm Americas, Inc.

Axiom is a registered trademark of Bayer Corporation.

Gramoxone, Surpass, Topnotch and Fulltime are registered trademarks of a Zeneca Group Company.

Roundup, Harness, Micro-Tech, Lasso and Partner are registered trademarks of Monsanto Company.

Dual and Magnum are registered trademarks of Novartis.

Frontier and Guardsman is a registered trademark of BASF Corporation.

Prowl is a registered trademark of American Cyanamid Company.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

BALANCE® 4SC Herbicide (PENDING) Resubmitted 7/14/99

